

STRONGID 48- pyrantel tartrate granule
Zoetis Inc.

Strongid® 48

Strongid® 48
(pyrantel tartrate)

Type A Medicated Article

Equine Anthelmintic

Active Drug Ingredient:

Pyrantel tartrate.....10.6% (48 g/lb)

Indications for Use:

For the prevention of *Strongylus vulgaris* larval infestation in horses.

For control of the following parasites in horses:

Large Strongyles (adults): *S. vulgaris*, *S. edentatus*

Small Strongyles (adults and fourth-stage larvae): *Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp., *Poteriostomum* spp., *Triodontophorus* spp.

Pinworms (adults and fourth-stage larvae): *Oxyuris equi*

Ascarids (adults and fourth-stage larvae): *Parascaris equorum*

CAUTION:

CAUTION: For use in manufacturing of equine feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

Storage

STORE IN A DRY, COOL PLACE
SEE BACK PANEL FOR FURTHER USE DIRECTIONS

Net Weight: 50 lb (22.6 kg)

Approved by FDA under NADA # 140-819

51047200

Mixing and Use Directions

Strongid is to be administered on a continuous basis either as a top-dress or mixed in the horse's daily grain ration at the rate of 1.2 mg/lb (2.64 mg/kg) of body weight daily. The duration of administration is for the period during which the horse is at risk of exposure to internal parasites.

Foals may be treated at such time when consistent intake of grain mix is occurring. This is generally between 2 and 3 months of age.

Strongid may be used in mares at any stage of pregnancy or lactation. Stallion fertility is not affected by the use of Strongid.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

Typical Mixing Directions to Deliver 1.2 mg/lb (2.64 mg/kg)

lb of Medicated Grain Mix per 100 lb of Body Weight	lb of Strongid 48	lb of Nonmedicated Feed	Resulting Concentration (g/ton)
2.00	2.5	1997.5	120
1.5	3.3	1996.7	160
0.50	10.0	1990.0	480
0.20	25.0	1975.0	1200

lb of Medicated Top-Dress per 100 lb of Body Weight	lb of Strongid 48	lb of Nonmedicated Feed	Resulting Concentration (g/ton)
0.04	125.0	1875.0	6000
0.025	200.0	1800.0	9600
0.02	250.0	1750.0	12000
0.0125	400.0	1600.0	19200

WARNING

Do not use in horses intended for human consumption.

OTHER WARNINGS

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

CAUTION

Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment, and control of parasitism. Do not mix in feeds containing bentonite.

51047200

Zoetis

Distributed by:
Zoetis Inc.
Kalamazoo, MI 49007
Product of Mexico

PRINCIPAL DISPLAY PANEL - 50lb bag label

Strongid® 48

(pyrantel tartrate)

Type A Medicated Article

Equine Anthelmintic

Active Drug Ingredient

Pyrantel tartrate.....10.6% (48 g/lb)

Indications for Use:

For the prevention of *Strongylus vulgaris* larval infestation in horses.

For control of the following parasites in horses:

Large *Strongylus* (adults): *S. vulgaris*, *S. edentatus*

Small *Strongylus* (adults and fourth-stage larvae): *Cyathostomum* spp., *Cylicocyclus* spp.,

Cylicostephanus spp., *Cylicodontophorus* spp., *Potamostrongylus* spp., *Tricodontophorus* spp.,

Pinworms (adults and fourth-stage larvae): *Oxyuris equi*

Ascarids (adults and fourth-stage larvae): *Parascaris equorum*

CAUTION: For use in manufacturing of equine feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety. Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-erecting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

STORE IN A DRY, COOL PLACE

SEE BACK PANEL FOR FURTHER USE DIRECTIONS

Net Weight: 50 lb (22.6 kg)

Approved by FDA under NADA # 140-819

51047200

zoetis

zoetis

Strongid® 48
(pyrantel tartrate)

STRONGID 48

pyrantel tartrate granule

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:54771-0459
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PYRANTEL TARTRATE (UNII: SC82VF0480) (PYRANTEL - UNII: 4QIH0N49E7)	PYRANTEL TARTRATE	105.8 g in 1 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-0459-1	22.6 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA140819	08/24/2018	

Labeler - Zoetis Inc. (828851555)

Revised: 8/2023

Zoetis Inc.