# BOSLEY MD REGULAR STRENGTH FOR WOMEN- minoxidil 2% solution Scientific Hair Research, LLC

-----

### Bosley MD Regular Strength for Women Minoxidil Topical Solution 2%

Minoxidil 2% w/v

Hair regrowth treatment

#### **USE**

Use to regrow hair on the top of the scalp (vertex only, see pcitre on side of box)

For external use only.

Flamable: Keep away from fire of flame.

Do not use if:

Your amount of hair loss is different than shown on the side of this carton, because this product may not work for you

You have no family history of hair loss

Your hair loss is sudden and/or patchy

Your hair loss is associated with childbirth

You do not know the reason for your hair loss

You are under 18 years of age. Do not use on babies or children

Your scalp is red, inflamed, infected, irritated, or painful

You use other medicines on the scalp

Ask a doctor before use if you have heart disease

When using this product

- Do not apply on other parts of the body
- Avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- Some people have expereinced changes in hair color and/or texture
- It takes time to re-grow hair. You may need to use this product for at least 4 months before you see results.
- The amount of hair regrowth is different for each person. This product will not work for everyone.

Stop use and ask a doctor if

- Chest pain, rapid heartbeat, faintness, or dizziness occurs
- Sudden, unexplained weight gain occurs
- Your hands or feet swell
- Scalp irritation or redness occurs
- Unwanted facial hair growth occurs
- You do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.

#### **DIRECTIONS**

- Apply one mL with dropper 2 times a day directly onto the scalp in the hair loss area
- Using more or more ofthen will not improve results
- Continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again
- See hair loss pictures on the side of this carton
- Before use, read all information on carton and enclosed leaflet
- Keep the carton. It contains important information
- In clincial studies of mostly white women aged 18-45 years with moderate degrees of hair loss, the following responses to Minoxidil Topical Solution 2% was reported: 19% of women reported moderate hair regrowth after using 2% minoxidil topical solution for 8 months (19% had moderate regrowth; 40% had minimal hair regrowth). This compares with the 7% of women reporting moderate hair regrowth after using a placebo, the liquid without minoxidil in it, for 8 months (7% had moderate regrowth; 33% had minimal regrowth).
- Store at controlled room temperature 20 o to 25 oC (68 o to 77 oF)

Alcohol, Propylene Glycol, Purified Water

Questions or comments? Call youre health professional or BosleMD at 1(800)BOSLEY1



#### minoxidil 2% solution

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69113-001

Route of Administration TOPICAL

## **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)	MINOXIDIL	2 g in 100 mL

# Inactive Ingredients Ingredient Name Strength WATER (UNII: 059QF0KO0R) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) ISOPROPYL ALCOHOL (UNII: ND2M416302)

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:69113-001- 01	2 in 1 BOX	05/21/2020				
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product					

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
ANDA	ANDA078176	05/21/2020				

# **Labeler -** Scientific Hair Research, LLC (961897647)

Revised: 4/2024 Scientific Hair Research, LLC