BENZ O STHETIC- benzocaine spray Geritrex LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Benz-O-Sthetic Spray

Drugs Facts

Active Ingredients Purpose Benzocaine 20%......Oral Anesthetic

Uses

Topical anesthetic for the temporary relief of occasional minor irritation and pain, associated with: canker sores

- sore mouth and throat
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances
- minor injury of the mouth and gums

Directions

May be used with extension tube for finer direct application. For adults and children 2 years and older anesthetic use in mouth or throat:

Pump spray 2-3 times up to 4 times. Repeat if needed for larger areas. Anesthetic effect is reached in 15-30 seconds. Children under 12 years of age should be supervised and use as directed by dentist/doctor.

Warnings

Do not use this product if patient has a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. Do not spray into or near eyes. Do not inhale.

Patient: When using extension tube, do not spray in the back of the throat unless your dentist or doctor directed you to do so.

When using this product:

- Avoid use near eyes
- Do not use more than recommended dosage
- Do not go beyond 7 days usage, unless directed by your doctor/dentist

Patient: Discontinue use and consult a physician if: sore mouth symptoms do not improve in 5 days; irritation, pain, or redness persists or worsens; swelling, rash, or fever develops.

Other Ingredients

Alcohol, benzyl alcohol, flavor, peppermint oil, polyethylene glycol, sodium saccharin

Other Information:

store at controlled room temperature 20-25°C (68-77°F)

KEEP OUT OF REACH OF CHILDREN



3.5"

Benz-O-Sthetic®

Topical Anesthetic

Non Aerosol Pump Spray

Fresh Cherry-Mint Flavor

- Fast Acting Spray Tastes Great
- Disposable Applicators
 - Safe and Effective

Benzocaine 20%

METERED DOSE: 40MG PER SPRAY

Net Wt. 2 FL OZ (59ml)

Uses Topical anesthetic for the temporary relief of occasional minor initiation and pain, associated with: ■ cauker sizes ■ sore mouth and threat. ■minor dental procedures ■ minor initiation of the mouth and gums caused by dentures or orthodontic appliances ■ minor injury of the mouth and gums.

Warnings
Do not use this product if patient has a history of allergy to local anesthetics such as procaine, butacaine, berzocaine, or other "caine" anesthetics. Do not spray into or near eyes. Do not inhale Patient: When using extension tube, do not spray in the back of the throat unless your dentist or doctor directed you to do so.

- When using this product Avoid use near eyes
 Do not use more than recommended dosage
 Do not use unless directed by your doctor/dentist
- Patient: Discontinue use and consult a physician if: sore mouth symptoms do not improve within 2 days; irritation, pain, or redness persists or worsens; swelling, rash, or fever develops.

KEEP OUT OF REACH OF CHILDREN. Caution: If accidental or excessive ingestion occurs (exceeds recommended dosage) contact a

physician for immediate advice or call a Poison Control Center. Directions May be used with extension habe for finer direct application. For Adults and children 2 years and older anesthelfic use in mouth or threat pump sing it 2.3 times up to 4 times. Repeat if needed for larger area. Anestheld celfest is cracked in 15-30 seconds. Children under 12 yrs of age should be supervised and use as directed by dentistictors.

Other information

Store at controlled room temperature 20-25°C (68-77°F)

Inactive ingredients alcohol, benzyl alcohol, flavor, peppermint oil, polyethylene glycol, Sodium Saccharin



Dieline Does Not Print

1.5"





PMS 299

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:54162-929

Route of Administration TOPICAL

Active Ingredient/Active Moiety

1	8			
	Ingredient Name		Basis of Strength	Strength
	BENZO CAINE (UNII: U3RS Y48 JW5) (BENZO CAINE - UNII:	U3RS Y48 JW5) BI	ENZOCAINE	11.8 g in 59 mL

Inactive Ingredients			
Ingredient Name	Strength		
BENZYL ALCOHOL (UNII: LKG8494WBH)			
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
PEPPERMINT OIL (UNII: AV092KU4JH)			
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)			

Product Characteristics				
Color		Score		
Shape		Size		
Flavor	CHERRY (CHERRY)	Imprint Code		
Contains				

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:54162-929- 02	59 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	06/30/2017	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part348	06/30/2017		

Labeler - Geritrex LLC (112796248)

Establishment				
Name	Address	ID/FEI	Business Operations	
Geritrex LLC		112796248	manufacture(54162-929)	

Revised: 9/2017 Geritrex LLC