

BOVAPRO 52 CONCENTRATE- iodine spray
AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

BovaPro 52 Concentrate

BOVAPRO 52 CONCENTRATE (1:9)

IODINE

TEAT DIP

WARNING

CAUSES SERIOUS EYE IRRITATION

FOR FARM AND INDUSTRIAL USE ONLY.

KEEP OUT OF REACH OF CHILDREN.

KEEP OUT OF DIRECT SUNLIGHT.

AVOID FREEZING - PRODUCT WILL BECOME THICKER WITH COLD.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS WHEN MIXED:

Active Ingredient - Iodine 0.5 % (5.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution

Emollients: Glycerine 2.0%

BovaPro is made with IoPlex technology and is NPE free. BovaPro is available as a dip, a spray, or in a concentrate.

Use BovaPro effectively to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

Directions:

General Use

IMPORTANT: THIS IS A CONCENTRATE AND MUST BE DILUTED WITH WATER PRIOR TO USE. Mix ONE part concentrate to NINE parts of clean soft water (mix by volume).

Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this

container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

BOVAPRO 52 CONCENTRATE (1:9)

IODINE
TEAT DIP

WARNING



CAUSES SERIOUS EYE IRRITATION

FOR FARM AND INDUSTRIAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
KEEP OUT OF DIRECT SUNLIGHT.

AVOID FREEZING - PRODUCT WILL BECOME
THICKER WITH COLD.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/
face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact
lenses, if present and easy to do. Continue rinsing. If eye irritation persists:
Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal
plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS WHEN MIXED:


Active Ingredient - Iodine 0.5 % (5.0 % Polyethoxy Polypropoxy
Polyethoxy Ethanol - Iodine Complex) Titration via USP33
Reissue, Topical Solution
Emollients: Glycerine 2.0%

BovaPro is made with IoPlex technology and is NPE free. BovaPro is available
as a dip, a spray, or in a concentrate.

NET CONTENTS:



8290C-BULK



Use BovaPro effectively to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

Directions:

General Use

IMPORTANT: THIS IS A CONCENTRATE AND MUST BE DILUTED WITH WATER PRIOR TO USE. Mix ONE part concentrate to NINE parts of clean soft water (mix by volume).

Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.



AgroChem

26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

BOVAPRO 52 CONCENTRATE (1:9)

IODINE
TEAT DIP

WARNING



CAUSES SERIOUS EYE IRRITATION

FOR FARM AND INDUSTRIAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
KEEP OUT OF DIRECT SUNLIGHT.

AVOID FREEZING - PRODUCT WILL BECOME
THICKER WITH COLD.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/
face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact
lenses, if present and easy to do. Continue rinsing. If eye irritation persists:
Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal
plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS WHEN MIXED:

Active Ingredient - Iodine 0.5 % (5.0 % Polyethoxy Polypropoxy
Polyethoxy Ethanol - Iodine Complex) Titration via USP33
Reissue, Topical Solution
Emollients: Glycerine 2.0%

BovaPro is made with IoPlex technology and is NPE free. BovaPro is available
as a dip, a spray, or in a concentrate.

NET CONTENTS:



Use BovaPro effectively to reduce the spread of mastitis causing organisms. Recommended for pre and post
milking applications.

Directions:

General Use

IMPORTANT: THIS IS A CONCENTRATE AND MUST BE DILUTED WITH WATER PRIOR TO USE. Mix ONE part concentrate
to NINE parts of clean soft water (mix by volume).

Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a
veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make
sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.
If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse
or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using
recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats
should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat
full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats
dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product.
Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow
leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last
milking. Dipping should be resumed at least twice daily one week before calving.



AgroChem

26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850

BOVAPRO 52 CONCENTRATE

iodine spray

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:17307-3293
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Iodine (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	Iodine	0.005 kg in 1 kg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	0.35 kg in 1 kg
GLYCERIN (UNII: PDC6A3C0OX)	0.09 kg in 1 kg
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	0.09 kg in 1 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3293-1	19 kg in 1 CONTAINER		
2	NDC:17307-3293-2	57 kg in 1 DRUM		
3	NDC:17307-3293-3	208 kg in 1 DRUM		
4	NDC:17307-3293-4	1040 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/01/2013	

Labeler - AgroChem Inc (805427700)

Establishment

Name	Address	ID/FEI	Business Operations
AgroChem Inc		805427700	api manufacture

Revised: 1/2018

AgroChem Inc