

**HEALTH MART ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride,
simethicone tablet
Strategic Sourcing Services LLC**

McKesson Anti-Diarrheal Anti-Gas Drug Facts

Active ingredients (in each caplet)

Loperamide HCl 2 mg

Simethicone 125 mg

Purposes

Anti-diarrheal

Anti-gas

Uses

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use I you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- **do not use if printed foil under cap is broken or torn**

Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin,

microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Health Mart®

Compare to Imodium® Multi-Symptom Relief active ingredients

MULTI-SYMPTOM RELIEF CAPLET

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg

Anti-Diarrheal

Anti-Gas

Relieves symptoms of diarrhea plus Cramps & Pressure, Bloating, Gas

Gluten Free

ACTUAL SIZE

12 Caplets*

(*Capsule-Shaped Tablets)

NDC 62011-0365-1



Health Mart.

Compare to Imodium® Multi-Symptom Relief active ingredients**

MULTI-SYMPTOM RELIEF CAPLET

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg

Anti-Diarrheal Anti-Gas

Relieves symptoms of diarrhea *plus* Cramps & Pressure, Bloating, Gas

Gluten Free



ACTUAL SIZE

12 Caplets* (*Capsule-Shaped Tablets)

ORF53 W6 C2

Drug Facts

Active ingredients (in each caplet) **Purposes**
 Loperamide HCl 2 mg..... Anti-diarrheal
 Simethicone 125 mg..... Anti-gas

Uses relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- mucus in feces
- a history of liver disease
- a history of an abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- each caplet contains: calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- do not use if printed blister unit is broken or torn

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® Multi-Symptom Relief.

MCKESSON

Another Quality Product
 Distributed by McKesson
 6555 State Highway 161, Irving, TX 75039
 Money Back Guarantee
 healthmart.com

Drug Facts (continued)

Inactive ingredients acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

Questions or comments? 1-800-719-9260



HEALTH MART ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0365
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	LOR1;2;125
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0365-1	12 in 1 CARTON	10/10/2018	

1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209837	10/10/2018	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 11/2022

Strategic Sourcing Services LLC