

NICOTINE POLACRILEX COATED MINT- nicotine polacrilex gum, chewing
Publix Supermarkets, Inc.

Nicotine Polacrilex Gum 2 mg and 4 mg Coated Mint - Publix

Drug Facts

Active ingredient (in each chewing piece) - 2 mg

Nicotine Polacrilex 2 mg (nicotine)

Active ingredient (in each chewing piece) - 4 mg

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may

need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions - 2 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Directions - 4 mg

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information - 2 mg

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Other information - 4 mg

- **each piece contains:** calcium 120 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide and xylitol.

Inactive ingredients - 4 mg

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 aluminum lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide and xylitol.

Principal display panel 2 mg

*Compare to the active ingredient in Nicorette® Gum

SUGAR FREE

nicotine gum

NICOTINE POLACRILEX GUM USP, 2 mg (nicotine

STOP SMOKING AID

COATED MINT FLAVOR

FOR THOSE WHO SMOKE THEIR FIRST CIGERATTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Chew **at least 9 pieces** of Nicotine Polacrilex per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a Support Program as directed in the enclosed User's Guide.
 - Not for sale to those under 18 years of age.
 - Proof of age required.
 - Not for sale in vending machines or from any source where proof of age cannot be verified.

Blister packaged for your protection. Do not use if individual seals are open or torn.

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

DISTRIBUTED BY PUBLIX SUPERMARKETS, INC.,

3300 PUBLIX CORPORATE PARKWAY,

LAKELAND, FL 33811, 1-888-267-3037

Package label 2 mg



SUGAR FREE

nicotinegum

NICOTINE POLACRILEX GUM

USP, 2 mg (nicotine)

STOP SMOKING AID



NDC 58062-428-44

SUGAR FREE

nicotinegum

NICOTINE POLACRILEX GUM

USP, 2 mg (nicotine)

STOP SMOKING AID

COATED
MINT FLAVOR

- **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.**
- **If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg**



ACTUAL SIZE

2 mg 100 PIECES OF GUM
2 mg EACH

*Compare to the active ingredient in Nicorette® Gum

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** – Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** – Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.



To remove the gum, tear off single unit.



Peel off backing starting at corner with loose edge.



Push gum through foil.

flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sac, titanium dioxide, xylitol

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.


■ mouth, throat, or jaw problems occur
■ you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
■ you have symptoms of an allergic reaction (such as difficulty breathing or rash)
Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. When used, pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ If you are under 18 years of age, ask a doctor before use
■ Before using this product, read the enclosed User's Guide for complete directions and other important information
■ Begin using the gum on your quit day
■ If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum

DISTRIBUTED BY PUBLIC SUPER MARKETS, INC., 3300 PUBLIC CORPORATE PARKWAY, LAKELAND, FL 33811, 1-888-267-3037
publix.com
PUBLIC GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK



machines or from any source where proof of age cannot be verified.



PUBLIX Nicotine Polacrilex Gum Coated Mint Flavor

Principal display panel 4 mg

*Compare to the active ingredient in Nicorette® Gum

SUGAR FREE

nicotinegum

NICOTINE POLACRILEX GUM USP, 4 mg (nicotine)

STOP SMOKING AID

COATED MINT FLAVOR

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum USP, 2 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. Use with a Support Program as directed in the enclosed User's Guide.
 - Not for sale to those under 18 years of age.
 - Proof of age required.
 - Not for sale in vending machines or from any source where proof of age cannot be verified.

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette Gum.

Blister packaged for your protection. Do not use if individual seals are open or torn

DISTRIBUTED BY PUBLIX SUPERMARKETS, INC.,

3300 PUBLIX CORPORATE PARKWAY,

LAKELAND, FL 33811, 1888-267-3037

Package label 4 mg



NICOTINE POLACRILEX GUM

USP, 4 mg (nicotine)

STOP SMOKING AID



SUGAR FREE

NICOTINE POLACRILEX GUM

USP, 4 mg (nicotine)

STOP SMOKING AID

COATED
MINT FLAVOR

- **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.**
- **If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum USP, 2 mg**



4 mg 100 PIECES OF GUM
4 mg EACH

*Compare to the active ingredient in Nicorette® Gum

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** – Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** – Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.



DISTRIBUTED BY PUBLIC SUPER MARKETS, INC., 3300 PUBLIC CORPORATE PARKWAY,
LAKELAND, FL 33811, 1-888-267-3037
lakes.com
PUBLIC GUARANTEE: COMPLETE SATISFACTION OR YOUR MONEY BACK


PLD-032A F0007628
4141540173 0

- Not for sale to those under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

Healthcare, LP, distributor of Neogen® Gum.

[illegible]

If you are under 18 years of age, ask a doctor before use.
 Before using the product, read the enclosed User's Guide for complete directions and other important information.
 • Do not use the gum on your gum (it is for oral use only).
 • Do not use the gum more than 30 minutes after waking up, use 2 mg nicotine gum.
 Directions



PUBLIX Nicotine Polacrilex Gum Coated Mint Flavor

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56062-428
Route of Administration	BUCCAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)		NICOTINE	2 mg
Inactive Ingredients			
Ingredient Name			Strength
ACACIA (UNII: 5C5403N26O)			
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)			
CARNAUBA WAX (UNII: R12CBM0EIZ)			
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)			
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)			
SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
SODIUM CARBONATE (UNII: 45P3261C7T)			
TALC (UNII: 7SEV7J4R1U)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
XYLITOL (UNII: VCQ006KQ1E)			
Product Characteristics			
Color	white	Score	no score
Shape	SQUARE	Size	14mm
Flavor	MINT (coated mint)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-428-44	10 in 1 CARTON	01/01/2014	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078699	01/01/2014	

NICOTINE POLACRILEX COATED MINT

nicotine polacrilex gum, chewing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56062-429
Route of Administration	BUCCAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics

Color	white	Score	no score
Shape	SQUARE	Size	14mm

Flavor		MINT (coated mint)	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-429-44	10 in 1 CARTON	01/01/2014	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA078697	01/01/2014	

Labeler - Publix Supermarkets, Inc. (006922009)

Revised: 12/2022

Publix Supermarkets, Inc.