LIQUID ACETAMINOPHEN- acetaminophen liquid LLORENS PHARMACEUTICALS INTERNATIONAL DIVISION

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active ingredient (per 5 mL teaspoonful)

Acetaminophen 160 mg

Purpose

Pain Reliever/Fever Reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 6 doses in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have liver disease.

Ask a doctor before use if you are taking the blood thinning drug warfarin.

When using this product: Do not exceed recommended dose.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children. In case of overdose, get medical help or contact a Paoison Control Center right away. Quick medical attention is critical for adults/children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- find dose in chart below
- if possible, use weight to dose, otherwise use age
- dosage may be repeasted every 4 hours, or as directed by a doctor
- do not give more than 5 doses in 24 hours

| Weight (lbs) | under 24 | 24-35 | 36-47 | 48-59 | 60-71 | 72-95 | over 95 |
|--------------------------------|----------------------|--------|-----------------------|--------|------------------------|------------------|------------------|
| Age (years) | under 2 | 2 to 3 | 4 to 5 | 6 to 8 | 9 to 10 | 11 | 12 to adult |
| Dosage Teaspoonful (tsp) | consult Physician | • | 1 1/2 tsp (7.5 mL) | • | 2 1/2 tsp (12.5 mL) | 3 tsp (15 mL) | 4 tsp (20 mL) |

Inactive ingredients: artificial and natural cherry flavor, citric acid, FD&C Red #40, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate, sucralose

Drug Facts

Active ingredient (per 5mL teaspoonful) Purpose Acetaminophen 160 mg., ...Pain Reliever / Fever Reducer

Uses • temporarily relieves minor aches and pains due to:
• headache • muscular aches • backache • minor pain of arthritis • the common cold • toothache • premenstrual and menstrual cramps • temporarily reduces fever

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if: • adult takes more than 6 doses in 24 hours, which is the maximum daily amount . child takes more than 5 doses in 24 hours, which is the maximum daily amount taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks everyday while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have a liver disease. Ask a doctor before use if you are taking the blood thinning drug warfarin.

When using this product: Do not exceed recomn Stop use and ask a doctor if . pain gets worse or lasts more than 10 days in adults • pain gets worse or lasts more than 5 days in children under 12 years • fever gets worse or lasts more than 3 days . new symptoms occur . redness or swelling is present. These could be signs of a serious condition.

NDC 54859-809-16



LIQUID ACETAMINOPHEN 160mg / 5mL

- ✓ Pain Reliever
- √ Aspirin Free
- √ Fever Reducer
- √ Alcohol Free

√ Compare to the active ingredient of Tylenol® Liquid.*

CHERRY FLAVOR

CONTENT: 16 FL OZ (473 mL)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use, Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away, Quick medical attention is critical for adults/children even if you do not notice any

Directions • do not take more than directed

• find dose in chart below • if possible, use weight to dose, otherwise use age

dosage may be repeated every 4 hours, or as directed by a doctor
 do not give more than 5 doses in 24 hours

| Weight (lbs) | under 24 | 24- 35 | 36- 47 | 48- 59 | 60- 71 | 72- 95 | over 95 |
|--------------------------------|----------------------|-----------------|---------------------|------------------|----------------------|------------------|------------------|
| Age (years) | Under 2 | 2 to 3 | 4 to 5 | 6 to 8 | 9 to 10 | 11 | 12 to adult |
| Dosage Teaspoonful (tsp) | consult Physician | 1 tsp (5 mL) | 1 ½ tsp (7.5 mL) | 2 tsp (10 mL) | 2 ½ tsp (12.5 mL) | 3 tsp (15 mL) | 4 tsp (20 mL) |

Other information

- store at room temperature 15°- 30°C (59°-86°F)
- protect from freezing
 protect from light
 TAMPER-EVIDENT: Do not use if foil seal over bottle opening is

torn, broken or missing. Pharmacist: Preserve and dispense in tight, light-resistant container with a child resistant cap as defined in the USP

Inactive ingredients: artificial and natural cherry flavor, citric acid, FD&C Red #40, methylparaben, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium citrate, sucralose

THIS IS A BULK CONTAINER NOT INTENDED FOR RETAIL

Manufactured by: Llorens Pharmaceutical International Division, Inc. 7080 NW 37th Court. Miami, FL 33147

Lot #:

Exp Date:

Code#: L-60 REV. 11/21



LIQUID ACETAMINOPHEN

acetaminophen liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:54859-809

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) ACETAMINOPHEN 160 mg in 5 mL

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |

METHYLPARABEN (UNII: A2I8C7HI9T) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3MJQ0SDW1A)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3) PROPYLPARABEN (UNII: Z8IX2SC10H)

WATER (UNII: 059QF0KO0R)

SODIUM CITRATE (UNII: 1Q73Q2JULR) **SUCRALOSE** (UNII: 96K6UQ3ZD4)

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date | | | | | | |
|---|--------------------------------------|---|-------------------------|-----------------------|--|--|--|--|--|--|
| 1 | | 473 mL in 1 BOTTLE; Type 0: Not a Combination Product | 10/01/2019 | | | | | | | |
| | | | | | | | | | | |
| | | | | Marketing Information | | | | | | |
| M | larketing | Information | | | | | | | | |
| M | larketing Marketing Category | Information Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | | | | | |
| | Marketing Category | Application Number or Monograph Citation | _ | | | | | | | |

Labeler - LLORENS PHARMACEUTICALS INTERNATIONAL DIVISION (037342305)

Registrant - LLORENS PHARMACEUTICALS INTERNATIONAL DIVISION (037342305)

Revised: 7/2022 LLORENS PHARMACEUTICALS INTERNATIONAL DIVISION