GABAPENTIN- gabapentin capsule GABAPENTIN- gabapentin tablet, film coated GABAPENTIN- gabapentin tablet DIRECT RX

GABAPENTIN

Gabapentin (gab" a pen' tin)Capsules

What is the most important information I should know about gabapentin capsules?

Do not stop taking gabapentin capsules without first talking to your healthcare provider.

Stopping gabapentin capsules suddenly can cause serious problems.

Gabapentin capsules can cause serious side effects including:

1. Suicidal Thoughts. Like other antiepileptic drugs, gabapentin capsules may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

Call a healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- \cdot thoughts about suicide or dying
- · attempts to commit suicide
- new or worse depression
- new or worse anxiety
- \cdot feeling agitated or restless
- · panic attacks
- · trouble sleeping (insomnia)
- · new or worse irritability
- · acting aggressive, being angry, or violent
- \cdot acting on dangerous impulses
- \cdot an extreme increase in activity and talking (mania)
- \cdot other unusual changes in behavior or mood

How can I watch for early symptoms of suicidal thoughts and actions?

 \cdot Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.

 \cdot Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

Do not stop taking gabapentin capsules without first talking to a healthcare provider.

· Stopping gabapentin capsules suddenly can cause serious problems. Stopping a

seizure medicine suddenly in a patient who has epilepsy can cause seizures that will not stop (status epilepticus).

 \cdot Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

2. Changes in behavior and thinking -Using gabapentin capsules in children 3 to 12 years of age can cause emotional changes, aggressive behavior, problems with concentration, restlessness, changes in school performance, and hyperactivity.

3. Gabapentin capsules may cause a serious or life-threatening allergic reaction that may affect your skin or other parts of your body such as your liver or blood cells. This may cause you to be hospitalized or to stop gabapentin capsule. You may or may not have a rash with an allergic reaction caused by gabapentin capsule. Call a healthcare provider right away if you have any of

The following symptoms:

- · skin rash
- hives
- · difficulty breathing
- · fever
- \cdot swollen glands that do not go away
- \cdot swelling of your face, lips, throat or tongue
- \cdot yellowing of your skin or of the whites of the eyes
- · unusual bruising or bleeding
- · severe fatigue or weakness
- \cdot unexpected muscle pain
- \cdot frequent infections

These symptoms may be the first signs of a serious reaction. A healthcare provider should examine you to decide if you should continue taking gabapentin capsules.

What are gabapentin capsules?

Gabapentin capsules are a prescription medicine used to treat:

 \cdot Pain from damaged nerves (postherpetic pain) that follows healing of shingles (a painful rash that comes after a herpes zoster infection) in adults.

 \cdot Partial seizures when taken together with other medicines in adults and children 3 years of age and older with seizures.

Who should not take gabapentin capsules?

Do not take gabapentin capsules if you are allergic to gabapentin or any of the other ingredients in gabapentin capsules. See the end of this Medication Guide for a complete list of ingredients in gabapentin capsules.

What should I tell my healthcare provider before taking gabapentin capsules?

Before taking gabapentin capsules, tell your healthcare provider if you:

- \cdot have or have had kidney problems or are on hemodialysis
- \cdot have or have had depression, mood problems, or suicidal thoughts or behavior
- have diabetes

 \cdot are pregnant or plan to become pregnant. It is not known if gabapentin capsules can harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking gabapentin capsules. You and your healthcare provider will decide if you should take gabapentin capsules while you are pregnant.

o Pregnancy Registry: If you become pregnant while taking gabapentin capsules, talk to your healthcare provider about registering with the North American Antiepileptic Drug (NAAED) Pregnancy Registry. The purpose of this registry is to collect information about the safety of antiepileptic drugs during pregnancy. You can enroll in this registry by calling 1-888-233-2334.

 \cdot are breast-feeding or plan to breast-feed. Gabapentin capsules can pass into breast milk. You and your healthcare provider should decide how you will feed your baby while you take gabapentin capsules.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking gabapentin capsules with certain other medicines can cause side effects or affect how well they work. Do not start or stop other medicines without talking to your healthcare provider.

Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take gabapentin capsules?

 \cdot Take gabapentin capsules exactly as prescribed. Your healthcare provider will tell you how much gabapentin capsules to take.

o Do not change your dose of gabapentin capsules without talking to your healthcare provider.

o Take gabapentin capsules with water.

 Gabapentin capsules can be taken with or without food. If you take an antacid containing aluminum and magnesium, such as Maalox®, Mylanta®, Gelusil®, Gaviscon®, or Di-Gel®, you should wait at least 2 hours before taking your next dose of gabapentin capsules.

If you take too much gabapentin capsules, call your healthcare provider or your local Poison Control Center right away at 1-800-222-1222.

What should I avoid while taking gabapentin capsules?

• Do not drink alcohol or take other medicines that make you sleepy or dizzy while taking gabapentin capsules without first talking with your healthcare provider. Taking gabapentin capsules with alcohol or drugs that cause sleepiness or dizziness may make your sleepiness or dizziness worse.

 \cdot Do not drive, operate heavy machinery, or do other dangerous activities until you know

how gabapentin capsules affects you. Gabapentin capsules can slow your thinking and motor skills.

What are the possible side effects of gabapentin capsules?

Gabapentin capsules may cause serious side effects including:

See "What is the most important information I should know about gabapentin capsules?"

 \cdot problems driving while using gabapentin capsules. See "What I should avoid while taking gabapentin capsules?"

 \cdot sleepiness and dizziness which could increase the occurrence of accidental injury, including falls

 \cdot The most common side effects of gabapentin capsules include:

 \cdot lack of coordination \bullet feeling tired

 \cdot viral infection • fever

- · feeling drowsy jerky movements
- \cdot nausea and vomiting \bullet difficulty with coordination
- \cdot difficulty with speaking \bullet double vision
- · tremor unusual eye movement
- \cdot swelling, usually of legs and feet

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of gabapentin capsules. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to Time Cap Laboratories 1 877 290-4008 and/or FDA at 1-800-FDA-1088.

How should I store gabapentin capsules?

· Store gabapentin capsules between 68°F to 77°F (20°C to 25°C).

Keep gabapentin capsules and all medicines out of the reach of children.

General information about the safe and effective use of gabapentin capsules

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use gabapentin capsules for a condition for which it was not prescribed. Do not give gabapentin capsules to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about gabapentin capsules. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about gabapentin capsules that was written for healthcare professionals.

What are the ingredients in gabapentin capsules?

Active ingredient: gabapentin, USP

Inactive ingredients in the capsules: lactose monohydrate, corn starch, and talc.

The 100 mg capsule shell contains gelatin and titanium dioxide.

The 300 mg capsule shell contains gelatin, titanium dioxide, and yellow iron oxide.

The 400 mg capsule shell contains gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. The imprinting ink contains FD&C Blue No. 2, propylene glycol and shellac.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Rx only

Manufactured for:

Time-Cap-Labs, INC,

7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by:

Marksans Pharma Ltd.

Plot No. L-82, L-83, Verna Indl. Estate,

Verna, Goa-403 722, India

Rev. 11/17

100 mg Capsules (White/White colored, size '3' hard gelatin capsules with "103" printed on body of capsules containing white to off white granular powder) Bottles of 100 & 500

degrees.	NDC 72189-158-30 Gabapentin	5 5 100 mpc 10 000 mg 5 5 100 721 89 - 158 - 30 30 Tebs 5 10 10 721 89 - 158 - 30 30 Tebs 5 10 10 10 10 10 10 10 10 10 10 10 10 10
tr of this drug to any corn it was prescribed, storeen 58-77 degrees lidreen.	800mg 30 Tabs	Gatemperson Bolding MBC 72189-158-30 30 Telm Let 31AU2206 Exp 2/26/23 Mbg NOC 71093-112-64
w prohibits tran the patient for w spe insert. Store out of reach of c	Generic For: Neurontin Each film-coated tablet contains: Gabapentin, USP800 mg	State Endemperature 000mmg NDE 72189 - 158 - 30 30 Taks NDE 72189 - 158 - 30 30 Taks Lot 31AU2208 Exp 2/28/23 Hitg NDC 71883 - 112 - 04
A TRUE STORE	Lets 31AU2208 Prode 4193 - 800 - 30 ackaged and DIRECK STATUS - 156 - 30 31AU2208 Browsorville Browsorville Browsorville Browsorville Browsorville	2 4 4 5 5 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7





Discard After: 10/31/21 72189-189-60

Dawsonville,

GA 30534

10JY2004

10/31/21

ASKJR

Lot# 10JY2084

Packaged and Distributed By:

202

Prod# 72189-0109-60

DIRECT

at 10JY2004 Exp 10/31/21 Mtg NDC 49483-605-50

NDC 72189-109-60 60 Caps

Lot 10JY2004 Exp 10/31/21 Mtg NDC 49483 - 605 - 50

pabapentin 100mg













Lot# SAMPLE

Packaged and

Distributed By:

Prod# 4193-400-30

DIRECT

Discerd After: 1/31/22

Dawsonville,

GA 30534

72189-113-30

SAMPLE

1/31/22

AGALM

SE Mfg NDC 49483-607-50



GABAPENTIN					
gabapentin capsule					
Product Information					
Product Information					
Product Type	HUMAN PRESCE DRUG	RIPTION Item Code (Source)	NDC:72 605)	2189-109(NDC:49483-
Route of Administration	ORAL				
Active Ingredient/Act	ive Moiety				
In	gredient Name		Basis of St	rength	Strength
GABAPENTIN (UNII: 6CW7F3G	59X) (GABAPENTIN - U	UNII:6CW7F3G59X)	GABAPENTIN		100 mg
Inactive Ingredients					
	Ingredient	Name		St	rength
TALC (UNII: 7SEV7J4R1U)					
STARCH, CORN (UNII: 08232					
TITANIUM DIOXIDE (UNII: 15 GELATIN (UNII: 2G86QN327L)	-				
PROPYLENE GLYCOL (UNII: 6					
SHELLAC (UNII: 46N107B710)					
FD&C BLUE NO. 2 (UNII: L06					
Product Characterist	ics				
Color	white	Score		no score	
	CAPSULE	Size		16mm	
Shape	CIN SOLL				
Shape Flavor		Imprint Code		103	

Packaging

- '	ackaging			
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72189-109- 90	90 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2020	
2	NDC:72189-109- 30	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2020	
3	NDC:72189-109- 72	120 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2020	
4	NDC:72189-109- 60	60 in 1 BOTTLE; Type 0: Not a Combination Product	06/12/2020	

Marketing Information

Citation	Marketing Start Date	Marketing End Date
4090007	06/12/2020	
40		

GABAPENTIN

gabapentin capsule

Product Information Item Code HUMAN PRESCRIPTION NDC:72189-113(NDC:49483-**Product Type** DRUG (Source) 607) **Route of Administration** ORAL **Active Ingredient/Active Moiety Ingredient Name Basis of Strength** Strength GABAPENTIN (UNII: 6CW7F3G59X) (GABAPENTIN - UNII:6CW7F3G59X) GABAPENTIN 400 mg **Inactive Ingredients Ingredient Name** Strength TALC (UNII: 7SEV7J4R1U) STARCH, CORN (UNII: 08232NY3SJ) LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) GELATIN (UNII: 2G86QN327L) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) FERRIC OXIDE YELLOW (UNII: EX43802MRT) **TITANIUM DIOXIDE** (UNII: 15FIX9V2JP)

FD&C BLUE NO. 2 (UNII: L06K8R7DQK)

SHELLAC (UNII: 46N107B710)

Product Characteris	tics		
Color	orange	Score	no score

Sł	nape	CAPS	ULE	Size		21mm
Fla	avor			Imprint Code	•	105
Contains						
Pa	ackaging					
#	ltem Code	Pac	kage Descrip	tion	Marketing Start Date	Marketing End Date
1	NDC:72189-113- 90	90 in 1 BOTTLI Product	E; Type 0: Not a C	Combination	10/15/2020	
2	NDC:72189-113- 72	120 in 1 BOTT Product	E; Type 0: Not a	Combination	10/15/2020	
3	NDC:72189-113- 30	30 in 1 BOTTLE Product	E; Type 0: Not a C	Combination	10/15/2020	
4	NDC:72189-113- 60	60 in 1 BOTTLE Product	E; Type 0: Not a C	Combination	10/15/2020	
M	larketing	Informati	ion			
	–				Ma uka tin n Cha ut	Ma ulua tina u Fuad
	Marketing Category	Аррисат	ion Number of Citation	r Monograph	Marketing Start Date	Marketing End Date
		ANDA090007	1		10/15/2020	

GABAPENTIN gabapentin tablet, film coated	4				
Product Information					
Product Type	HUMAN PRESCRIPTION DRUG	ltem Code (Source)		NDC:72189-158 112)	(NDC:71093-
Route of Administration	ORAL				
Active Ingredient/Active	Moiety				
Ingre	dient Name		Basis	of Strength	Strength
GABAPENTIN (UNII: 6CW7F3G59X)	(GABAPENTIN - UNII:6CW7F3	G59X)	GABAPE	NTIN	800 mg
Inactive Ingredients					
indetive ingredients	Ingredient Name				Strength
HYDROXYPROPYL CELLULOSE,		5N6OH)			
TALC (UNII: 7SEV7J4R1U)					
POLYETHYLENE GLYCOL, UNSP	ECIFIED (UNII: 3WJQ0SDW1A))			
HYDROXYPROPYL CELLULOSE,	LOW SUBSTITUTED (UNII: 2	2165RE0K14)			
COPOVIDONE K25-31 (UNII: D9C	330MD8B)				
MAGNESIUM STEARATE (UNII: 70					
MAGNESIUM STEARATE (UNII: /(0097M6I30)				
POLOXAMER 407 (UNII: TUF2IVWS	·				
· · ·	·				

P	roduct Chara	acteristics					
	olor		hite	Score			2 pieces
	hape		VAL	Size			10mm
	avor			Imprint Code			NT;151
Co	ontains						
Pa	ackaging						
#	ltem Code	Pa	ckage Desc	ription	Marketiı Da	ng Start Ite	Marketing End Date
1	NDC:72189-158- 90	90 in 1 BOTTL Product	E; Type 0: Not	a Combination	12/01/2020		
	30						
2	NDC:72189-158- 72	120 in 1 BOTT Product	TLE; Type 0: No	ot a Combination	12/01/2020		
2 3	NDC:72189-158-	Product			12/01/2020 12/01/2020		
_	NDC:72189-158- 72 NDC:72189-158-	Product 30 in 1 BOTTL Product	E; Type 0: Not	a Combination	12/01/2020		
3	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158-	Product 30 in 1 BOTTL Product 60 in 1 BOTTL	E; Type 0: Not	a Combination	12/01/2020		
3	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product	E; Type 0: Not E; Type 0: Not	a Combination	12/01/2020		
3	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product	E; Type 0: Not E; Type 0: Not ion	a Combination a Combination r or Monogra	12/01/2020 12/01/2020 12/01/2020	ting Start	-
3 4 M	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product	E; Type 0: Not E; Type 0: Not ion tion Numbe Citati	a Combination a Combination r or Monogra	12/01/2020 12/01/2020 12/01/2020	Date	Marketing Enc Date
3 4 M	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60 Iarketing Marketing Category	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product Informat Applica	E; Type 0: Not E; Type 0: Not ion tion Numbe Citati	a Combination a Combination r or Monogra	12/01/2020 12/01/2020 12/01/2020 ph Marke	Date	-
3 4 M	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60 Iarketing Marketing Category IDA	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product Applica ANDA20324	E; Type 0: Not E; Type 0: Not ion tion Numbe Citati	a Combination a Combination r or Monogra	12/01/2020 12/01/2020 12/01/2020 ph Marke	Date	-
3 4 M AN	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60 Iarketing Category NDA ABAPENTI	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product Informat Applica ANDA20324	E; Type 0: Not E; Type 0: Not ion tion Numbe Citati	a Combination a Combination r or Monogra	12/01/2020 12/01/2020 12/01/2020 ph Marke	Date	-
3 4 M AN	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60 Iarketing Marketing Category IDA	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product Informat Applica ANDA20324	E; Type 0: Not E; Type 0: Not ion tion Numbe Citati	a Combination a Combination r or Monogra	12/01/2020 12/01/2020 12/01/2020 ph Marke	Date	-
3 4 M	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60 Iarketing Category NDA ABAPENTI	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product Informat Applica ANDA20324 t	E; Type 0: Not E; Type 0: Not ion tion Numbe Citati	a Combination a Combination r or Monogra	12/01/2020 12/01/2020 12/01/2020 ph Marke	Date	-
3 4 M AN G	NDC:72189-158- 72 NDC:72189-158- 30 NDC:72189-158- 60 Iarketing Category IDA ABAPENTI bapentin table	Product 30 in 1 BOTTL Product 60 in 1 BOTTL Product Informat Applica ANDA20324 t	E; Type 0: Not E; Type 0: Not ion tion Numbe Citati	a Combination	12/01/2020 12/01/2020 12/01/2020 ph Marke	Date 20	-

Active ingredient/Active Molecy							
Ingredient Name	Basis of Strength	Strength					
GABAPENTIN (UNII: 6CW7F3G59X) (GABAPENTIN - UNII:6CW7F3G59X)	GABAPENTIN	800 mg					

Inactive Ingredients					
Ingredient Name	Strength				
POLOXAMER 407 (UNII: TUF2IVW3M2)					
MANNITOL (UNII: 30WL53L36A)					
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)					

HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
CROSPOVIDONE (UNII: 2S7830E561)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
COPOVIDONE K25-31 (UNII: D9C330MD8B)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	2 pieces
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	SG;178
Contains			

Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72189-178- 90	90 in 1 BOTTLE; Type 0: Not a Combination Product	02/08/2021	
2	NDC:72189-178- 72	120 in 1 BOTTLE; Type 0: Not a Combination Product	02/08/2021	
3	NDC:72189-178- 30	30 in 1 BOTTLE; Type 0: Not a Combination Product	02/08/2021	

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date
ANDA	ANDA205101	02/08/2021	

Labeler - DIRECT RX (079254320)

Registrant - DIRECT RX (079254320)

Establishment							
Name	Address	ID/FEI	Business Operations				
DIRECT RX		079254320	repack(72189-113, 72189-158, 72189-178, 72189-109)				

Revised: 6/2023

DIRECT RX