


CORDRAN- flurandrenolide tape
Allergan, Inc.

Cordran[®] Tape
Flurandrenolide Tape, USP
Rx only

DESCRIPTION

Cordran Tape (Flurandrenolide Tape, USP) is a transparent, inconspicuous, plastic surgical tape. It contains Cordran (Flurandrenolide, USP), a potent corticosteroid for topical use. Flurandrenolide occurs as white to off-white, fluffy crystalline powder and is odorless. Flurandrenolide is practically insoluble in water and in ether. One gram dissolves in 72 mL of alcohol and in 10 mL of chloroform. The molecular weight of flurandrenolide is 436.52.

The chemical name of flurandrenolide is Pregn-4-ene-3,20-dione, 6-fluoro-11,21 dihydroxy-16,17-[(1-methylethylidene)bis(oxy)]-, (6 α , 11 β , 16 α)-; its empirical formula is C₂₄H₃₃FO₆. The structural formula is as follows:



Each square centimeter contains 4 mcg (0.00916 μ mol) flurandrenolide uniformly distributed in the adhesive layer. The tape is made of a thin, matte-finish polyethylene film that is slightly elastic and highly flexible.

The adhesive is a synthetic copolymer of acrylate ester and acrylic acid that is free from substances of plant origin. The pressure-sensitive adhesive surface is covered with a protective paper liner to permit handling and trimming before application.

CLINICAL PHARMACOLOGY

Cordran is primarily effective because of its anti-inflammatory, antipruritic, and vasoconstrictive actions.

The mechanism of the anti-inflammatory effect of topical corticosteroids is not completely understood. Various laboratory methods, including vasoconstrictor assays, are used to compare and predict potencies and/or clinical efficacies of the topical corticosteroids. There is some evidence to suggest that a recognizable correlation exists between vasoconstrictor potency and therapeutic efficacy in man. Corticosteroids with anti-inflammatory activity may stabilize cellular and lysosomal membranes. There is also the suggestion that the effect on the membranes of lysosomes prevents the release of proteolytic enzymes and, thus, plays a part in reducing inflammation.

The tape serves as both a vehicle and an occlusive dressing. Retention of insensible perspiration by the tape results in hydration of the stratum corneum and improved diffusion of the medication. The skin is protected from scratching, rubbing, desiccation, and chemical irritation. The tape acts as a mechanical splint to fissured skin. Since it prevents removal of the medication by washing or the rubbing action of clothing, the tape formulation provides a sustained action.

Pharmacokinetics—The extent of percutaneous absorption of topical corticosteroids is determined by many factors, including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings.

Topical corticosteroids can be absorbed from normal intact skin. Inflammation and/or other disease processes in the skin increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. Thus, occlusive dressings may be a valuable therapeutic adjunct for treatment of resistant dermatoses (*see* **DOSAGE AND ADMINISTRATION**).

Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to those of systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. They are metabolized primarily in the liver and then excreted in the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS AND USAGE

For relief of the inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses, particularly dry, scaling localized lesions.

CONTRAINDICATIONS

Topical corticosteroids are contraindicated in patients with a history of hypersensitivity to any of the components of these preparations.

Use of Cordran Tape is not recommended for lesions exuding serum or in intertriginous areas.

PRECAUTIONS

General -

Systemic absorption of topical corticosteroids has produced reversible hypothalamic-pituitary-adrenal (HPA) axis suppression, manifestations of Cushing's syndrome, hyperglycemia, and glucosuria in some patients.

Conditions that augment systemic absorption include application of the more potent steroids, use over large surface areas, prolonged use, and the addition of occlusive dressings.

Therefore, patients receiving a large dose of a potent topical steroid applied to a large surface area or under an occlusive dressing should be evaluated periodically for evidence of HPA axis suppression by using urinary-free cortisol and ACTH stimulation tests. If HPA axis suppression is noted, an attempt should be made to withdraw the drug, to reduce the frequency of application, or to substitute a less potent steroid.

Recovery of HPA axis function is generally prompt and complete on discontinuation of the drug. Infrequently, signs and symptoms of steroid withdrawal may occur, so that supplemental systemic corticosteroids are required.

Pediatric patients may absorb proportionately larger amounts of topical corticosteroids and thus be more susceptible to systemic toxicity (*see* ***Pediatric Use*** under **PRECAUTIONS**).

If irritation develops, topical corticosteroids should be discontinued and appropriate therapy instituted.

In the presence of dermatologic infections, the use of an appropriate antifungal or antibacterial agent

should be instituted. If a favorable response does not occur promptly, Cordran should be discontinued until the infection has been adequately controlled.

Information for the Patient -

Patients using topical corticosteroids should receive the following information and instructions:

- This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes.
- Patients should be advised not to use this medication for any disorder other than that for which it was prescribed.
- The treated skin area should not be bandaged or otherwise covered or wrapped in order to be occlusive unless the patient is directed to do so by the physician.
- Patients should report any signs of local adverse reactions, especially under occlusive dressing.
- Parents of pediatric patients should be advised not to use tight-fitting diapers or plastic pants on a patient being treated in the diaper area, because these garments may constitute occlusive dressings.

Laboratory Tests -

The following tests may be helpful in evaluating the HPA axis suppression:

Urinary-free cortisol test

ACTH stimulation test

Carcinogenesis, Mutagenesis, and Impairment of Fertility -

Long-term animal studies have not been performed to evaluate the carcinogenic potential or the effect on fertility of topical corticosteroids.

Studies to determine mutagenicity with prednisolone and hydrocortisone have revealed negative results.

Usage in Pregnancy -

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. The more potent corticosteroids have been shown to be teratogenic after dermal application in laboratory animals. There are no adequate and well-controlled studies in pregnant women on teratogenic effects from topically applied corticosteroids. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Drugs of this class should not be used extensively for pregnant patients or in large amounts or for prolonged periods of time.

Nursing Mothers -

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities *not* likely to have a deleterious effect on the infant. Nevertheless, caution should be exercised when topical corticosteroids are administered to a nursing woman.

Pediatric Use -

Pediatric patients may demonstrate greater susceptibility to topical-corticosteroid-induced HPA axis suppression and Cushing's syndrome than do mature patients because of a larger skin surface area to body weight ratio.

Hypothalamic-pituitary-adrenal (HPA) axis suppression, Cushing's syndrome, and intracranial hypertension have been reported in pediatric patients receiving topical corticosteroids. Manifestations of adrenal suppression in pediatric patients include linear growth retardation, delayed weight gain, low plasma-cortisol levels, and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include bulging fontanelles, headaches, and bilateral papilledema.

Administration of topical corticosteroids to pediatric patients should be limited to the least amount compatible with an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of pediatric patients.

ADVERSE REACTIONS

The following local adverse reactions are reported infrequently with topical corticosteroids but may occur more frequently with the use of occlusive dressings. These reactions are listed in an approximate decreasing order of occurrence: burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis.

The following may occur more frequently with occlusive dressings: maceration of the skin, secondary infection, skin atrophy, striae, miliaria.

OVERDOSAGE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (see **PRECAUTIONS**).

DOSAGE AND ADMINISTRATION

Occlusive dressings may be used for the management of psoriasis or recalcitrant conditions.

If an infection develops, the use of Cordran Tape and other occlusive dressings should be discontinued and appropriate antimicrobial therapy instituted.

Replacement of the tape every 12 hours produces the lowest incidence of adverse reactions, but it may be left in place for 24 hours if it is well tolerated and adheres satisfactorily. When necessary, the tape may be used at night only and removed during the day.

If ends of the tape loosen prematurely, they may be trimmed off and replaced with fresh tape.

The directions given below are included for the patient to follow unless otherwise instructed by the physician.

APPLICATION OF CORDRAN TAPE

IMPORTANT: Skin should be clean and <i>dry</i> before tape is applied. Tape should always be cut, never torn.
--

DIRECTIONS FOR USE:

- Prepare skin as directed by your physician or as follows: Gently clean the area to be covered to remove scales, crusts, dried exudates, and any previously used ointments or creams. A germicidal soap or cleanser should be used to prevent the development of odor under the tape. Shave or clip the hair in the treatment area to allow both good contact with the skin and comfortable removal. If shower or tub bath is to be taken, it should be completed before the tape is applied. The skin should be dry before application of the tape.
- Remove tape from flip-top vial and cut a piece slightly larger than area to be covered. Round off corners.
- Return remainder of tape roll to vial and tightly close the lid.
- Pull white paper from transparent tape. Be careful that tape does not stick to itself.
- Apply tape, keeping skin smooth; press tape into place.

Keep Cordran Tape in the original container. Do not repackage. Protect from moisture. Keep vial tightly

closed in a dry place.

REPLACEMENT OF TAPE:

Unless instructed otherwise by your physician, replace tape after 12 hours. Cleanse skin and allow it to dry for 1 hour before applying new tape.

IF IRRITATION OR INFECTION DEVELOPS, REMOVE TAPE AND CONSULT PHYSICIAN.

HOW SUPPLIED

Tape:

4 mcg/sq cm—small roll, 24 in x 3 in (60 cm x 7.5 cm), in a flip-top vial

NDC 0023-5870-24

4 mcg/sq cm—large roll, 80 in x 3 in (200 cm x 7.5 cm), in a flip-top vial

NDC 0023-5870-80

Store at 20-25°C (68-77°F). [See USP controlled room temperature.]

Keep out of reach of children.

Rx only

REFERENCES

Bard JW: Flurandrenolide tape in the treatment of lichen simplex chronicus. *J Ky Med Assoc* 1969;67:668.

Baxter DL, Stoughton RB: Mitotic index of psoriatic lesions treated with anthralin, glucocorticosteroid and occlusion only. *J Invest Dermatol* 1970;54:410.

Compilation of clinical reports on Cordran Tape received by Eli Lilly and Company.

Halprin KM, Fukui K, Ohkawara A: Flurandrenolone (Cordran) tape and carbohydrate metabolizing enzymes. *Arch Dermatol* 1969;100:336.

Labow TA, Eisert J, Sanders SL: Flurandrenolide tape in treatment of psoriasis. *NY State J Med* 1969;69:3138.

Ronchese F: Flurandrenolone tape therapy. *RI Med J* 1969;52:389.

Sellers FM: Investigative study of flurandrenolone tape in a series of ambulatory outpatients. *J Indiana State Med Assoc* 1970;63:34.

Weiner MA: Flurandrenolone tape, a new preparation for occlusive therapy, *J Invest Dermatol* 1966; 47:63.

Distributed by:

Allergan USA, Inc.

Madison, NJ 07940

Manufactured by:

3M Company

St. Paul, MN 55144 USA

© 2018 Allergan. All rights reserved.

Cordran[®] is a registered trademark of Aqua Pharmaceuticals, LLC.

Allergan[®] and its design are trademarks of Allergan, Inc.

Content Updated: May 2018

v1.0USPI5870

For the Patient

Cordran® Tape ***Flurandrenolide Tape, USP***


DIRECTIONS FOR USE

APPLICATION OF MEDICATED TRANSPARENT TAPE

Rx only

IMPORTANT: Skin should be clean and *dry* before tape is applied. Tape should always be cut, never torn.

- Prepare skin as directed by your physician or as follows: Gently clean the area to be covered to remove scales, crusts, dried exudates, and any previously used ointments or creams. A germicidal soap or cleanser should be used to prevent the development of odor under the tape. Shave or clip the hair in the treatment area to allow both good contact with the skin and comfortable removal. If a shower or tub bath is to be taken, it should be completed before the tape is applied. The skin should be dry before application of the tape.
- Remove tape from flip-top vial and cut a piece slightly larger than area to be covered. Round off corners.



- Return remainder of tape roll to vial and tightly close the lid.



- Pull white paper backing from transparent tape. Be careful that tape does not stick to itself.



- Apply tape, keeping skin smooth; press tape into place.




REPLACEMENT OF TAPE

Unless instructed otherwise by your physician, replace tape after 12 hours. Cleanse skin and allow it to dry for 1 hour before applying new tape.

DISPENSING THE TAPE

To use the roll as a dispenser, pull the tape as illustrated.



IF IRRITATION OR INFECTION DEVELOPS, REMOVE TAPE AND CONSULT PHYSICIAN.

Store at 20 - 25°C (68 - 77°F). [See USP controlled room temperature.]

Keep Cordran Tape in the original container. Do not repackage. Protect from moisture. Keep vial tightly closed in a dry place.

Keep out of reach of children.

Rx only

Distributed by:
Allergan USA, Inc.
Madison, NJ 07940

Manufactured by:
3M Company
St. Paul, MN 55144 USA

© 2018 Allergan. All rights reserved.

Cordran® is a registered trademark of Aqua Pharmaceuticals, LLC.

Allergan® and its design are trademarks of Allergan, Inc.

Content Updated: May 2018

v1.0PPI5870

PRINCIPAL DISPLAY PANEL

Cordran® Tape 4 mcg/sq cm
24 inches x 3 inches (60 cm x 7.5 cm)
NDC 0023-5870-24



PRINCIPAL DISPLAY PANEL

Cordran® Tape 4 mcg/sq cm

80 inches x 3 inches (200 cm x 7.5 cm)
NDC 0023-5870-80

NDC 0023-5870-80

Cordran[®] Tape

Flurandrenolide Tape, USP

4 mcg per sq cm

LARGE ROLL
80 in x 3 in (200 cm x 7.5 cm)

Rx only

USUAL DOSAGE:
Apply to affected area every 12 hours. See accompanying literature.
Keep out of reach of children.
Each sq cm of CORDRAN[®] TAPE contains 4 mcg of flurandrenolide dispersed in the adhesive layer.
Store at 20-25°C (68-77°F). [See USP controlled room temperature.]
Keep Cordran Tape in the original container. Do not repackage. Protect from moisture. Keep vial tightly closed in a dry place.

Distributed by:
Allergan USA, Inc.
Irvine, CA 92612

© 2016 Allergan. All rights reserved.
Cordran[®] is a registered trademark of Aqua Pharmaceuticals, LLC.

LOT:
EXP:



55503US10

PRINCIPAL DISPLAY PANEL

Cordran[®] Tape 4 mcg/sq cm
80 inches x 3 inches (200 cm x 7.5 cm)
NDC 0023-5870-80

4 mcg per sq cm

Cordran[®] Tape

Flurandrenolide Tape, USP

NDC 0023-5870-80 80 Inch



65965US10

USUAL DOSAGE:
Apply to affected area every 12 hours. See accompanying literature.
Keep out of reach of children.
Each sq cm of CORDRAN[®] TAPE contains 4 mcg of flurandrenolide dispersed in the adhesive layer.
Store at 20-25°C (68-77°F). [See USP controlled room temperature.]
Keep Cordran Tape in the original container. Do not repackage. Protect from moisture. Keep vial tightly closed in a dry place.

Cordran[®] Tape

Flurandrenolide Tape, USP

4 mcg per sq cm

LARGE ROLL
80 in x 3 in (200 cm x 7.5 cm)

Rx only

Distributed by:
Allergan USA, Inc.
Irvine, CA 92612

© 2016 Allergan. All rights reserved.
Cordran[®] is a registered trademark of Aqua Pharmaceuticals, LLC.



N 3 00235 87080 6

Full Prescribing Information With Attached Patient Information Enclosed.

CORDRAN
flurandrenolide tape

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0023-5870
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FLURANDRENOLIDE (UNII: 8EUL29XUQT) (FLURANDRENOLIDE - UNII:8EUL29XUQT)	FLURANDRENOLIDE	4 ug in 1 cm2

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0023-5870-24	450 cm2 in 1 CARTON; Type 0: Not a Combination Product	08/15/2016	
2	NDC:0023-5870-80	1500 cm2 in 1 CARTON; Type 0: Not a Combination Product	08/15/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA016455	08/15/2016	

Labeler - Allergan, Inc. (144796497)

Revised: 12/2019

Allergan, Inc.