

**RANITIDINE- ranitidine tablet**  
**RANITIDINE COOL MINT- ranitidine tablet**  
**Granules India Ltd**

-----  
**Ranitidine Tablets**

**ACTIVE INGREDIENT(S)**

Ranitidine 150 mg

**PURPOSE**

Acid reducer

**USE(S)**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**WARNINGS**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**DO NOT USE**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**ASK A DOCTOR BEFORE USE IF**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn

- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.

### **STOP USE AND ASK DOCTOR IF**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

### **PREGNANCY/BREASTFEEDING**

ask a health professional before use.

### **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **DIRECTIONS**

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water
- to **prevent** symptoms, swallow 1 tablet with a glass of water
- **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- do not chew tablet (for cool mint tablets only)
- children under 12 years: ask a doctor

### **OTHER INFORMATION**

- do not use if individual blister unit is open or torn.
- do not use if printed foil under bottle cap is open or torn
- store at 20 °-25 °C (68 °-77 °F)
- avoid excessive heat or humidity

- this product is sodium and sugar free

## **INACTIVE INGREDIENTS**

**For 150 mg:** croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide

**For 150 mg cool mint:** acacia, croscarmellose sodium, FD&C Blue No. 1, magnesium stearate, menthol, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide

## **Questions or comments?**

- Contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM EST.

This product is not manufactured or distributed by Boehringer Ingelheim ® consumer health care products, distributor of Zantac 150mg and 150mg Cool Mint

## **Read the directions and warnings before use.**

Keep the carton. It contains important information including tips for managing heartburn.

## **Tips for managing heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables.
- Eat slowly and do not eat big meals
- if you are overweight, lose weight
- if you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

## **PRINCIPAL DISPLAY PANEL**

NDC 62207-774-58

\*Compare to the active ingredient of Zantac 150 Cool Mint\*

COOL MINT TABLETS

## Ranitidine Tablets, USP 150 mg

Acid Reducer

PREVENTS &amp; RELIEVES HEARTBURN

associated with acid indigestion  
and sour stomach

20 (2x10) Tablets

Un Varnish area

LOT:  
EXP:

## Drug Facts

**Active ingredient (in each tablet)...Purpose**  
Ranitidine 150 mg  
(as ranitidine hydrochloride 168 mg).....Acid reducer**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach

brought on by eating or drinking certain foods and beverages

**Warnings****Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

2000000XXXXX

## Drug Facts (continued)

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with nighttime, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach or kidney disease

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take the product for more than 14 days

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
  - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
  - do not chew tablet
  - children under 12 years: ask a doctor

**Other information**

- do not use 1 individual blister unit as open orion.
- store at 20°-25°C (68°-77°F) ■ avoid excessive heat or humidity
- this product is sodium and sugar free

**Inactive ingredients**

anhydrous calcium sulfate, FD&C Blue No. 1, magnesium stearate, methylcellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide.

**Questions or comments?**

Contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM EST.

This product is not manufactured or distributed by Boehringer Ingelheim consumer health care products, distributor of Zantac 150 Cool Mint®. Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

Labeling Format Information	
Format	Description
Font Style	Helvetica Narrow, Bold, Italic
Drug Facts	10 pt
Drug Facts (continued)	9 pt
Header	8 pt
Body Text	6 pt
Table (solid square)	5 pt
Table	2 pt
Table	0.5 pt
Drug Title	10 pt
Justification	Left
Revision	02

**Tips for managing heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables.
- Eat slowly and do not eat big meals
- if you are overweight, lose weight

- if you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach



NDC 62207-774-58

2000000000X



acacia, croscarmellose sodium, FD&C Blue No. 1, magnesium stearate, menthol, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide

Distributed By:  
Granules USA, Inc.  
Parsippany, NJ 07054  
Toll-free: 1-877-770-3183

37/FR/AP/2003/FR

Format	Description
Font Style	8-point Arial or Arial, 6 pt.
Drop Caps	0 pt
Drop Caps (continued)	0 pt
Header	9 pt
Section Header	7 pt
Bullet (left margin)	6 pt
Text	8 pt
Table	8 pt
Equation	10 pt
Justification	Left
Paper Size	11"

Labeling Format Information	
Format	Description
Font Style	Helvetica Narrow, sans, bold
Drug Facts	10 pt
Drug Facts (continued)	9 pt
Header	7 pt
Body Text	6 pt
Bullet (solid square)	5 pt
Barline	2 pt
Headerline	0.5 pt
Drug Title	15 pt
Justification	Left
Revision	02



LOT: EXP: Un varnish coding area



307 PPIA/P12003/R  
MADE IN INDIA  
India  
Hyderabad - 500 081,  
Granules India Limited  
Panipat, NY 07054  
Toll-free: 1-877-770-3183  
Distributed By:  
Granules USA, Inc.

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

**NDC 62207-774-41**

**COOL MINT TABLETS**  
**MAXIMUM STRENGTH**  
**Ranitidine Tablets, USP**  
**150 mg**

**Acid Reducer**  
**PREVENTS & RELIEVES HEARTBURN**  
associated with acid indigestion  
and sour stomach

**Releases a Cooling Sensation in  
Mouth and Throat**

<sup>†</sup>Compare to the active ingredient of  
Zantac 150 Cool Mint<sup>®</sup>

**24 Tablets**  
**(24 Doses)**



**Drug Facts**

**Active ingredient** ..... **Purpose**  
(in each tablet)  
Ranitidine 150 mg  
(as ranitidine hydrochloride 168 mg) ... Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness** ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain** ■ frequent wheezing, particularly with heartburn ■ unexplained weight

**Drug Facts (continued)**

loss ■ nausea or vomiting ■ stomach pain  
■ kidney disease

**Ask a doctor or pharmacist before use if you**  
are taking a prescription drug. Acid reducers may  
interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health  
professional before use.

**Keep out of reach of children.** In case of overdose,  
get medical help or contact a Poison Control  
Center right away.

**Directions**

- adults and children 12 years and over:  
■ to **relieve** symptoms, swallow 1 tablet with a glass of water
- to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- do not chew tablet
- children under 12 years: ask a doctor

**Other information**

- do not use if printed foil under bottle cap is open or

**Drug Facts (continued)**

form ■ store at 20°-25°C (68°-77°F) ■ avoid excessive heat or humidity ■ this product is sodium and sugar free

**Inactive ingredients**  
acacia, croscarmellose sodium, FD&C Blue No. 1, magnesium stearate, menthol, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide

**Questions or comments?**  
Contact 1-877-770-3183  
Mon-Fri 8:00 AM EST to 5:00 PM EST.

<sup>†</sup>This product is not manufactured or distributed by  
Boehringer Ingelheim consumer health care products,  
distributor of Zantac 150 Cool Mint<sup>®</sup>

**Tips for managing heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables.
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach





Un varnish area



200000000000X

NDC 62207-773-47

MAXIMUM STRENGTH

**Ranitidine Tablets, USP**  
**150 mg**

**Acid Reducer**

**PREVENTS & RELIEVES HEARTBURN**  
associated with acid indigestion  
and sour stomach

**500 Tablets**  
(500 Doses)

<sup>1</sup>Compare to the active ingredient  
of Zantac 150<sup>®</sup>



**Drug Facts**

**Active ingredient (in each tablet).....Purpose**  
Ranitidine 150 mg  
(as ranitidine hydrochloride 168 mg).....**Acid reducer**

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers.

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulder; or lightheadedness • frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting • stomach pain • kidney disease

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

**Other information**

- do not use if printed oil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F) • avoid excessive heat or humidity
- this product is sodium and sugar free

**Inactive ingredients**

microcrystalline cellulose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide

**Questions or comments?**

Contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM EST

<sup>1</sup>This product is not manufactured or distributed by Boehringer Ingelheim consumer health care products, distributor of Zantac 150<sup>®</sup>

**Tips for managing heartburn**

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods and drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

Manufactured By:  
**Granules India Limited**  
Hyderabad - 500 081, India  
**MADE IN INDIA**


301RRUP2003/F/R

Distributed By:  
**Granules USA, Inc.**  
Parampany, NJ 07654  
Toll-free: 1-877-770-3183


Labeling format information	
Form	Description
Form	Granules, 150 mg, white, oval
150 mg Tablet	0.25
150 mg Tablet (chewable)	0.25
Tablet	0.25
150 mg Tablet	0.25
Tablet with imprint	0.25
Tablet	0.25
Tablet	0.25
150 mg Tablet	0.25
150 mg Tablet	0.25
150 mg Tablet	0.25



Labeling Format Information	
Format	Description
Font Style	Helvetica Narrow, Bold, Italic
Drug Facts	10 pt
Drug Facts (continued)	9 pt
Header	7 pt
Body Text	6 pt
Bullet (solid square)	5 pt
Barline	2 pt
Headline	0.5 pt
Drug Title	15 pt
Justification	Left
Revision	02




UNWINDING  
DIRECTION




- PRINTING AREA  
FILL ALL TEXT AND GRAPHIC INSIDE GREEN AREA
- GLUE AREA  
BLANK
- PAPER AREA

APP. DATE : 02-07-2015  
UNICK FTX-A-FORM AND PRINTERS LTD.

IP part-1 FRONT




IP part-1 BACK



Labeling Format Information	
Format	Description
Font Style	Helvetica Narrow, Bold, Italic
Drug Facts	10 pt
Drug Facts (continued)	9 pt
Header	7 pt
Body Text	6 pt
Bullet (solid square)	5 pt
Barline	2 pt
Headline	0.5 pt
Drug Title	15 pt
Justification	Left
Revision	02

UNWINDING  
DIRECTION




PRINTING AREA  
FILL ALL TEXT AND GRAPHIC INSIDE  
GREEN AREA

GLUE AREA  
BLANK


PAPER AREA

APP. DATE : 02-07-2015  
UNICK FIX-A-FORM AND PRINTERS LTD.

IP part-1 FRONT



IP part-1 BACK



Labeling Format Information	
Page	Description
Page 01	Product Name, Strength, and Form
Page 02	Drug Facts
Page 03	Warnings
Page 04	Directions
Page 05	Other Information
Page 06	QR Code
Page 07	Manufacturer Information

NDC 62207-774-47

COOL MINT TABLETS  
MAXIMUM STRENGTH

Ranitidine Tablets, USP  
150 mg

Acid Reducer

PREVENTS & RELIEVES HEARTBURN

associated with acid indigestion and  
sour stomach

Releases a Cooling Sensation in  
Mouth and Throat

500 Tablets  
(500 Doses)

\*Compare to the active ingredient of  
Zantac 150 Cool Mint\*



Important: Read all product information before using.

### Drug Facts

**Active ingredient (in each tablet).....Purpose**

Ranitidine 150 mg  
(as ranitidine hydrochloride 168 mg).....Acid reducer

### Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

### Warnings

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

### Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with

Manufactured By:  
**Granules India Limited**  
Hyderabad - 500 081, India  
**MADE IN INDIA**

Distributed By:  
**Granules USA, Inc.**  
Parsippany, NJ 07054  
Toll-free: 1-877-770-3183



200 00000 XXXX

M. L. No.: 37/R/RAP/2003/F/R

LOT:  
EXP:

164 x 64 mm

### Drug Facts (continued)

shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain  
■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducer may interact with certain prescription drugs

### Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water
  - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- do not chew tablet
- children under 12 years: ask a doctor

### Other information

- do not use if printed lot underbottle cap is open or torn
- store at 20°-25°C (68°-77°F) ■ avoid excessive heat or humidity
- this product is sodium and sugar free

### Inactive ingredients

acacia, croscarmellose sodium, FD&C Blue No. 1, magnesium stearate, menthol, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide

### Questions or comments?

Contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST.

\*This product is not manufactured or distributed by Boehringer Ingelheim consumer health care products, distributor of Zantac 150 Cool Mint\*

164 x 64 mm



NDC 62207-773-47

MAXIMUM STRENGTH

# Ranitidine Tablets, USP

150 mg

## Acid Reducer

**PREVENTS & RELIEVES HEARTBURN**  
associated with acid indigestion  
and sour stomach

500 Tablets  
(500 Doses)

<sup>†</sup> Compare to the active ingredient  
of Zantac 150\*



Important: Read all product information before using.

### Drug Facts

#### Active ingredient (in each tablet).....Purpose

Ranitidine 150 mg  
(as ranitidine hydrochloride 168 mg).....Acid reducer

#### Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

#### Warnings

**Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers

#### Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

#### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness** ■ chest pain or shoulder pain with

Manufactured By:  
**Granules India Limited**  
Hyderabad – 500 081, India  
**MADE IN INDIA**

Distributed By:  
**Granules USA, Inc.**  
Parsippany, NJ 07054  
Toll-free: 1-877-770-3183



200000 00XXXX

M. L. No.: 37/FR/AP/2003/FR

LOT:

EXP:

164 x 64 mm

### Drug Facts (continued)

shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent **chest pain** ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducer may interact with certain prescription drugs

#### Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

#### Other information

- do not use if printed foil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F) ■ avoid excessive heat or humidity
- this product is sodium and sugar free

#### Inactive ingredients

croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talcum, titanium dioxide

#### Questions or comments?

Contact 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST.

\*This product is not manufactured or distributed by Boehringer Ingelheim consumer health care products, distributor of Zantac 150\*

# RANITIDINE TABLETS, USP 150 mg Cool mint

Each film coated tablet contains

Ranitidine 150 mg

(as Ranitidine hydrochloride USP 168 mg)

Batch No. : XXXXXXXX

No. of Units : 10,000's

Mfg.Dt. : MM/YYYY

Exp.Dt. : MM/YYYY

NDC No. : 62207-774-32

Mfg.Lic.No. : 37/RR/AP/2003/F/R

Storage : Store at 20°-25°C (68°-77°F) Avoid excessive heat or humidity

Caution : For repacking only.

Country of Origin: INDIA

Product code : 700000001630

# RANITIDINE TABLETS, USP 150 mg

Each film coated tablet contains:

Ranitidine 150 mg

(as Ranitidine hydrochloride USP 168 mg)

Batch No. : XXXXXXXX

No. of Units : 10,000's

Mfg.Dt. : MM/YYYY

Exp.Dt. : MM/YYYY

NDC No. : 62207-773-32

Mfg.Lic.No. : 37/RR/AP/2003/F/R

Storage : Store at 20°-25°C (68°-77°F) Avoid excessive heat or humidity

Caution : For repacking only.

Country of Origin: INDIA

Product code : 700000001629



# RANITIDINE TABLETS, USP 150 mg Cool mint

Each film coated tablet contains

Ranitidine 150 mg

(as Ranitidine hydrochloride USP 168 mg)

Batch No. : XXXXXXXX

Mfg.Dt. : MM/YYYY

Exp.Dt. : MM/YYYY

NDC No. : 62207-774-36

Mfg.Lic.No. : 37/RR/AP/2003/F/R

No. of Units : 5x10,000's

Gross Wt. : 00.000 Kg

Tare Wt. : 00.000 Kg

Net Wt. : 00.000 Kg

Box number : 000

Storage : Store at 20°-25°C (68°-77°F) Avoid excessive heat or humidity

Caution : For repacking only.



(01)50362207774364(17)000000(10)XXXXXXX

Country of Origin: INDIA



(00)189038320000000004

Product code : 700000001630

# RANITIDINE TABLETS, USP 150 mg

Each film coated tablet contains

Ranitidine 150 mg

(as Ranitidine hydrochloride USP 168 mg)

Batch No. : XXXXXXXX

Mfg.Dt. : MM/YYYY

Exp.Dt. : MM/YYYY

NDC No. : 62207-773-36

Mfg.Lic.No. : 37/RR/AP/2003/F/R

No. of Units : 5x10,000's

Gross Wt. : 00.000 Kg


Tare Wt. : 00.000 Kg

Net Wt. : 00.000 Kg

Box number : 000

Storage : Store at 20°-25°C (68°-77°F) Avoid excessive heat or humidity

Caution : For repacking only.



(01)50362207773367(17)000000(10)XXXXXXX

Country of Origin: INDIA



(00)189038320000000004

Product code : 700000001629

# RANITIDINE

ranitidine tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:62207-773
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>RANITIDINE HYDROCHLORIDE</b> (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	150;G
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62207-773-58	2 in 1 CARTON	06/26/2019	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:62207-773-41	1 in 1 CARTON	06/26/2019	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:62207-773-47	1 in 1 CARTON	06/26/2019	
3		500 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:62207-	5 in 1 CARTON	06/26/2019	

4	773-36	3 IN 1 CARTON	06/26/2019	
4	NDC:62207-773-32	10000 in 1 BAG; Type 0: Not a Combination Product		

## RANITIDINE TABLETS, USP 150 mg

Each film coated tablet contains:

Ranitidine 150 mg

(as Ranitidine hydrochloride USP 168 mg)

Batch No. : XXXXXXXX

No. of Units : 10,000's

Mfg.Dt. : MM/YYYY

Exp.Dt. : MM/YYYY

NDC No. : 62207-773-32

Mfg.Lic.No. : 37/RR/AP/2003/F/R

Storage : Store at 20°-25°C (68°-77°F) Avoid excessive heat or humidity

Caution : For repacking only.

Country of Origin: INDIA

Product code : 700000001629

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210243	06/26/2019	

## RANITIDINE COOL MINT

ranitidine tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62207-774
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	150;G
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62207-774-58	2 in 1 CARTON	06/26/2019	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:62207-774-41	1 in 1 CARTON	06/26/2019	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:62207-774-47	1 in 1 CARTON	06/26/2019	
3		500 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:62207-774-36	5 in 1 CARTON	06/26/2019	
4		10000 in 1 BAG; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210243	06/26/2019	

**Labeler** - Granules India Ltd (915000087)

