# GOOD SENSE CHILDRENS PAIN AND FEVER- acetaminophen suspension L. Perrigo Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## Perrigo Children's Pain & Fever Drug Facts

### Active ingredient (in each 5 mL)

Acetaminophen 160 mg

## **Purpose**

Pain reliever/fever reducer

#### Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

#### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if your child has ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if your child has

liver disease

## Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

## When using this product

## do not exceed recommended dose (see overdose warning)

## Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

#### Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

#### **Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

<sup>\*</sup>or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

#### Other information

- each 5 mL contains: sodium 3 mg
- store at  $20-25^{\circ}$ C (68-77 $^{\circ}$ F)
- do not use if printed neckband is broken or missing

## **Inactive ingredients**

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

#### Questions or comments?

1-800-719-9260

## **Principal Display Panel**

Ages 2 – 11 Years

Children's Pain & Fever

Pain Reliever/Fever Reducer

Acetaminophen 160 mg per 5 mL

Oral Suspension

Fast, effective relief of children's pain and fever

Alcohol Free

Aspirin Free

Ibuprofen Free

Cherry Flavor

Compare to active ingredient of Children's Tylenol® Oral Suspension

100% SATISFACTION GUARANTEED

4 FL OZ (118 mL)





## **COODSENSE®**

GOODSENSE.

Ages 2-11 Years

Children's

Acetaminophen 160 mg per 5 mL

Oral Suspension

Fast, effective relief of children's pain and fever



Children's Tylenol Oral Suspension

4 FL 0Z (118 mL)

GOODSENSE.

Ages 2-11 Years

**Children's** Pain&Fever

> DIRECTIONS: STEP 1. Shake well before using



STEP 2. Remove child protective сар



STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT:

e see dosing chart or ask a doctor for your child's correct dose. Drug Facts

Active ingredient (in each 5 mL)

Purpose

- Uses temporarily:

  reduces fever

  relieves minor a ches and pains due to:

  rel
- Warnings

Liver warning: This product contains aceta minophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximudable sensor.

Incre than a user of the daily amount daily amount with other drugs containing acetaminophen Allergy afert. Acetaminophen may cause severe skin reactions. Symptoms may include:

### Skin reddening ### Distars ### Taskin reaction occurs, stop use and seek medical help

If a skin reaction occurs, stop use and seek medical help

Sore throat warning: If sore throat is severe, persist for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- Do not use

  with any other drug containing acetaminophen
  (prescription or nonprescription). If you are not sure
  whether a drug contains acetaminophen, ask a doctor
  or pharmacist.

  If you richl has ever had an allengic reaction to this
  product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

- Stop use and ask a doctor if
- Top use and ask a doctor if in pain gets worse or lasts more than 5 days fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- this product does not contain directions or complete warmings for adult use
   do not give more than directed (see overdose warming). warning)

  shake well before using

  mL = millliter

  find right dose on chart below. If possible, use weight

- min ngnt dose on chart below. It possible, use weig to dose; otherwise, use age. remove the child protective cap and squeeze your child's dose into the dosing que repeat dose every 4 hours while symptoms last do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-05	11	1E ml

\*or as directed by a doctor

Attention: use only endosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

#### Other information

- each 5 mL contains: sodium 3 mg
   store at 20-25°C (68-77°F)
   do not use if printed neckband is broken or missing

Inactive ingredients
anhydrous citric acid, butyliparaben, calcium sulfate,
carrageenan, FD&C red #40, flavor, glycerin,
high fructose com syup, kydroxyethyl cellulose,
microcrystalline cellulose and carboxymethylcellulose
sodium, propylene glycol, purified water, sodium
berizoate, sorbitol solution, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

Gluten Free

Distributed By

Perrigo® Allegan, MI 49010





17526 C2 C10

## GOOD SENSE CHILDRENS PAIN AND FEVER

acetaminophen suspension

## **Product Information**

Product Type HUMAN OTC DRUG NDC:0113-0608 Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)		
BUTYLPARABEN (UNII: 3QPI1U3FV8)		
CALCIUM SULFATE (UNII: WAT0 DDB505)		
CARRAGEENAN (UNII: 5C69 YCD2YJ)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)		
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		
SODIUM PHO SPHATE, TRIBASIC (UNII: A752Q30A6X)		

Product Characteristics			
Color	RED (opaque)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date	
1	NDC:0113-0608-26	1 in 1 CARTON	02/15/2016		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:0113-0608-28	1 in 1 CARTON	05/28/2013		
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	10/08/1993	

## Labeler - L. Perrigo Company (006013346)

Revised: 8/2018 L. Perrigo Company