

**KETO-TREAT- propylene glycol liquid  
IBA**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**KETO-TREAT  
(PROPYLENE GLYCOL)  
U.S.P.**

**CONTENTS:**

CONTENTS:  
Propylene Glycol.....100%

**USE DIRECTIONS:**

**PREVENTATIVE**

As an aid in the prevention of Acetonemia (Ketosis) in dairy cattle, administer 4 to 8 fluid ounces per head daily, beginning two weeks before freshening, and continue for 6 weeks after freshening (calving). Use as a drench or mix in the feed.

**TREATMENT**

In treatment for Acetonemia (Ketosis) in dairy cattle, administer 4 to 16 fluid ounces per day, per head for 10 days as a drench.

**FOR VETINARY USE ONLY  
NOT FOR HUMAN USE**

**NOTE:**some cows may not respond to treatment as a result of complicating factors such as milk fever, hardware disease, etc. The conditions must be treated in conjunction with the Acetonemis (Ketosis).

Consult your veterinarian if condition does not respond. As with all drenches, care should be exercised not to lift mouth of animal excessively high. Administer drench slowly, preferably with a device designated with drenching.

**KEEP BOTTLE TIGHTLY CLOSED FOR ANIMAL USE ONLY**

**KEEP OUT OF REACH OF CHILDREN**

**Keto-Treat**

For the prevention and treatment of acetonemia (ketosis) in dairy cows

## USE DIRECTIONS

### PREVENTATIVE

As an aid in the prevention of Acetonemia (Ketosis) in dairy cattle, administer 4 to 8 fluid ounces per head daily, beginning two weeks before freshening, and continue for 6 weeks after freshening (calving). Use as a drench or mix in the feed.

### TREATMENT

In treatment for Acetonemia (Ketosis) in dairy cattle, administer 4 to 16 fluid ounces per day, per head for 10 days as a drench.



# KETO-TREAT

(Propylene Glycol)

# U.S.P.

**FOR VETERINARY  
USE ONLY**

**NOT FOR HUMAN USE**

#### CONTENTS:

Propylene Glycol ..... 100%

#### NOTE:

Some cows may not respond to treatment as a result of complicating factors such as milk fever, hardware disease, etc. The conditions must be treated in conjunction with the Acetonemia (Ketosis).

Consult your veterinarian if condition does not respond. As with all drenches, care should be exercised not to lift mouth of animal excessively high. Administer drench slowly, preferably with a device designed for drenching.

**KEEP BOTTLE TIGHTLY CLOSED  
FOR ANIMAL USE ONLY**

**KEEP OUT OF REACH  
OF CHILDREN**

**FOR THE PREVENTION  
AND TREATMENT OF  
ACETONEMIA (KETOSIS)  
IN DAIRY COWS**

MANUFACTURED FOR:  
IBA INC.  
ANIMAL HEALTH DIVISION  
103 GILMORE DRIVE  
SUTTON, MASSACHUSETTS 01590



TAKE TIME

OBSERVE LABEL  
DIRECTIONS

MFG. DATE:

LOT NO.:

NET CONTENTS:

321EW

**KETO-TREAT**

propylene glycol liquid

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:29983-1038
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3) (PROPYLENE GLYCOL - UNII:6DC9Q167V3)	PROPYLENE GLYCOL	100 kg in 100 kg

### Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:29983-1038-2	3.95 kg in 1 PACKAGE		
2	NDC:29983-1038-6	59.19 kg in 1 DRUM		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
unapproved drug other		06/26/2015	

**Labeler - IBA (019494160)**

### Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Morgan Gallacher Inc. DBA in Custom Chemical Formulators Inc.		028311595	manufacture, api manufacture, pack

Revised: 11/2023

IBA