EEZY SUN SPF 30- meradimate, octinoxate, octocrylene, titanium dioxide, zinc oxide cream EEZY SUN SPF 50- meradimate, octinoxate, octocrylene, titanium dioxide, zinc oxide cream BIO EARTH MANUFACTURING (PTY) LTD

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### **Eezy Sun**

### **Actives**

Meradimate, Octinoxate, Octocrylene, Titanium Dioxide, Zinc Oxide

### **Inactives**

Bulbine Fruitescence leaf oil, Cellulose gum, Cetearyl alcohol, Glycerin, Glycerol Stearate, Methylparaben, Phenoxyethanol, Propylparaben, Vit E, Vit A, Water, White petrolatum, Xanthan gum

# Dosage and Administration

Directions

Apply liberally 15 minutes before sun exposure Reapply

PLEASE RECYCLE

DISTRIBUTED BY SANRIA26 (PTY) LTD 9428 ASHFORD PLACE BRENTWOOD, TN 37027 INFO@EEZYSUN.COM PRODUCT OF SOUTH AFRICA

After 40 minutes of swimming or sweating Immediately after towel drying At least every 2 hours.

Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

Limit time in the sun, espesially from 10 a.m - 2 p.m

Wear long-sleeved shirts, pants, hats and sunglasses Children under 6 months of age: Ask a docto

## Warnings


For external use only

Do not use: On Damaged on broken skin.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

### **Product Label SPF 30**



# **Drug Facts**

Directions Purpose

**Drug Facts** (continued)

■ Apply liberally 15 minutes before sun exposure
■ Reapply
■ After 40 minutes of swimming or sweating
■ Immediately after towel drying

■ At least every 2 hours.

■ Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

Limit time in the sun, espesially from
10 a.m - 2 p.m

Wear long-sleeved shirts, pants, hats and

- sunglasses

  Children under 6 months of age: Ask a doctor

### Other Information

■ Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients
Bulbine fruitescense leaf oil, cellulose gum,
cetearyl alcohol, glycerin, glycerol stearate,
methylparaben, phenoxyethanol, propylparaben
vitamin E, vitamin A, water, white petrolatum, xanthan gum,

DBE UV SMART ST



10 packets, 0.27 fl oz (8 ml) each









Active Ingredients
Meradimate 3%, Octinoxate 7.5%,
Octocrylene 4%, Titanium Oxide 5%,
Zinc Oxide 9%

Uses

USes

■ Helps prevent sunburn
■ If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings
For external use only
Do not use ■ on damaged or broken skin
When using this product ■ keep out of eyes. Rinse

with water to remove.

Stop use and ask a doctor if rash occurs Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.







**Product Label SPF 50** 



# **Drug Facts**

Directions Purpose

Active Ingredients
Meradimate 5%, Octinoxate 7.5%,
Octocrylene 8%, Titanium Oxide 7%
Zinc Oxide 11 %

### Uses

 Helps prevent sunburn
 If used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings
For external use only
Do not use ■ on damaged or broken skin
When using this product ■ keep out of eyes. Rinse

with water to remove.

Stop use and ask a doctor if rash occurs Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.



### Drug Facts (continued)

- Directions

  Apply liberally 15 minutes before sun exposure

  Reapply

  After 40 minutes of swimming or sweating

  Immediately after towel drying

  At least every 2 hours.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection
- measures including:

  Limit time in the sun, espesially from
  10 a.m 2 p.m

  Wear long-sleeved shirts, pants, hats and
- sunglasses

  Children under 6 months of age: Ask a doctor

### Other Information

■ Protect the product in this container from excessive heat and direct sun.

Inactive Ingredients
Bulbine fruitescense leaf oil, cellulose gum,
cetearyl alcohol, glycerin, glycerol stearate,
methylparaben, phenoxyethanol, propylparaben,
vitamin E, vitamin A, water, white petrolatum, xanthan gum,

\$50



\$50 \$50 SUNSCREEN **BROAD SPECTRUM SPF 50** WATER RESISTANT
(40 MINUTES)

10 packets, 0.27 fl oz (8 ml) each

. SNAP





**HOW TO OPEN THE SNAP PACKET** 





### **EEZY SUN SPF 30**

meradimate, octinoxate, octocrylene, titanium dioxide, zinc oxide cream

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70116-001	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
MERADIMATE (UNII: J9QGD60OUZ) (MERADIMATE - UNII:J9QGD60OUZ)	MERADIMATE	30 mg in 1 mL		
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	75 mg in 1 mL		
Octocrylene (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	Octocrylene	40 mg in 1 mL		
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP) (TITANIUM DIO XIDE - UNII:15FIX9 V2JP)	TITANIUM DIO XIDE	50 mg in 1 mL		

Inactive Ingredients			
Ingredient Name	Strength		
BULBINE FRUTESCENS WHOLE (UNII: M2U1C7UW6Y)			
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679 OBS 311)			
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)			
GLYCERIN (UNII: PDC6A3C0OX)			
GLYCERYL MONOSTEARATE (UNII: 230 O U9 XXE4)			
METHYLPARABEN (UNII: A2I8 C7HI9 T)			
PHENOXYETHANOL (UNII: HIE492ZZ3T)			
PROPYLPARABEN (UNII: Z8IX2SC1OH)			
.ALPHATOCOPHEROL (UNII: H4N855PNZ1)			
<b>VITAMIN A</b> (UNII: 81G40 H8 B0 T)			
WATER (UNII: 059QF0KO0R)			
PETROLATUM (UNII: 4T6H12BN9U)			
XANTHAN GUM (UNII: TTV12P4NEE)			

	P	ackaging			
l	#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
l	1	NDC:70116-001-01	8 mL in 1 POUCH; Type 0: Not a Combination Product	12/14/2017	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	12/14/2017	

# **EEZY SUN SPF 50**

meradimate, octinoxate, octocrylene, titanium dioxide, zinc oxide cream

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70116-002	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
MERADIMATE (UNII: J9QGD60OUZ) (MERADIMATE - UNII:J9QGD60OUZ)	MERADIMATE	50 mg in 1 mL	
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	75 mg in 1 mL	
Octocrylene (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	Octocrylene	80 mg in 1 mL	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP) (TITANIUM DIO XIDE - UNII:15FIX9 V2JP)	TITANIUM DIO XIDE	70 mg in 1 mL	
Zinc Oxide (UNII: SOI2LOH54Z) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	110 mg in 1 mL	

Inactive Ingredients		
Ingredient Name	Strength	
BULBINE FRUTESCENS WHO LE (UNII: M2U1C7UW6 Y)		
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)		
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)		
GLYCERIN (UNII: PDC6A3C0OX)		
GLYCERYL MONOSTEARATE (UNII: 230 OU9 XXE4)		
METHYLPARABEN (UNII: A2I8 C7HI9T)		
PHENOXYETHANOL (UNII: HIE492ZZ3T)		
PROPYLPARABEN (UNII: Z8IX2SC1OH)		
.ALPHATO COPHEROL (UNII: H4N855PNZ1)		
VITAMIN A (UNII: 81G40H8B0T)		
WATER (UNII: 059QF0KO0R)		
PETROLATUM (UNII: 4T6H12BN9U)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Ш	Packaging			
	# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
	1 NDC:70116-002-01	8 mL in 1 POUCH; Type 0: Not a Combination Product	12/14/2017	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	12/14/2017	

# Labeler - BIO EARTH MANUFACTURING (PTY) LTD (639768436)

Establishment					
Name	Address	ID/FEI	Business Operations		
BIO EARTH MANUFACTURING (PTY) LTD		639768436	manufacture(70116-001, 70116-002)		

Revised: 12/2017 BIO EARTH MANUFACTURING (PTY) LTD