

SUNMARK POVIDONE-IODINE TOPICAL- povidone-iodine solution
Strategic Sourcing Services LLC

Sunmark Povidone-Iodine 10% Topical Solution

sunmark Povidone-Iodine 10% Topical Solution

Active Ingredient

Povidone Iodine 10% (*Equivalent to 1% titratable Iodine)

Purpose

Antiseptic

Uses

First aid antiseptic to prevent infection in minor cuts, burns.

Warnings

For external use only.

Do not:

- use in the eyes.
- use longer than 1 week unless directed by a doctor.
- use on individuals who are allergic or sensitive to iodine.
- apply over large areas of the body.

Stop use and consult a doctor if

- condition persists or gets worse.
- irritation and redness develop and persist for more than 72 hours.

Ask a doctor in case of:

- deep or puncture wounds
- animal bites
- serious burns.

Keep out of reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- Clean the affected area
- Apply a small amount on the area 1 to 3 times daily.
- May be covered with a sterile bandage.
- If bandaged, let dry first.

Other Information

Store at room temperature, 20-25 ° C (68-77 ° F).

Inactive Ingredients:

Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water.

COMPARE TO BETADINE ACTIVE INGREDIENT* sunmark povidone-iodine 10% Topical Solution FIRST AID ANTISEPTIC 16 FL OZ (PT) 473mL *This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark Betadine

sunmark Povidone-Iodine 10% Topical Solution 8oz (49348-622-37) | sunmark Povidone-Iodine 10% Topical Solution 16oz (49348-622-38)

sunmark™ COMPARE TO BETADINE® ACTIVE INGREDIENT* NDC 49348-622-37

povidone-iodine

10% Topical Solution

FIRST AID ANTISEPTIC

8 FL OZ (237 mL)

Active Ingredient	Purpose
Povidone Iodine 10% (*Equivalent to 1% titratable iodine)	Antiseptic

Uses
First aid antiseptic to prevent infection in minor cuts, burns.

Warnings For external use only.
Do not:

- use in the eyes.
- use longer than 1 week unless directed by a doctor.
- use on individuals who are allergic or sensitive to iodine.
- apply over large areas of the body.

Stop use and consult a doctor if

- condition persists or gets worse.
- irritation and redness develop and persist for more than 72 hours.

Ask a doctor in case of:

- deep or puncture wounds ■ animal bites ■ serious burns.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- Clean the affected area.
- Apply a small amount on the area 1 to 3 times daily.
- May be covered with a sterile bandage.
- If bandaged, let dry first.

Other Information store at room temperature, 20-25° C (68-77° F).

Inactive Ingredients: Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water.

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark Betadine®

0 10939 14233 7

McKesson
Empowering Healthcare
Distributed By: McKesson
One East Street, San Francisco, CA 94104
Money Back Guarantee



**COMPARE TO BETADINE®
ACTIVE INGREDIENT*
NDC 49348-622-38**

povidone-iodine

10% Topical Solution

Drug Facts

Active Ingredient Povidone Iodine 10% (*Equivalent to 1% titratable iodine)	Purpose Antiseptic
Uses First aid antiseptic to prevent infection in minor cuts, burns.	
Warnings For external use only.	
Do not:	
<ul style="list-style-type: none"> ■ use in the eyes. ■ use longer than 1 week unless directed by a doctor. ■ use on individuals who are allergic or sensitive to iodine. ■ apply over large areas of the body. 	
Stop use and consult a doctor if	
<ul style="list-style-type: none"> ■ condition persists or gets worse. ■ irritation and redness develop and persist for more than 72 hours. 	
Ask a doctor in case of:	
<ul style="list-style-type: none"> ■ deep or puncture wounds ■ animal bites ■ serious burns. 	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions	
<ul style="list-style-type: none"> ■ Clean the affected area. ■ Apply a small amount on the area 1 to 3 times daily. ■ May be covered with a sterile bandage. ■ If bandaged, let dry first. 	
Other Information store at room temperature, 20-25° C (68-77° F).	
Inactive Ingredients: Citric Acid, Disodium Phosphate, Glycerin, Nonoxynol-9, Sodium Hydroxide and Purified Water.	



0 10939 14333 4

FIRST AID ANTISEPTIC

16 FL OZ (1 PT) 473 mL

MEKESON
Empowering Healthcare

Distributed By McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee

*This product is not manufactured or distributed by Purdue Frederick, owner of the registered trademark Betadine®

SUNMARK POVIDONE-IODINE TOPICAL

povidone-iodine solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-622
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	1 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
NONOXYNOL-9 (UNII: 48Q180SH9T)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

Packaging

	Marketing Start	Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-622-37	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/15/2017	
2	NDC:49348-622-38	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/15/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	06/28/2012	

Labeler - Strategic Sourcing Services LLC (116956644)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(49348-622) , analysis(49348-622) , pack(49348-622) , label(49348-622)

Revised: 12/2023

Strategic Sourcing Services LLC