

**R-PEN- penicillin g potassium powder, for solution**  
**Huvepharma, Inc.**

-----  
**R-Pen®**

**0.5 BU Front Panel**

**Antibiotic**

**R-Pen®**  
**(penicillin G potassium)**

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

**ANTIBIOTIC**  
**for Drinking Water**

**Nonsterile**

**Active Ingredient:**

0.500 billion units  
penicillin G potassium

Mix 1 packet to 333 gallons water

Store between 20°C - 25°C  
(68°F - 77°F) with excursions  
permitted between 15°C - 40°C  
(59°F - 104°F).

**For Oral Use in Turkeys Only**

**Keep Out of Reach of Children**

Made in Canada

Huvepharma®  
Manufactured for:  
Huvepharma, Inc.  
525 Westpark Dr, Suite 230  
Peachtree City, GA 30269

**Take Time**



**Observe Label  
Directions**

Restricted Drug (California) -  
Use only as directed.

**Not for human use**

Approved by FDA under ANADA # 200-106

KCP-J00018 (Rev. 05-2023)

Huvepharma and R-Pen are registered trademarks of Huvepharma EOOD  
104078A

### **Gravity Flow Watering Systems:**

Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

### **RESIDUE WARNINGS:**

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

### **PRECAUTIONS:**

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

### **INDICATIONS:**

For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

### **DOSAGE AND ADMINISTRATION:**

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

### **DIRECTIONS:**

Combine contents of entire package and approximately 1.5 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

### **Automatic Watering Systems:**

Pour the concentrated solution into a glass or plastic container, then add enough water to make 2.6 gallons (9.9 liters) of solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.

## **1.0 BU Front Panel**

### **Antibiotic**

**R-Pen®  
(penicillin G potassium)**

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a**

**licensed veterinarian.**

## **ANTIBIOTIC for Drinking Water**

### **Nonsterile**

#### **Active Ingredient:**

1.0 billion units  
penicillin G potassium

Mix 1 packet to 666 gallons water

Store between 20°C - 25°C  
(68°F - 77°F) with excursions  
permitted between 15°C - 40°C  
(59°F - 104°F).

### **For Oral Use in Turkeys Only**

### **Keep Out of Reach of Children**

Made in Canada

### **Huvepharma**

Manufactured for:  
Huvepharma, Inc.  
525 Westpark Dr, Suite 230  
Peachtree City, GA 30269

**Take Time**



**Observe Label  
Directions**

Restricted Drug (California) -

Use only as directed.

**Not for human use**

Approved by FDA under ANADA # 200-106

KCP-J00019 (Rev. 05-2023)

Huvepharma and R-Pen are registered trademarks of Huvepharma EOOD

104079A

### **Gravity Flow Watering Systems:**

Pour the concentrated solution into enough water to make 666 gallons (2521 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

**RESIDUE WARNINGS:**

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

**PRECAUTIONS:**

For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

**INDICATIONS:**

For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

**DOSAGE AND ADMINISTRATION:**

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

**DIRECTIONS:**

Combine contents of entire package and approximately 3 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

**Automatic Watering Systems:**

Pour the concentrated solution into a glass or plastic container, then add enough water to make 5.2

gallons (19.7 liters) of solution. [This amount of solution will medicate 666 gallons (2521 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.

**0.5 BU Packet Label Image**

**Gravity Flow Watering Systems:**  
 Pour the concentrated solution into enough water to make 333 gallons (1260 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.  
 Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

**RESIDUE WARNINGS:**  
 Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.  
 Do not use in turkeys producing eggs for human consumption.

**PRECAUTIONS:**  
 For best results, the treatment should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

**Antibiotic**

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**R-Pen<sup>®</sup>**  
**(penicillin G potassium)**

**ANTIBIOTIC FOR DRINKING WATER**  
**Nonsterile**

**Active Ingredient:**  
 0.500 billion units penicillin G potassium

Mix 1 packet to 333 gallons water  
 Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).  
**For Oral Use in Turkeys Only**  
**Keep Out of Reach of Children**

Made in Canada

**HUVEPHARMA<sup>®</sup>**  
 Manufactured for:  
 Huvepharma, Inc.  
 525 Westpark Drive, Suite 230  
 Peachtree City, GA 30269



**Restricted Drug (California).**  
 Use only as directed.  
**Not for human use.**

Approved by FDA under ANADA # 200-106  
 KCP-J00018 (Rev. 05-2023)  
 Huvepharma and R-Pen are registered trademarks of Huvepharma EOOD 104078A

**INDICATIONS:**  
 For treatment of erysipelas in turkeys (caused by *Erysipelothrix rhusiopathiae*).

**DOSAGE AND ADMINISTRATION:**  
 Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

**DIRECTIONS:**  
 Combine contents of entire package and approximately 1.5 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

**Automatic Watering Systems:**  
 Pour the concentrated solution into a glass or plastic container, then add enough water

to make 2.6 gallons (9.9 liters) of solution. [This amount of solution will medicate 333 gallons (1260 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.

**HUVEPHARMA<sup>®</sup>**

**R-Pen<sup>®</sup>**

**1.0 BU Packet Label Image**

**Gravity Flow Watering Systems:**

**RESIDUE WARNINGS:**

**PRECAUTIONS:**  
 For best results, the treatment

Pour the concentrated solution into enough water to make 666 gallons (2521 liters) of drinking water. In gravity flow watering systems, prepare fresh solutions every 12 hours.

Drinking water prepared as directed above will contain 1,500,000 units of penicillin G per gallon (3.8 liters).

Treated turkeys must not be slaughtered for food during treatment and for one day after last treatment.

Do not use in turkeys producing eggs for human consumption.

should be started at the first sign of infection. If improvement is not noted after 3 to 4 days of treatment, consult a poultry pathologist or veterinarian. Keep this and all other medications out of the reach of children.

## Antibiotic

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

# R-Pen<sup>®</sup>

(penicillin G potassium)

### ANTIBIOTIC for Drinking Water

Nonsterile

**Active Ingredient:**  
1.0 billion units  
penicillin G potassium

Mix 1 packet to 666 gallons water  
Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

**For Oral Use in Turkeys Only**  
**Keep Out of Reach of Children**

Made in Canada



Manufactured for:  
Huvepharma, Inc.  
525 Westpark Drive, Suite 230  
Peachtree City, GA 30269

### Take Time



**Observe Label  
Directions**

Restricted Drug (California).  
Use only as directed.  
**Not for human use.**

Approved by FDA under ANADA # 200-106  
KCP-J00019 (Rev. 05-2023)

Huvepharma and R-Pen are registered trademarks of  
Huvepharma EOOD

104079A

### INDICATIONS:

For treatment of erysipelas in turkeys (caused by *Erysipelothrix*)

### DIRECTIONS:

Combine contents of entire package and approximately 3 pints of water

add enough water to make 5.2 gallons (19.7 liters) of solution. [This amount of solution will medicate 666

... (caused by *Myoporum  
rhusiopathiae*).

**DOSAGE AND ADMINISTRATION:**

Administer orally at a dosage of 1,500,000 units of penicillin per gallon (3.8 liters) of drinking water for 5 consecutive days.

... and approximately 6 pints of water in glass or plastic container. Stir to dissolve. Allow the concentrated solution to stand until the foam disappears.

**Automatic Watering Systems:** Pour the concentrated solution into a glass or plastic container, then

... gallons (2521 liters) of drinking water.] The automatic waterer should be adjusted to deliver 1 oz (30 ml) of stock solution per gallon (3.8 liters) of drinking water. In automatic watering systems, prepare fresh solutions every 12 hours.



## R-PEN

penicillin g potassium powder, for solution

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:23243-6768
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PENICILLIN G POTASSIUM</b> (UNII: VL775ZTH4C) (PENICILLIN G - UNII:Q42T66VG0C)	PENICILLIN G	500000000 [iU] in 500000000 [iU]

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-6768-1	25 in 1 PAIL		
1		500000000 [iU] in 1 PACKET		
2	NDC:23243-6768-2	12 in 1 PAIL		
2		10000000000 [iU] in 1 PACKET		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200106	04/13/2017	

**Labeler** - Huvepharma, Inc. (619153559)

**Registrant** - Huvepharma EOOD (552691651)