

CARE ONE IBUPROFEN INFANTS- ibuprofen suspension
American Sales Company

American Sales Company Ibuprofen Oral Suspension Drug Facts

Active ingredient (in each 1.25 mL)

Ibuprofen 50 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- measure with the dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child's mouth, toward the inner cheek

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart		
Weight (lbs)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

Other information

- store at 20-25°C (68-77°F)
- do not freeze
- do not use if printed neckband is broken or missing
- see bottom of box for lot number and expiration date

Inactive ingredients

anhydrous citric acid, artificial flavor, butylparaben, FD&C red #40, glycerin, hypromellose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the Active Ingredient in Motrin® Infants' Drops

For Ages 6 to 23 mos.

INFANTS'

CONCENTRATED DROPS

IBUPROFEN ORAL SUSPENSION

50 mg per 1.25 mL

PAIN RELIEVER/FEVER REDUCER (NSAID)

BERRY FLAVOR

Lasts up to 8 hours

Alcohol Free

Use Only with Enclosed Syringe

See New Warnings Information

CAREONE®

For Ages 6 to 23 mos. **INFANTS'**
CONCENTRATED DROPS

IBUPROFEN ORAL SUSPENSION

50 mg per 1.25 mL

PAIN RELIEVER/FEVER REDUCER (NSAID)

Child-Resistant Safety Cap
Read Instructions Carefully

Compare to the Active
Ingredient in Motrin®
Infants' Drops**

CAREONE®

For Ages 6 to 23 mos.

INFANTS'
CONCENTRATED DROPS

IBUPROFEN ORAL SUSPENSION

50 mg per 1.25 mL

**PAIN RELIEVER/FEVER REDUCER
(NSAID)**

BERRY FLAVOR

Lasts up to 8 hours

Alcohol Free

Use Only with
Enclosed Syringe



0.5 FL OZ

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children ages 6 months to 23 months of age.

Drug Facts

Active ingredient (in each 1.25 mL)	Purposes
Ibuprofen 50 mg (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily: ■ reduces fever
■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to

See New Warnings Information

(15 mL)


children under 3 years of age unless directed by doctor. ▶


LOT NO.

EXP.

: 05705 D4 C5

**CODE
AREA**





Drug Facts (continued)

Do not use ■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ child has asthma ■ child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ give with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint ■ vomits blood ■ has bloody or black stools
 - has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed

- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- measure with the dosing device provided. Do not use with any other device.
- dispense liquid slowly into the child's mouth, toward the inner cheek
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart

Weight (lbs)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL



Drug Facts (continued)

Other information

- store at 20-25°C (68-77°F)
- do not freeze
- do not use if printed neckband is broken or missing
- see bottom of box for lot number and expiration date

Inactive ingredients

anhydrous citric acid, artificial flavor, butylparaben, FD&C red #40, glycerin, hypromellose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Motrin® Infants' Drops.


GLUTEN FREE

DISTRIBUTED BY:
AMERICAN SALES CO.
4201 WALDEN AVE.,
LANCASTER, NY 14086
www.care1.info
©2011 S&S Brands, Inc.

**Quality guaranteed
or your money back.**



3 4 1520 31081 2



CARE ONE IBUPROFEN INFANTS

ibuprofen suspension

Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:41520-057
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (IBUPROFEN)	IBUPROFEN	50 mg in 1.25 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
BUTYLPARABEN	
FD&C RED NO. 40	
GLYCERIN	
HYPROMELLOSES	
POLYSORBATE 80	
PROPYLENE GLYCOL	
WATER	
SODIUM BENZOATE	
SORBITOL	
SUCROSE	
XANTHAN GUM	

Product Characteristics

Color	PINK (light)	Score	
Shape		Size	
Flavor	FRUIT (mixed)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-057-05	1 in 1 CARTON		

1	15 mL in 1 BOTTLE		
---	-------------------	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075217	05/11/2004	

Labeler - American Sales Company (809183973)

Revised: 10/2012

American Sales Company