

L-S 50- lincomycin hydrochloride and spectinomycin sulfate tetrahydrate powder, for solution
Zoetis Inc.

L-S 50
WATER SOLUBLE® POWDER
lincomycin-spectinomycin

L-S 50
WATER SOLUBLE® POWDER
lincomycin-spectinomycin
Antibacterial and Antimycoplasmal

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
FOR USE IN CHICKENS ONLY

INDICATIONS AND USAGE

For use in chickens up to 7 days of age as an aid in the control of: Airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin. Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.

DOSAGE AND ADMINISTRATION

Dosage

Provide 2 grams (g) antibiotic activity per gallon of drinking water. Administer as the sole source of water for the first 5 to 7 days of life.

Administration

Amount of L-S 50	Amount of Drinking Water
1 bottle	25 gallons
For proportioners delivering 1 ounce of solution per gallon of drinking water, dissolve contents of 5 bottles in each gallon of proportioner solution.	

Important: Chickens should consume water at the following approximate rate to insure intake of the required dose of lincomycin-spectinomycin indicated:

Broilers and Layer Replacements (Light and Heavy)

Age (Weeks)	Daily Water Intake Gallons/1000	Dosage mg Antibiotic/lb
1	5	50-65

PRECAUTION

Discard medicated drinking water daily and replace with fresh medicated drinking water.

WARNING

Not for human use.

Storage

Store at Controlled Room Temperature 20° to 25° C (68° to 77° F).

Restricted Drug (California)-Use Only as Directed

Made in China

Distributed by: Zoetis Inc.

Kalamazoo, MI 49007

500809

40039820

PRINCIPAL DISPLAY PANEL - 75 g Bottle Label

TAKE TIME

OBSERVE LABEL

DIRECTIONS

This bottle contains:

Lincomycin hydrochloride, equivalent to
lincomycin
16.7 grams

Spectinomycin sulfate tetrahydrate,
equivalent to spectinomycin
33.3 grams

Total Antibiotic Activity
50.0 grams

zoetis

Approved by FDA under NADA # 046-109

**PEEL FROM THE RIGHT
FOR SPANISH LANGUAGE**

**NET WT. 2.65 OZ
(75.0 grams)**



DIRECTIONS FOR USE
INDICATIONS AND USAGE: For use in chickens up to 7 days of age as an aid in the control of: Airsacculitis caused by either *Mycoplasma synoviae* or *Mycoplasma gallisepticum* susceptible to lincomycin-spectinomycin. Complicated Chronic Respiratory Disease (Air Sac Infection) caused by *Escherichia coli* and *M. gallisepticum* susceptible to lincomycin-spectinomycin.
DOSAGE AND ADMINISTRATION
Dosage: Provide 2 grams (g) antibiotic activity per gallon of drinking water. Administer as the sole source of water for the first 5 to 7 days of life.
Administration:

Amount of L-S 50	Amount of Drinking Water
1 bottle	25 gallons

For proportioners delivering 1 ounce of solution per gallon of drinking water, dissolve contents of 5 bottles in each gallon of proportioner solution.

Important: Chickens should consume water at the following approximate rate to insure intake of the required dose of lincomycin-spectinomycin indicated:

Broilers and Layer Replacements (Light and Heavy)		
Age (Weeks)	Daily Water Intake Gallons/1000	Dosage mg Antibiotic/lb
1	5	50-65

500809
40012436

PRECAUTION: Discard medicated drinking water daily and replace with fresh medicated drinking water.
WARNING: Not for human use. Store at Controlled Room Temperature 20° to 25° C (68° to 77° F).
Restricted Drug (California)-Use Only as Directed

Made in China
Distributed by: Zoetis Inc.
Kalamazoo, MI 49007



LOT:
EXP:

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian. **FOR USE IN CHICKENS ONLY**
 This bottle contains:
 Lincomycin hydrochloride, equivalent to lincomycin 16.7 grams
 Spectinomycin sulfate tetrahydrate, equivalent to spectinomycin 33.3 grams
zoetis Total Antibiotic Activity 50.0 grams
 NADA #46-109, Approved by FDA
NET WT. 2.65 OZ (75.0 grams)
 PEEL FROM THE RIGHT FOR SPANISH LANGUAGE



INDICACIONES DE USO
INDICACIONES Y USO: Para usar en pollos de hasta 7 días de edad, con el fin de ayudar a controlar: La alveolitis causada por *Mycoplasma synoviae* o *Mycoplasma gallisepticum* susceptible a la lincomicina-espectinomicina. Enfermedad Respiratoria Crónica Complicada (Infección de los alvéolos) causada por *Escherichia coli* y *M. gallisepticum* susceptible a la lincomicina-espectinomicina.
DOSIS Y ADMINISTRACIÓN
Dosis: Proporcionar 2 gramos (g) de actividad antibiótica por cada galón de agua de bebida. Adminístrese como fuente única de agua durante los primeros 5 a 7 días de vida.
Administración:

Cantidad de L-S 50	Cantidad de agua de bebida
1 frasco	25 galones

Para los surtidores que entregan 1 onza de solución por cada galón de agua potable, disuelva el contenido de 5 frascos en cada galón de solución del surtidor fraccionador.

Importante: Los pollos deben consumir agua en las siguientes cantidades aproximadas, para asegurar la ingestión de la dosis requerida de la lincomicina-espectinomicina indicada:

Pollos de engorde (broiler) y reemplazo de ponedoras (aves livianas y pesadas)		
Edad (semanas)	Ingestión diaria de agua (galones/1000)	Dosis (mg antibiótico/libra)
1	5	50-65

PRECAUCIÓN: Deseche el agua de beber restante todos los días y cámbiela por agua de beber fresca que contiene el medicamento.

ADVERTENCIA: No usar en seres humanos. Almacénese a temperaturas controladas entre 20° y 25° C (68° a 77° F). Fármaco restringido (California)—Úsese solamente según las indicaciones. Fabricado en China. Dist. por: Zoetis Inc. Kalamazoo, MI 49007

ADVERTENCIA: La ley Federal (EE. UU.) restringe el uso de este fármaco por o bajo prescripción de un veterinario con licencia. **PARA USO EN POLLOS SOLAMENTE**
 Este frasco contiene:
 Clorhidrato de lincomicina, equivalente a lincomicina 16.7 gramos
 Sulfato de espectinomicina tetrahidratada, equivalente a espectinomicina 33.3 gramos
zoetis Actividad antibiótica total 50.0 gramos
 Núm. NADA #46-109, Aprobado por la FDA
PESO NETO 75.0 gramos (2.65 onzas)

L-S 50

lincomycin hydrochloride and spectinomycin sulfate tetrahydrate powder, for solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:54771-1108
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LINCOMYCIN HYDROCHLORIDE (UNII: M6T05Z2B68) (LINCOMYCIN - UNII:BOD072YW0F)	LINCOMYCIN	16.7 g in 75 g
SPECTINOMYCIN SULFATE TETRAHYDRATE (UNII: YS91P54918) (SPECTINOMYCIN - UNII:93AK1U6QF)	SPECTINOMYCIN	33.3 g in 75 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-1108-1	75 g in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA046109	06/27/1973	

Labeler - Zoetis Inc. (828851555)

Revised: 12/2023

Zoetis Inc.