

INFLUENZINUM- influenza a virus a/hong kong/2671/2019 ivr-208 (h3n2) hemagglutinin antigen (formaldehyde inactivated), influenza a virus a/guangdong-maonan/swl1536/2019 cnic-1909 (h1n1) hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/washington/02/2019 hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 hemagglutinin antigen (formaldehyde inactivated) pellet

Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

INFLUENZINUM 15C

INFLUENZINUM 15C HPUS**

(**contains 0.443 mg of the active ingredient per pellet)

After Effects of Flu or Flu-Like Symptoms*

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose

Made in France

1-800-BOIRON-1 (1-800-264-7661),

BoironUSA.com Info@boiron.com

Distributed by Boiron, Inc. Newtown Square, PA 19073



Symptoms:



Lot:

Exp:

3

06960

90912

5

Contains approx. 80 pellets.

US

Peel for Drugs Facts and instructions for use.

NEOPATHIC PRACTICE,
NOT FDA EVALUATED.

Drug Facts

Active ingredient:** See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: ■ Do not use if pellet dispenser seal is broken.

Drug Facts (continued)

Inactive ingredients: lactose, sucrose

Questions or comments?

BoironUSA.com Info@Boiron.com

1-800-BOIRON-1 (1-800-264-7661)

**C, K, CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

Distributed by Boiron Inc.

Newtown Square, PA 19073

How to
dispense pellets?



1 Turn tube upside down.



2 Twist until 5 pellets are dispensed into cap.



3 Carefully remove cap and use it to pour pellets under the tongue.

INFLUENZINUM

influenza a virus a/hong kong/2671/2019 ivr-208 (h3n2) hemagglutinin antigen (formaldehyde inactivated), influenza a virus a/guangdong-maonan/swl1536/2019 cnic-1909 (h1n1) hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/washington/02/2019 hemagglutinin antigen (formaldehyde inactivated), influenza b virus b/phuket/3073/2013 hemagglutinin antigen (formaldehyde inactivated) pellet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-2713
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: 9HB0XUS9TM) (INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:9HB0XUS9TM)	INFLUENZA B VIRUS B/PHUKET/3073/2013 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)	15 [hp_C] in 15 [hp_C]
INFLUENZA B VIRUS B/WASHINGTON/02/2019 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: 1G2CU2LIB5) (INFLUENZA B VIRUS B/WASHINGTON/02/2019 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:1G2CU2LIB5)	INFLUENZA B VIRUS B/WASHINGTON/02/2019 HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)	15 [hp_C] in 15 [hp_C]
INFLUENZA A VIRUS A/HONG KONG/2671/2019 IVR-208 (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: THG3PU0KOK) (INFLUENZA A VIRUS A/HONG KONG/2671/2019 IVR-208 (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:THG3PU0KOK)	INFLUENZA A VIRUS A/HONG KONG/2671/2019 IVR-208 (H3N2) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)	15 [hp_C] in 15 [hp_C]
INFLUENZA A VIRUS A/GUANGDONG-MAONAN/SWL1536/2019 CNIC-1909 (H1N1) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) (UNII: R0BT251SA5) (INFLUENZA A VIRUS A/GUANGDONG-MAONAN/SWL1536/2019 CNIC-1909 (H1N1) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED) - UNII:R0BT251SA5)	INFLUENZA A VIRUS A/GUANGDONG-MAONAN/SWL1536/2019 CNIC-1909 (H1N1) HEMAGGLUTININ ANTIGEN (FORMALDEHYDE INACTIVATED)	15 [hp_C] in 15 [hp_C]

Inactive Ingredients

Ingredient Name	Strength
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	white	Score	
Shape	ROUND	Size	4mm
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0220-2713-41	15 [hp_C] in 1 TUBE; Type 0: Not a Combination Product	10/01/2020	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic			10/01/2020	

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment			
Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-2713)