MAXIMUM STRENGTH ANTACID- aluminum hydroxide, magnesium hydroxide, dimethicone suspension CVS PHARMACY

cvs max antacid

Active ingredients (in each 10 mL dose)

Aluminum hydroxide 800 mg (equivalent to dried gel, USP)

Magnesium hydroxide 800 mg

Simethicone 80 mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Do not take more than 60 mL (6 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals, at bedtime or as directed by a doctor
- children under 12 years of age: ask a doctor

Other information

- each 10 mL dose contains: magnesium 340 mg, sodium 10 mg
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

package Label



Compare to the active ingredients in Mylanta® Maximum Strength*

MAXIMUM STRENGTH Antacid _iquid

ALUMINA / MAGNESIA / SIMETHICONE

Antacid/Anti-gas Fast-Acting

Soothing relief of:

- Heartburn
- Acid Indigestion
- Pressure and Bloating



Mint Flavor

12 FL 0Z (355 mL)

205-06242-3 REV GC-0421

Drug Facts

TAMPER EVIDENT: DO NOT USE IF BREAKAWAY Band on Bottle Cap is missing or broken.

Active ingredients (in each 10 mL dose)

Purposes

Aluminum hydroxide (equivalent to dried gel, USP) 800 mg Antacid Magnesium hydroxide 800 mgAntacid Simethicone 80 mg Antigas

Uses relieves: • heartburn

- sour stomach
- acid indigestion
 symptoms referred to as gas

Warnings

Do not take more than 60 mL (6 doses) in a 24 hour period or use the maximum dosage for more than 2 weeks.

Ask doctor before use if you have

kidney disease
 a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before use

- adults and children 12 years of age and older: take 10 mL to 20 mL (1 to 2 doses) between meals, at bedtime or as directed by a doctor
- children under 12 years of age: ask a doctor

Other information

- each 10 mL dose contains: magnesium 340 mg, sodium 10 mg
- store at room temperature
 protect from freezing
- keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor, hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by Johnson & Johnson Merck, distributor of Mylanta®.

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 ©2015 CVS/Pharmacy CVS.com[®] 1-800-SHOP CVS

905-06242-4 REV GC-0521



CVS Quality



MAXIMUM STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-619

Route of Administration ORAL

Active Ingredient/Active Moiety

Basis of Ingredient Name Strength Strength

ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE -UNII:5QB0T2IUN0)

ALUMINUM HYDROXIDE 800 mg in 10 mL

MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients	
Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC10H)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:69842-619- 12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	09/01/2020	

Labeler - CVS PHARMACY (062312574)

Registrant - GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(69842-619)	

Revised: 10/2023 CVS PHARMACY