


32OZ HYDROGEN PEROXIDE- 32oz hydrogen peroxide liquid
Collective Science Integration

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.



1.75" x 4"

Drug Facts	
Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent
Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth	
Warnings For external use only	
Do not use • in the eyes or apply over large areas of the body • longer than 1 week	
Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns	
Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions First aid antiseptic • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first	
Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor	
Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.	
Inactive ingredients purified water	
Dist. by Collective Science Integration El Monte, CA 91731 Made in USA www.csihealth.org	
8 60005 37344 4	

1.75" x 4"



Drug Facts	
Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent
Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth	
Warnings For external use only	
Do not use • in the eyes or apply over large areas	

HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



32 FL. OZ. (1 QT) / 946 mL

2" x 4.5"

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions First aid antiseptic • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration

El Monte, CA 91731

Made in USA

www.csihealth.org



2" x 4.5"



HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



16 FL. OZ. (1 PT) / 473 mL

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions First aid antiseptic • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration

El Monte, CA 91731 Made in USA





1.75" x 4"



HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



32 FL. OZ. (1 QT) / 946 mL

2" x 4.5"



1.75" x 4"

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731
Made in USA
www.csihealth.org



2" x 4.5"

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

HYDROGEN PEROXIDE
TOPICAL SOLUTION USP

**FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT**

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



16 FL. OZ. (1 PT) / 473 mL

1.75" x 4"

of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731 Made in USA
www.csihealth.org



8 60005 37344 4

1.75" x 4"



HYDROGEN PEROXIDE
TOPICAL SOLUTION USP

**FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT**

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings
For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns


Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.


Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor


Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.




2" x 4.5"




2" x 4.5"



1.75" x 4"



1.75" x 4"



HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



32 FL. OZ. (1 QT) / 946 mL

2" x 4.5"



COLLECTIVE
SCIENCE
INTEGRATION

HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions First aid antiseptic • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731
Made in USA
www.csihealth.org



2" x 4.5"

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions First aid antiseptic • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

16 FL. OZ. (1 PT) / 473 mL

1.75" x 4"



HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



32 FL. OZ. (1 QT) / 946 mL

2" x 4.5"



Dist. by Collective Science Integration
El Monte, CA 91731 Made in USA
www.csihealth.org



1.75" x 4"

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731
Made in USA
www.csihealth.org



2" x 4.5"

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

HYDROGEN PEROXIDE
TOPICAL SOLUTION USP

**FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT**

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



16 FL. OZ. (1 PT) / 473 mL

1.75" x 4"

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731 Made in USA
www.csihealth.org



8 60005 37344 4

1.75" x 4"



HYDROGEN PEROXIDE
TOPICAL SOLUTION USP

**FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT**

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings
For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold

MADE IN THE
USA

32 FL. OZ. (1 QT) / 946 mL

2" x 4.5"

away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731
Made in USA
www.csihealth.org

8 60005 37248 5

2" x 4.5"

 **COLLECTIVE SCIENCE INTEGRATION**

HYDROGEN PEROXIDE
TOPICAL SOLUTION USP

**FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT**

For treatment of minor
cuts & abrasions
For use as a gargle or rinse

MADE IN THE
USA

16 FL. OZ. (1 PT) / 473 mL

1.75" x 4"

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings
For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731 Made in USA
www.csihealth.org

8 60005 37344 4

1.75" x 4"

 **COLLECTIVE SCIENCE INTEGRATION**

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



32 FL. OZ. (1 QT) / 946 mL

2" x 4.5"



HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions First aid antiseptic • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration

El Monte, CA 91731

Made in USA

www.csihealth.org



2" x 4.5"

Drug Facts

Active ingredients Purpose

Hydrogen peroxide (stabilized) 3%.....First aid antiseptic
Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions First aid antiseptic • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive inaredients purified water

16 FL. OZ. (1 PT) / 473 mL

1.75" x 4"

Dist. by Collective Science Integration
El Monte, CA 91731 Made in USA
www.csihealth.org



1.75" x 4"



HYDROGEN PEROXIDE

TOPICAL SOLUTION USP

FIRST AID ANTISEPTIC
ORAL DEBRIDING AGENT

For treatment of minor
cuts & abrasions
For use as a gargle or rinse



32 FL. OZ. (1 QT) / 946 mL

2" x 4.5"

Drug Facts

Active ingredients	Purpose
Hydrogen peroxide (stabilized) 3%.....	First aid antiseptic Oral debriding agent

Uses • first aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only

Do not use • in the eyes or apply over large areas of the body • longer than 1 week

Ask a doctor before use if you have deep puncture wounds, animal bites or serious burns

Stop us and ask a doctor if • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash, or fever develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions **First aid antiseptic** • clean the affected area • apply a small amount of product on the affected area 1-3 times a day • may be covered with a sterile bandage • if bandaged, let dry first

Oral debriding agent: adults, children over 2 years of age: • mix with equal amount of water • swish around in the mouth over the affected area for at least 1 minute, then spit out • use up to 4 times daily after meals and at bedtime • children under 12 years of age should be supervised in the use • children under 2 years of age: consult a dentist, doctor

Other information • keep tightly closed and at controlled room temperature. Do not shake bottle. Hold away from face when opening.

Inactive ingredients purified water

Dist. by Collective Science Integration
El Monte, CA 91731
Made in USA
www.csihealth.org



2" x 4.5"

32OZ HYDROGEN PEROXIDE

32oz hydrogen peroxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:90055-208
Route of Administration	ORAL, TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	3 mL in 3 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:90055-208-32	946 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	09/24/2020	

Labeler - Collective Science Integration (133591508)**Establishment**

Name	Address	ID/FEI	Business Operations
P&P International		133591508	manufacture(90055-208)

Revised: 9/2020

Collective Science Integration