

EXCEDRIN PM TRIPLE ACTION CAPLETS- acetaminophen, aspirin (nsaid) and diphenhydramine citrate tablet, coated
Haleon US Holdings LLC

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Diphenhydramine citrate 38 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain Reliever

Pain Reliever

Nighttime sleep-aid

Uses

- for the temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)

- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have glaucoma
- you have a breathing problem such as emphysema or chronic bronchitis
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
taking

- a prescription drug for:
 - diabetes
 - gout
 - arthritis

- any other drug, or are under a doctor's care for any serious condition
- any product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer
- sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- pain gets worse or last for more than 10 days
- painful area is red or swollen
- ringing in the ears or a loss of hearing occurs
- any new symptoms occur

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- do not use in children under 12 years of age
- adults and children 12 years of age and over: take 2 caplets at bedtime, with a full glass of water
- do not take more than 2 caplets in 24 hours, unless directed by a doctor

Other information

- **each caplet contains:** 75 mg calcium
- store at 20°-25°C (68°-77°F).
- close cap tightly after use.
- read all product information before using. Keep this box for important information.

Inactive ingredients

benzoic acid, calcium carbonate, FD&C blue no. 1 aluminum lake, ferric oxide yellow, hypromellose, low substituted hydroxypropyl cellulose, magnesium stearate, maltodextrin, medium-chain triglycerides, polydextrose, polysorbate 80, povidone, pregelatinized corn starch, silicified microcrystalline cellulose, stearic acid, talc, titanium dioxide, zinc stearate

Questions or comments?

1-800-468-7746

Principal Display Panel

EXCEDRIN

PM HEADACHE

Acetaminophen 250 mg, Aspirin 250 mg (NSAID)

and Diphenhydramine Citrate 38 mg

Pain Reliever/Nighttime Sleep-Aid

Caffeine-Free

Non-Habit Forming

ACTUAL SIZE

24 CAPLETS

TAMPER-EVIDENT BOTTLE

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH “SEALED for YOUR PROTECTION” IS BROKEN OR MISSING

Distributed by: Haleon, Warren, NJ 07059

©2023 Haleon or licensor. Trademarks are owned by or licensed to the Haleon group of companies. 62000000201994

HALEON

EXCEDRIN

PM HEADACHE



Acetaminophen 250 mg, Aspirin 250 mg (NSAID)
and Diphenhydramine Citrate 38 mg
Pain Reliever / Nighttime Sleep-Aid

Caffeine-Free
Non-Habit Forming

ACTUAL SIZE

24
CAPLETS



EXCEDRIN PM TRIPLE ACTION CAPLETS

acetaminophen, aspirin (nsaid) and diphenhydramine citrate tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0067-2056
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
DIPHENHYDRAMINE CITRATE (UNII: 4OD433S209) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE CITRATE	38 mg

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	

MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ZINC STEARATE (UNII: H92E6QA4FV)	

Product Characteristics

Color	BLUE (light blue)	Score	no score
Shape	RECTANGLE (oblong shaped tablet)	Size	17mm
Flavor		Imprint Code	EXPM
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-2056-24	1 in 1 CARTON	09/24/2014	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0067-2056-91	1 in 1 CARTON	09/24/2014	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/24/2014	

Labeler - Haleon US Holdings LLC (079944263)