ESOMEPRAZOLE- esomeprazole magnesium capsule, delayed release INNOVUS PHARMACEUTICALS, INC.

Esomeprazole Magnesium

Drug Facts

Active ingredient (in each capsule)

*Esomeprazole 20 mg (Each delayed-release capsule contains 22.3 mg esomeprazole magnesium trihydrate)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are:

• taking a prescription drug. Acid reducers may interact with certain prescription

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect
- 14-Day Course of Treatment
- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use.
- Store at 20°C to 25°C (68°F to 77°F). [See USP controlled room temperature.]
- keep the carton. It contains important information.

Inactive Ingredients

FD & C blue no 1, FD & C red no 3, ferrosoferric oxide, gelatin, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, mono and di

glycerides, polysorbate 80, potassium hydroxide, propylene glycol, shellac, sugar spheres (corn starch and sucrose), talc, titanium dioxide and triethyl citrate.

Questions?

Call toll-free weekdays 8:30 AM to 5 PM EST at **1-800-818-4555.**

TAMPER-EVIDENT FEATURES: Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or blue band around center of each capsule is broken or missing.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Distributed by: Innovus Pharmaceuticals, Inc Englewood, CO 80112

PRINCIPAL DISPLAY PANEL - 42 Capsule Bottle Carton

COMPARE TO THE ACTIVE INGREDIENT OF Nexium® 24HR[†]

NDC 57483-900-42

EsomepraCare™ Esomeprazole Magnesium Delayed-Release Capsules 20 MG* USP / ACID REDUCER

42 CAPSULES
Three 14-day courses of treatment

3 x 14 Count Bottles Inside

Treats Frequent Heartburn May take 1 to 4 days for full effect

24 HR



EsomepraCare[™] **Esomeprazole Magnesium**

Delayed-Release Capsules

20 MG* USP ACID REDUCER

- Tips for Managing Heartburn

 Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.

- some acolic truits and vegetates.

 Est slowly and do not eat bit gmeals.

 Do not eat late at night or just before beditine.

 Do not lie flat or bend over soon after eating.

 Raise the head of your bed

 Wear loose-fitting clothing around your stomach.

 If you are overweight, lies weight

 If you are overweight, lies weight

 If you are overweight, lies weight

Distributed by

Innovus Pharmaceuticals, Inc Englewood, CO 80112



COMPARE TO THE ACTIVE INGREDIENT OF Nextum® 24HR†

NDC 57483-900-42

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule)

Esomeprazole 20 mg _ (Each delayed-release capsule contains 22.3 mg esomeprazole magnesium trihydrate)

Uses

- treats frequent heartburn (occurs 2 or n days a week)
- not intended for immediate relief of heartburn this drug may take 1 to 4 days for full effect

Warnings Allergy alert: Do not use if you are allergic to

- Do not use if you have:

 trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain These may be signs of a serious condition. See your

EsomepraCare™

Esomeprazole Magnesium Delayed-Release Capsules

20 MG* USP / ACID REDUCER



42 CAPSULES

3 x 14 Count Bottles Inside

Treats Frequent Heartburn May take 1 to 4 days for full effect

† This product is not manufactured or distributed by Pfizer, distributor of Nexium® 24HR.

Expiration Date:



Drug Facts (continued)

- Ask a doctor before use if you have m had heartburn over 3 months. This may be
- a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
 unexplained weight loss
- nausea or vomiting

stomach pain

Ask a doctor or pharmacist before use if you are

■ taking a prescription drug. Acid reducers may interact with certain

Stop use and ask a doctor if your heartburn continues or

- your heartburn continues or worsens
 you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14
- days
 may take 1 to 4 days for full effect
 14-Day Course of Treatment
- swallow 1 capsule with a glass of water before eating in the morning

Drug Facts (continued)

- do not take more than 1 capsule a day
 swallow whole, Do not crush or chew capsules,
 do not use for more than 14 days unless directed by your doctor
 Repeated 14-Day Courses (If needed)
 you may repeat a 14-day course every 4 months
 do not take for more than 14 days or more often than every 4
 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information


- read the directions and warnings before use store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room
- keep the carton, it contains important information.

Inactive ingredients

Triactive Ingredients
TD & C blue not 0, FD & C red no 3, ferrosoferric oxide, gelatin,
hydroxypropt cellulose, hyprometose, magnesium stearate, methacrylic
acid copolymer dispersion, mono and di glycerides, polysorbate 80, potassium hydroxide, propylene glycol, shellac, sugar sphere starch and sucrose), talc, titanium dioxide and triethyl citrate. spheres (corn

Questions? Call toll free weekdays 8:30 AM to 5 PM EST at

TAMPER-EVIDENT FEATURES: Do not use if seal under bottle imprinted with "SEALED for YOUR PROTECTION" or blue



ESOMEPRAZOLE

esomeprazole magnesium capsule, delayed release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:57483-900

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ESOMEPRAZOLE MAGNESIUM (UNII: R6DXU4WAY9) (ESOMEPRAZOLE - UNII:N3PA6559FT)	ESOMEPRAZ OLE	20 mg		

FERROSOFERRIC OXIDE (UNII: XMOM87F357) GELATIN, UNSPECIFIED (UNII: 2G86QN327L) HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K) MAGNESIUM STEARATE (UNII: 70097M6I30) METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J) POLYSORBATE 80 (UNII: 6OZP39ZG8H) POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B710) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD) FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K) MAGNESIUM STEARATE (UNII: 70097M6I30) METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J) POLYSORBATE 80 (UNII: 6OZP39ZG8H) POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B710) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K) MAGNESIUM STEARATE (UNII: 70097M6I30) METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J) POLYSORBATE 80 (UNII: 60ZP39ZG8H) POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B710) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
MAGNESIUM STEARATE (UNII: 70097M6I30) METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J) POLYSORBATE 80 (UNII: 60ZP39ZG8H) POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B710) TALC (UNII: 75EV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J) POLYSORBATE 80 (UNII: 6OZP39ZG8H) POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B710) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H) POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B710) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: O8232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	MAGNESIUM STEARATE (UNII: 70097M6I30)	
POTASSIUM HYDROXIDE (UNII: WZ H3C48M4T) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B71O) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z 96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3) SHELLAC (UNII: 46N107B710) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z 96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
SHELLAC (UNII: 46N107B710) TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z 96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: O8232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	SHELLAC (UNII: 46N107B710)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM) STARCH, CORN (UNII: 08232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	TALC (UNII: 7SEV7J4R1U)	
STARCH, CORN (UNII: O8232NY3SJ) SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SUCROSE (UNII: C151H8M554) FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	STARCH, CORN (UNII: O8232NY3SJ)	
	SUCROSE (UNII: C151H8M554)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
	FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	

Product Characteristics			
Color	PINK	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	RG;50
Contains			

P	Packaging					
#	tem Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:57483-900- 42	3 in 1 CARTON	08/14/2020			
1		14 in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA212866	08/14/2020		

Labeler - INNOVUS PHARMACEUTICALS, INC. (962507187)

Establishment					
Name	Address	ID/FEI	Business Operations		
Ohm Laboratories Inc.		184769029	MANUFACTURE(57483-900)		

Revised: 5/2022 INNOVUS PHARMACEUTICALS, INC.