CLARITIN REDITABS- loratadine tablet, orally disintegrating Bayer Healthcare LLC.

Claritin ®

RediTabs ®

Drug Facts

Active ingredient (in each tablet)

Loratadine 10 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25°C (68° to 77°F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients

anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA.

PRINCIPAL DISPLAY PANEL - 10 Tablet Carton

Original Prescription Strength

NDC 11523-7157-2

Non-Drowsy* Claritin ® RediTabs® loratadine 10 mg/antihistamine

Indoor & Outdoor Allergies

*When taken as directed. See Drug Facts Panel.

24 Hour

Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes

• Itchy Throat or Nose

No Water Needed Melts in Your Mouth

10 ORALLY DISINTEGRATING TABLETS

Claritin Redilabs Redilabs

10 ORALLY DISINTEGRATING TABLETS FOR 10 DAYS OF RELIEF

Original Prescription Strength

Non-Drowsy*
Claritin®
Redilabs®

loratadine 10 mg/antihistamine

Indoor & Outdoor Allergies

NDC 11523-7157-2





Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose





*When taken as directed. See Drug Facts Panel.

10 ORALLY DISINTEGRATING TABLETS

The graphics on the front panel of this carton constitute trademarks of MSD Consumer Care, Inc.

© Copyright & Distributed by MSD Consumer Care, Inc., PO Box 377, Memphis, TN 38151 USA, a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All rights reserved. Product of Ireland.

27194-05





3. Place the tablet on tongue and close mouth. The tablet will disintegrate.



Centify push tablet our



Peel back



Follow these directions carefully. Do not attempt to push the tablet through the foll.



No Water Needed
 Melts in your mouth



QUESTIONS OF COMMENTS?
1-800-CLARITIN (1-800-252-7484) or www.claritin.com

•	sak a doctor	consumers with liver or kidney disease
	ask a doctor	children under 6 years of age
	1 tablet daily; not more than 1 tablet in 24 hours	and over

DIRECTIONS= place 1 tablet on tongue; tablet disintegrates, with or without water

If pregnant or breast-feeding, ask a health professional before use, Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.


medical neip right away.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek

directed may cause drowsiness.

When using this product do not take more than directed. Taking more than

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.



HSINAAV

Drug Facts (continued)

■ safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn Other information

■ store between 20° to 25° C (68° to 77° F)

 complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration ■ use tablet immediately after opening individual blister

Inactive ingredients anhydrous citric acid, gelatin, mannitol, mint flavor

enimstairlitnA, Active ingredient (in each tablet) Purpose Drug Facts

Loratadine 10 mg.

temporarily relieves these symptoms due to hay fever or other SASI


upper respiratory allergies:

ason ynnun = itchy, watery eyes

 itching of the nose or throat Buizeeus ■

Do not use if you have ever had an allergic reaction to this product or any of Warnings

its ingredients.



PRINCIPAL DISPLAY PANEL - 60 Tablet Twin Pack Carton TWIN PACK

Two 30ct Cartons 60 RediTabs ®

Original Prescription Strength

NDC 11523-4329-1

Non-Drowsy*
Claritin ®
RediTabs®
loratadine 10 mg/antihistamine

Indoor & Outdoor Allergies

*When taken as directed. See Drug Facts Panel.

24 Hour

Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat

or Nose

No Water Needed Melts in Your Mouth

60 ORALLY DISINTEGRATING TABLETS

TWIN PACK Two 30ct Cartons 60 RediTabs

Original Prescription Strength

NDC 11523-4329-1

Non-Drowsy* Claritin

Redilabs*

loratadine 10 mg/antihistamine

Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

No Water Needed Melts in Your Mouth



Indoor & Outdoor Allergies

 * When taken as directed. See Drug Facts Panel. 32437-00

60 ORALLY
DISINTEGRATING TABLETS

NO

VARNISH

Drug Facts

Active ingredient (in each tablet)

Purpose

USES temporarily relieves these symptoms due to hay lever or other upper respiratory allergies:

- rurny nose
- itchy, watery eyes
- = sneezing = itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a dector if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-leeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years 1 tablet daily; not more than 1 tablet in 24 hours children under 6 years of age ask a doctor consumers with liver or ask a doctor kidney disease

Drug Facts (continued)

Other information

- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F
- use tablet immediately after opening individual pilster
 complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients

anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Claritin Redilabs

- No Water Needed
- · Melts in your mouth




Follow these directions carefully. Do not attempt to push the tablet through the foil.









CLARITIN REDITABS

loratadine tablet, orally disintegrating

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11523-7157
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg

Inactive Ingredients	
Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
MANNITOL (UNII: 30WL53L36A)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor	MINT	Imprint Code	C10
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523- 7157-2	1 in 1 CARTON	11/27/2002	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11523- 7157-3	3 in 1 CARTON	11/27/2002	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:11523- 7157-4	2 in 1 CARTON	11/27/2002	
3		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:11523- 7157-7	4 in 1 CARTON	11/27/2002	
4		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:11523- 7157-8	5 in 1 CARTON	11/27/2002	
5		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
6	NDC:11523- 7157-9	7 in 1 CARTON	11/27/2002	
6		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Application Number or Monograph Marketing Start Marketing Category Citation Date Date			
NDA	NDA020704	11/27/2002	

CLARITIN REDITABS

loratadine tablet, orally disintegrating

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11523-4329
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

GELATIN (UNII: 2G86QN327L)	
MANNITOL (UNII: 30WL53L36A)	

Product Characteristics					
Color	white	Score	no score		
Shape	ROUND	Size	12mm		
Flavor	MINT	Imprint Code	C10		
Contains					

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:11523- 4329-1	2 in 1 PACKAGE, COMBINATION	11/27/2002				
1		3 in 1 CARTON					
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product					
2	NDC:11523- 4329-2	6 in 1 CARTON	11/27/2002				
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product					

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
NDA	NDA020704	11/27/2002			

Labeler - Bayer Healthcare LLC. (112117283)

Revised: 11/2023 Bayer Healthcare LLC.