MILK OF MAGNESIA- magnesium hydroxide liquid DOLGENCORP, INC.

dg milk of mag original

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL tablespoonful contains: magnesium 500 mg
- avoid freezing
- store at room temperature tightly closed

Inactive ingredients

purified water, sodium hypochlorite

Questions or Comments?

1-888-309-9030

package label



Compare to the active ingredient of Phillips'® Milk of Magnesia*

Milk of Magnesia

Magnesium Hydroxide 1200 mg per 15 mL **Saline Laxative**

- Comfortable, cramp free relief
- Sugar free
- Stimulant free

Original Flavor

12 FL OZ (355 mL)

207-19112-0 REV0221

mg per 15 mL

Drug Facts

TAMPER EVIDENT: Do not use if imprinted neckband is missing

Purpose

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg Saline laxative

- · relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- · kidney disease a magnesium-restricted diet · stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product, These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • shake well before use

- . do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

9 1 1		
age	dose	
adults and children 12 years and over	30 mL to 60 mL	
children 6 to 11 years	15 mL to 30 mL	
children under 6 years	ask a doctor	

Other information

- · each 15 mL contains; magnesium 500 mg
- · avoid freezing · store at room temperature tightly closed

Inactive ingredients purified water, sodium hypochlorite

Ouestions or comments? 1-888-309-9030

*This product is not manufactured or distributed by Bayer Healthcare LLC., distributors of Phillips'® brand Milk of Magnesia.

DISTRIBUTED BY OLD EAST MAIN CO. 100 MISSION RIDGE, GOODLETTSVILLE, TN 37072

100% SATISFACTION GUARANTEED!

(888) 309-9030

907-19112-0 REV GC0221

MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:55910-649

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients			
Ingredient Name	Strength		
WATER (UNII: 059QF0KO0R)			
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)			

l	Packaging				
	# Item Code Package Description		Marketing Start Date	Marketing End Date	
	1 NDC 12	:55910-649-	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	02/01/2021	

Labeler - DOLGENCORP, INC. (068331990)

Registrant - GCP Laboratories (965480861)

Establishment					
Name	Address	ID/FEI	Business Operations		
GCP Laboratories		965480861	manufacture(55910-649)		

Revised: 11/2023 DOLGENCORP, INC.