

PROMETHAZINE HYDROCHLORIDE- promethazine hydrochloride injection, solution

XGen Pharmaceuticals DJB, Inc.

Promethazine Hydrochloride Injection, USP

WARNINGS: RESPIRATORY DEPRESSION IN PEDIATRICS AND SEVERE TISSUE INJURY, INCLUDING GANGRENE

Respiratory Depression - Pediatrics

Promethazine hydrochloride injection should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression. Post-marketing cases of respiratory depression, including fatalities, have been reported with use of promethazine in pediatric patients less than 2 years of age. Caution should be exercised when administering promethazine hydrochloride injection to pediatric patients 2 years of age and older (see WARNINGS - Respiratory Depression).


Severe Tissue Injury, Including Gangrene

- Severe chemical irritation and damage to tissues regardless of the route of parenteral administration has been reported in patients treated with promethazine hydrochloride injection including gangrene, tissue necrosis, and thrombophlebitis; and in some cases, surgical intervention including fasciotomy, skin graft, and/or amputation have been required (see WARNINGS-Severe Tissue Injury, Including Gangrene).
- The use of promethazine hydrochloride injection by the following routes of administration is **CONTRAINDICATED**:
 - o Intravenous injection at concentrations greater than 1 mg/mL
 - o Intra-arterial injection
 - o Subcutaneous injection (see CONTRAINDICATIONS).
- The preferred route of administration is by deep intramuscular administration.
- Promethazine hydrochloride injection may be administered intravenously after dilution through an intravenous catheter inserted in a large vein. Preferably through a central venous catheter (see DOSAGE AND ADMINISTRATION).
- If pain occurs at the injection site during intravenous infusion, immediately discontinue the infusion and evaluate for possible arterial injection or perivascular extravasation, and initiate appropriate medical management.

DESCRIPTION

Promethazine hydrochloride injection, USP is a sterile, pyrogen-free solution for deep

intramuscular or intravenous administration. Promethazine hydrochloride (10H-phenothiazine-10-ethanamine, N, N, α -trimethyl-, monohydrochloride, ($\pm$)-) is a racemic compound and has the following structural formula:



C₁₇H₂₀N₂S.HCl

MW=320.88

Each mL contains promethazine hydrochloride, either 25 mg or 50 mg, edetate disodium 0.1 mg, calcium chloride 0.04 mg, sodium metabisulfite 0.25 mg and phenol 5 mg in Water for Injection, USP. pH 4.0 to 5.5; buffered with acetic acid-sodium acetate. Sealed under nitrogen.

Promethazine hydrochloride injection is a clear, colorless solution. The product is light sensitive. It should be inspected before use and discarded if either color or particulate is observed.

CLINICAL PHARMACOLOGY

Promethazine hydrochloride is a phenothiazine derivative which possesses antihistaminic, sedative, antimotion-sickness, antiemetic, and anticholinergic effects. Promethazine is a competitive H₁ receptor antagonist, but does not block the release of histamine. Structural differences from the neuroleptic phenothiazines result in its relative lack (1/10 that of chlorpromazine) of dopamine antagonist properties. Clinical effects are generally apparent within 5 minutes of an intravenous injection and within 20 minutes of an intramuscular injection. Duration of action is four to six hours, although effects may persist up to 12 hours. Promethazine hydrochloride is metabolized in the liver, with the sulfoxides of promethazine and N-desmethylpromethazine being the predominant metabolites appearing in the urine. Following intravenous administration in healthy volunteers, the plasma half-life for promethazine has been reported to range from 9 to 16 hours. The mean plasma half-life for promethazine after intramuscular administration in healthy volunteers has been reported to be 9.8 ± 3.4 hours.

INDICATIONS AND USAGE

Promethazine hydrochloride injection is indicated for the following conditions:

1. Amelioration of allergic reactions to blood or plasma.
2. In anaphylaxis as an adjunct to epinephrine and other standard measures after the acute symptoms have been controlled.
3. For other uncomplicated allergic conditions of the immediate type when oral therapy is impossible or contraindicated.
4. For sedation and relief of apprehension and to produce light sleep from which the patient can be easily aroused.
5. Active treatment of motion sickness.
6. Prevention and control of nausea and vomiting associated with certain types of

anesthesia and surgery.

7. As an adjunct to analgesics for the control of postoperative pain.
8. Preoperative, postoperative, and obstetric (during labor) sedation.
9. Intravenously in special surgical situations, such as repeated bronchoscopy, ophthalmic surgery, and poor-risk patients, with reduced amounts of meperidine or other narcotic analgesic as an adjunct to anesthesia and analgesia.

CONTRAINDICATIONS

The use of promethazine hydrochloride injection is contraindicated:

- In pediatric patients less than 2 years of age due to the risk of respiratory depression (see **WARNINGS - Respiratory Depression**).
- For use as an intravenous injection at concentrations greater than 1 mg/mL due to the risk of perivascular extravasation, unintentional intra-arterial injection, and intraneuronal or perineuronal infiltration (see **WARNINGS-Severe Tissue Injury, Including Gangrene and DOSAGE AND ADMINISTRATION**).
- For use as an intra-arterial injection due to the likelihood of severe arteriospasm and the possibility of resultant gangrene (**see WARNINGS-Severe Tissue Injury, Including Gangrene**).
- For use as a subcutaneous injection because chemical irritation and necrotic lesions have been reported (see **WARNINGS-Severe Tissue Injury, Including Gangrene**).
- In patients in a comatose state.
- In patients who have demonstrated an idiosyncratic reaction or hypersensitivity to promethazine hydrochloride or other phenothiazines.

WARNINGS

Respiratory Depression

Pediatrics

Promethazine hydrochloride injection should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression. Post-marketing cases of respiratory depression, including fatalities, have been reported with use of promethazine in pediatric patients less than 2 years of age. A wide range of weight-based doses of promethazine hydrochloride injection have resulted in respiratory depression in these patients.

Caution should be exercised when administering promethazine hydrochloride injection to pediatric patients 2 years of age and older. It is recommended that the lowest effective dose of promethazine hydrochloride injection be used in pediatric patients 2 years of age and older. Avoid concomitant administration of other drugs with respiratory depressant effects because of an association with respiratory depression, and sometimes death, in pediatric patients.

Other

Because of the risk of potentially fatal respiratory depression, use of promethazine

hydrochloride injection in patients with compromised respiratory function or patients at risk for respiratory failure (e.g. COPD, sleep apnea) should be avoided.

Severe Tissue Injury, Including Gangrene

- **Promethazine hydrochloride injection has been reported to cause severe chemical irritation and damage to tissues, including gangrene, regardless of the route of parenteral administration. Intra-arterial and subcutaneous administration have resulted in more significant complications. The use of promethazine hydrochloride injection by intravenous injection at concentrations greater than 1 mg/mL, intra-arterial injection, or subcutaneous injection is contraindicated (see CONTRAINDICATIONS).**
- **Irritation and damage can result from perivascular extravasation, intra-arterial injection, and intraneuronal or perineuronal infiltration. Inadvertent intra-arterial injection can cause severe arteriospasm. Other reported adverse reactions included burning, pain, erythema, swelling, sensory loss, palsies, paralysis, severe spasm of distal vessels, thrombophlebitis, venous thrombosis, phlebitis, abscesses, tissue necrosis, and gangrene. In some cases, surgical intervention, including fasciotomy, skin graft, and/or amputation have been required.**
- **The preferred route of administration of promethazine hydrochloride injection is by deep intramuscular administration.**
- **Promethazine hydrochloride injection may be administered intravenously after dilution through an intravenous catheter inserted in a large vein. Preferably through a central venous catheter (see DOSAGE AND ADMINISTRATION).**
- **If pain occurs at the injection site during intravenous infusion, immediately discontinue the infusion, evaluate for possible arterial injection or perivascular extravasation, and initiate appropriate medical management.**

CNS Depression

Promethazine hydrochloride injection may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery.

The impairment may be amplified by concomitant use of other central-nervous-system depressants such as alcohol, sedative/hypnotics (including barbiturates), general anesthetics, narcotics, narcotic analgesics, tricyclic antidepressants, and tranquilizers; therefore such agents should either be eliminated or given in reduced dosage in the presence of promethazine hydrochloride (see **PRECAUTIONS- Information for Patients and Drug Interactions**).

Lower Seizure Threshold

Promethazine hydrochloride injection may lower seizure threshold and should be used with caution in persons with seizure disorders or in persons who are using concomitant medications, such as narcotics or local anesthetics, which may also affect seizure threshold.

Bone-Marrow Depression

Promethazine hydrochloride injection should be used with caution in patients with bone-

marrow depression. Leukopenia and agranulocytosis have been reported, usually when promethazine hydrochloride has been used in association with other known marrow-toxic agents.

Neuroleptic Malignant Syndrome

A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with promethazine hydrochloride alone or in combination with antipsychotic drugs. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis and cardiac dysrhythmias).

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include 1) immediate discontinuation of promethazine hydrochloride, antipsychotic drugs, if any, and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

Since recurrences of NMS have been reported with phenothiazines, the reintroduction of promethazine hydrochloride should be carefully considered.

Sulfite Sensitivity

Promethazine hydrochloride injection contains sodium metabisulfite, a sulfite that may cause allergic-type reactions, including anaphylactic symptoms and life-threatening or less severe asthma episodes, in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people.

Visual Inspection

This product is light sensitive and should be inspected before use and discarded if either color or particulate is observed.

Cholestatic Jaundice

Administration of promethazine has been associated with reported cholestatic jaundice.

PRECAUTIONS

General

Drugs having anticholinergic properties should be used with caution in patients with narrow-angle glaucoma, prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal obstruction, and bladder-neck obstruction.

Promethazine hydrochloride injection should be used cautiously in persons with cardiovascular disease or impairment of liver function.

Information for Patients

Patients should be advised of the risk of respiratory depression, including potentially fatal respiratory depression in children less than 2 years of age (see **WARNINGS-Respiratory Depression**).

Patients should be advised of the risk of severe tissue injury, including gangrene (see **WARNINGS-Severe Tissue Injury, including Gangrene**). Patients should be advised to immediately report persistent or worsening pain or burning at the injection site.

Promethazine hydrochloride injection may cause marked drowsiness or impair the mental or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery. Pediatric patients should be supervised to avoid potential harm in bike riding or in other hazardous activities. The concomitant use of alcohol, sedative/hypnotics (including barbiturates), general anesthetics, narcotics, narcotic analgesics, tricyclic antidepressants, and tranquilizers may enhance impairment (see **WARNINGS-CNS Depression and PRECAUTIONS-Drug Interactions**).

Patients should be advised to report any involuntary muscle movements (see **ADVERSE REACTIONS-Paradoxical Reactions**).

Patients should be advised to avoid prolonged exposure to the sun (see **ADVERSE REACTIONS-Dermatologic**).

Drug Interactions

CNS Depressants

Promethazine hydrochloride injection may increase, prolong, or intensify the sedative action of central-nervous-system depressants, such as alcohol, sedative/hypnotics (including barbiturates), general anesthetics, narcotics, narcotic analgesics, tricyclic antidepressants, and tranquilizers; therefore, such agents should be avoided or administered in reduced dosage to patients receiving promethazine hydrochloride. When given concomitantly with promethazine hydrochloride injection, the dose of barbiturates should be reduced by at least one-half, and the dose of narcotics should be reduced by one-quarter to one-half. Dosage must be individualized. Excessive amounts of promethazine hydrochloride injection relative to a narcotic may lead to restlessness and motor hyperactivity in the patient with pain; these symptoms usually disappear with adequate control of the pain.

Epinephrine

Because of the potential for promethazine hydrochloride to reverse epinephrine's vasopressor effect, epinephrine should NOT be used to treat hypotension associated with promethazine hydrochloride injection overdose.

Anticholinergics

Concomitant use of other agents with anticholinergic properties should be undertaken with caution.

Monoamine Oxidase (MAO) Inhibitors

Drug interactions, including an increased incidence of extrapyramidal effects, have been reported when some MAO Inhibitors and phenothiazines are used concomitantly. This possibility should be considered with promethazine hydrochloride injection.

Drug/Laboratory Test Interactions

The following laboratory tests may be affected in patients who are receiving therapy with promethazine hydrochloride injection:

Pregnancy Tests

Diagnostic pregnancy tests based on immunological reactions between HCG and anti-HCG may result in false-negative or false-positive interpretations.

Glucose Tolerance Test

An increase in blood glucose has been reported in patients receiving promethazine hydrochloride.

Carcinogenesis, Mutagenesis and Impairment of Fertility

Long-term animal studies have not been performed to assess the carcinogenic potential of promethazine hydrochloride injection, nor are there other animal or human data concerning carcinogenicity, mutagenicity, or impairment of fertility. Promethazine hydrochloride injection was nonmutagenic in the Salmonella test system of Ames.

Pregnancy

Teratogenic Effects - Pregnancy Category C

Teratogenic effects have not been demonstrated in rat-feeding studies at doses of 6.25 and 12.5 mg/kg (approximately 2.1 and 4.2 times the maximum recommended human daily dose) of promethazine hydrochloride injection. Daily doses of 25 mg/kg intraperitoneally have been found to produce fetal mortality in rats.

There are no adequate and well-controlled studies of promethazine hydrochloride injection in pregnant women. Because animal reproduction studies are not always predictive of human response, promethazine hydrochloride injection should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Adequate studies to determine the action of the drug on parturition, lactation and development of the animal neonate have not been conducted.

Nonteratogenic Effects

Promethazine hydrochloride injection administered to a pregnant woman within two weeks of delivery may inhibit platelet aggregation in the newborn.

Labor and Delivery

Promethazine hydrochloride injection may be used alone or as an adjunct to narcotic analgesics during labor (see DOSAGE AND ADMINISTRATION). Limited data suggest that

use of promethazine hydrochloride injection during labor and delivery does not have an appreciable effect on the duration of labor or delivery and does not increase the risk of need for intervention in the newborn. The effect on later growth and development of the newborn is unknown. (**See** Pregnancy-Nonteratogenic Effects.)

Nursing Mothers

It is not known whether promethazine hydrochloride injection is excreted in human milk. Because many drugs are excreted in human milk, and because of the potential for serious adverse reactions in nursing infants from promethazine hydrochloride injection, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Promethazine hydrochloride injection is contraindicated for use in pediatric patients less than 2 years of age, because of the potential for fatal respiratory depression. Promethazine hydrochloride injection should be used with caution in pediatric patients 2 years of age and older (see **WARNINGS - Respiratory Depression**).

Antiemetics are not recommended for treatment of uncomplicated vomiting in pediatric patients, and their use should be limited to prolonged vomiting of known etiology. The extrapyramidal symptoms which can occur secondary to promethazine hydrochloride injection administration may be confused with the CNS signs of undiagnosed primary disease, e.g. encephalopathy or Reye's syndrome. The use of promethazine hydrochloride injection should be avoided in pediatric patients whose signs and symptoms may suggest Reye's syndrome or other hepatic diseases.

Excessively large dosages of antihistamines, including promethazine hydrochloride injection, in pediatric patients may cause sudden death (see **OVERDOSAGE**). Hallucinations and convulsions have occurred with therapeutic doses and overdoses of promethazine hydrochloride injection in pediatric patients. In pediatric patients who are acutely ill associated with dehydration, there is an increased susceptibility to dystonias with the use of promethazine hydrochloride injection.

Geriatric Use (patients approximately 60 years or older)

Since therapeutic requirements for sedative drugs tend to be less in geriatric patients, the dosage should be reduced for these patients.

ADVERSE REACTIONS

Respiratory Depression

Promethazine hydrochloride injection is contraindicated in pediatric patients less than 2 years of age, because of the potential for fatal respiratory depression. Promethazine hydrochloride injection should be used with caution in pediatric patients 2 years of age and older (see **WARNINGS - Respiratory Depression**).

Severe Tissue Injury, Including Gangrene

Promethazine hydrochloride injection has been reported to cause severe chemical

irritation and damage to tissues, including gangrene, regardless of the route of parenteral administration. Intra-arterial and subcutaneous injection have resulted in more significant complications (**see CONTRAINDICATIONS**).

Reported adverse reactions included burning, pain, erythema, swelling, sensory loss, palsies, paralysis, severe spasm of distal vessels, thrombophlebitis, venous thrombosis, phlebitis, abscesses, tissue necrosis, and gangrene (**see WARNINGS-Severe Tissue Injury, Including Gangrene**).

Central Nervous System

Drowsiness is the most prominent CNS effect of this drug. Sedation, somnolence, blurred vision, dizziness, confusion, disorientation, and extrapyramidal symptoms such as oculogyric crisis, torticollis, and tongue protrusion; lassitude, tinnitus, incoordination, fatigue, euphoria, nervousness, diplopia, insomnia, tremors, convulsive seizures, excitation, catatonic-like states, hysteria. Hallucinations have also been reported.

Cardiovascular

Increased or decreased blood pressure, tachycardia, bradycardia, faintness.

Dermatologic

Dermatitis, photosensitivity, urticaria.

Hematologic

Leukopenia, thrombocytopenia, thrombocytopenic purpura, agranulocytosis.

Gastrointestinal

Dry mouth, nausea, vomiting, jaundice.

Respiratory

Asthma, nasal stuffiness, respiratory depression (potentially fatal) and apnea (potentially fatal). (See **WARNINGS - Respiratory Depression**.)

Other

Angioneurotic edema. Neuroleptic Malignant Syndrome (potentially fatal) has also been reported. (See **WARNINGS - Neuroleptic Malignant Syndrome**.)

Paradoxical Reactions

Hyperexcitability and abnormal movements have been reported in patients following a single administration of promethazine hydrochloride injection. Consideration should be given to the discontinuation of promethazine hydrochloride injection and to the use of other drugs if these reactions occur. Respiratory depression, nightmares, delirium, and agitated behavior have also been reported in some of these patients.

OVERDOSAGE

Signs and symptoms of overdosage range from mild depression of the central nervous

system and cardiovascular system to profound hypotension, respiratory depression, unconsciousness and sudden death. Other reported reactions include hyperreflexia, hypertonia, ataxia, athetosis, and extensor-plantar reflexes (Babinski reflex).

Stimulation may be evident, especially in pediatric patients and geriatric patients. Convulsions may rarely occur. A paradoxical-type reaction has been reported in pediatric patients receiving single doses of 75 mg to 125 mg orally, characterized by hyperexcitability and nightmares.

Atropine-like signs and symptoms dry-mouth; fixed, dilated pupils; flushing; etc., as well as gastrointestinal symptoms, may occur.

Treatment

Treatment of overdose is essentially symptomatic and supportive. Only in cases of extreme overdose or individual sensitivity do vital signs, including respiration, pulse, blood pressure, temperature, and EKG, need to be monitored. Attention should be given to the reestablishment of adequate respiratory exchange through provision of a patent airway and institution of assisted or controlled ventilation. Diazepam may be used to control convulsions. Acidosis and electrolyte losses should be corrected. Note that any depressant effects of promethazine hydrochloride injection are not reversed by naloxone.

Avoid analeptics, which may cause convulsions. The treatment of choice for resulting hypotension is administration of intravenous fluids, accompanied by repositioning if indicated. In the event that vasopressors are considered for the management of severe hypotension which does not respond to intravenous fluids and repositioning, the administration of norepinephrine or phenylephrine should be considered. EPINEPHRINE SHOULD NOT BE USED, since its use in a patient with partial adrenergic blockade may further lower the blood pressure. Extrapyrimal reactions may be treated with anticholinergic antiparkinson agents, diphenhydramine, or barbiturates. Oxygen may also be administered. Limited experience with dialysis indicates that it is not helpful.

DOSAGE AND ADMINISTRATION

Important Administration Information for Adult and Pediatric Patients 2 Years of Age and Older

- The preferred route of administration of promethazine hydrochloride injection is by deep intramuscular administration (**see WARNINGS-Severe Tissue Injury, Including Gangrene**).
- Promethazine hydrochloride injection may be administered intravenously after dilution as recommended below (**see Preparation and Administration**). If pain occurs at the injection site during intravenous infusion, immediately discontinue the infusion, and evaluate for possible arterial injection or perivascular extravasation.

Allergic Conditions

The average adult dose is 25 mg. This dose may be repeated within two hours if necessary, but continued therapy, if indicated, should be via the oral route as soon as existing circumstances permit. After initiation of treatment, dosage should be adjusted to the smallest amount adequate to relieve symptoms. The average adult dose for

amelioration of allergic reactions to blood or plasma is 25 mg.

Sedation

In hospitalized adult patients, nighttime sedation may be achieved by a dose of 25 to 50 mg of promethazine hydrochloride injection.

Nausea and Vomiting

For control of nausea and vomiting, the usual adult dose is 12.5 to 25 mg, not to be repeated more frequently than every four hours. When used for control of postoperative nausea and vomiting, the dosage of analgesics and barbiturates should be reduced accordingly (see **PRECAUTIONS - Drug Interactions**).

Antiemetics should not be used in vomiting of unknown etiology in children and adolescents (see **PRECAUTIONS - Pediatric Use**).

Preoperative and Postoperative Use

As an adjunct to preoperative or postoperative medication, 25 to 50 mg of promethazine hydrochloride injection in adults may be combined with appropriately reduced doses of analgesics and atropine-like drugs as desired. Dosage of concomitant analgesic or hypnotic medication should be reduced accordingly (see **PRECAUTIONS - Drug Interactions**).

Promethazine hydrochloride is contraindicated for use in pediatric patients less than two years of age.

Obstetrics

Promethazine hydrochloride injection in doses of 50 mg will provide sedation and relieve apprehension in the early stages of labor. When labor is definitely established, 25 to 75 mg (average dose, 50 mg) promethazine hydrochloride injection may be given with an appropriately reduced dose of any desired narcotic (see **PRECAUTIONS - Drug Interactions**). If necessary, promethazine hydrochloride injection with a reduced dose of analgesic may be repeated once or twice at four-hour intervals in the course of a normal labor. A maximum total dose of 100 mg of promethazine hydrochloride injection may be administered during a 24-hour period to patients in labor.

Pediatric Patients

Promethazine hydrochloride injection is contraindicated for use in pediatric patients less than 2 years of age (see WARNINGS - Respiratory Depression). Caution should be exercised when administering promethazine hydrochloride injection to pediatric patients 2 years of age or older. It is recommended that the lowest effective dose of promethazine hydrochloride be used in pediatric patients 2 years of age and older and concomitant administration of other drugs with respiratory depressant effects be avoided (see WARNINGS - Respiratory Depression).

In pediatric patients 2 years of age and older, the dosage should not exceed half that of the suggested adult dose. As an adjunct to premedication, the suggested dose is 1.1 mg per kg of body weight in combination with an appropriately reduced dose of narcotic or barbiturate and the appropriate dose of an atropine-like drug (see **PRECAUTIONS -**

Drug Interactions). Antiemetics should not be used in vomiting of unknown etiology in pediatric patients.

Preparation and Administration Instructions for Diluted Intravenous Infusion in Adults

Inspect the solution for particulate matter and discoloration, before dilution, after dilution, and before administration. Discard the vial, ampule, or bag if particulates and/or discoloration are observed.

1. Determine the recommended dose of promethazine hydrochloride injection.
2. Aseptically withdraw the required volume from the vial or ampule and transfer it into an infusion bag.
 - Use a filter needle when withdrawing promethazine hydrochloride injection from the ampule.
3. Dilute with the recommended volume of 0.9% Sodium Chloride Injection, see Table 1. Avoid mixing and/or diluting with any other drugs or solutions other than 0.9% Sodium Chloride Injection.
4. Gently invert the infusion bag.
5. Check patency of the access site before administration.
6. Administer the diluted solution through an intravenous catheter, inserted in a large vein (preferably through a central venous catheter), over 20 to 40 minutes; for maximum infusion rates, see Table 1.
 - Do NOT administer via an intravenous catheter in the hand or wrist.
7. If pain occurs at the injection site during intravenous infusion, immediately discontinue the infusion and evaluate for possible arterial injection or perivascular extravasation.

Table 1. Preparation and Infusion Information by Adult Dose of Promethazine Hydrochloride Injection

Dose of Promethazine Hydrochloride Injection	Volume of 0.9% Sodium Chloride Injection for Dilution	Maximum Concentration of the Diluted Promethazine Hydrochloride Injection Solution	Maximum Rate of Infusion
12.5 mg	50 mL	1 mg/mL	2.5 mL/minute
25 mg	50 mL		2.5 mL/minute
50 mg	50 mL		2.5 mL/minute
75 mg	100 mL		5 mL/minute

Preparation and Administration Instructions for Diluted Intravenous Infusion to Pediatric Patients 2 Years of Age and Older

Inspect the solution for particulate matter and discoloration, before dilution, after dilution, and before administration. Discard the vial, ampule, syringe or bag if particulates and/or discoloration are observed.

1. Determine the recommended dose of promethazine hydrochloride injection for pediatric patients 2 years of age and older.
2. Aseptically withdraw the required volume from the vial or ampule and transfer it into

- an appropriately sized syringe or bag for use with an infusion pump.
- Use a filter needle when withdrawing promethazine hydrochloride injection from the ampule.
3. Dilute with the recommended volume of 0.9% Sodium Chloride Injection, see Table 2. Avoid mixing and/or diluting with any other drugs or solutions other than 0.9% Sodium Chloride Injection.
 4. Gently invert the syringe or bag.
 5. Check patency of the access site before administration.
 6. Administer the diluted solution through an intravenous catheter, inserted in a large vein (preferably through a central venous catheter) with a maximum infusion rate of 1.25 mL/minute.
 - Do NOT administer via an intravenous catheter in the hand or wrist.
 7. If pain occurs at the injection site during intravenous infusion, immediately discontinue the infusion and evaluate for possible arterial injection or perivascular extravasation.

Table 2. Preparation and Infusion Information by Pediatric Dose of Promethazine Hydrochloride Injection*

Dose of Promethazine Hydrochloride Injection	Volume of 0.9% Sodium Chloride Injection for Dilution	Maximum Concentration of the Diluted Promethazine Hydrochloride Injection Solution	Maximum Rate of Infusion
Up to 25 mg	25 mL	1 mg/mL	1.25 mL/minute
25 mg to 50 mg	50 mL		

***Avoid mixing and/or diluting with any other drugs or solutions besides 0.9% Sodium Chloride Injection**

Storage of Diluted Promethazine Hydrochloride Injection

Diluted solution may be stored up to 4 hours at room temperature, 20°C to 25°C (68°F to 77°F), or refrigerated at 2°C to 8°C (36° to 46°F), up to 24 hours.

HOW SUPPLIED

Promethazine Hydrochloride Injection, USP - 25 mg/mL, 1 mL fill in a 1 mL ampul (NDC 39822-5525-2) packaged in cartons of 25 (NDC 39822-5525-3)

Promethazine Hydrochloride Injection, USP - 50 mg/mL, 1 mL fill in a 1 mL ampul (NDC 39822-5550-5) packaged in cartons of 25 (NDC 39822-5550-6)

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Protect from light. Keep covered in carton until time of use.

Do not use if solution has developed color or contains a precipitate.

Manufactured in Germany

Distributed by:


XGen Pharmaceuticals DJB, Inc.
Big Flats, NY 14814 USA

Rev.: 12/2023

PZ-PI-04

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Promethazine Hydrochloride Injection, USP 50 mg/mL ampul label NDC 39822-5550-5



Promethazine Hydrochloride Injection, USP 50 mg/mL carton NDC 39822-5550-6



PROMETHAZINE HYDROCHLORIDE

promethazine hydrochloride injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:39822-5550
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PROMETHAZINE HYDROCHLORIDE (UNII: R61ZEH7I1I) (PROMETHAZINE - UNII:FF28EQ494)	PROMETHAZINE HYDROCHLORIDE	50 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
EDETATE SODIUM (UNII: MP1J8420LU)	0.1 mg in 1 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M)	0.01 mg in 1 mL
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	0.25 mg in 1 mL
PHENOL (UNII: 339NCG44TV)	5 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
SODIUM ACETATE (UNII: 4550K0SC9B)	
ACETIC ACID (UNII: Q40Q9N063P)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:39822-5550-6	25 in 1 CARTON	01/30/2015	
1	NDC:39822-5550-5	1 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040737	01/30/2015	

Labeler - XGen Pharmaceuticals DJB, Inc. (117380305)

Revised: 1/2024

XGen Pharmaceuticals DJB, Inc.