

MILK OF MAGNESIA- magnesium hydroxide liquid
TOP CARE (TOPCO ASSOCIATES LLC)

top milk of mag original

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- usually produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL tablespoonful contains: **magnesium 500 mg**
- avoid freezing
- store at room temperature tightly closed

Inactive ingredients

purified water, sodium hypochlorite

Questions or Comments?

1-800-540-3765

package label

NDC 36800-649-12



COMPARE TO PHILLIPS'® ORIGINAL ACTIVE INGREDIENT*

STIMULANT FREE



Milk of Magnesia®

SALINE LAXATIVE
MAGNESIUM HYDROXIDE 1200 mg
**Comfortable,
Cramp Free Relief
Sugar Free**



Scan here for more information or call 1-888-423-0139
296-19112-0 REV1220

12 FL OZ (355 mL)

ORIGINAL FLAVOR

Drug Facts

TAMPER EVIDENT: Do not use if imprinted neckband is missing or broken.

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg Saline laxative

Purpose

- Use**
- relieves occasional constipation (irregularity)
 - generally produces bowel movement in ½ to 6 hours

Warnings

- Ask a doctor before use if you have**
- a magnesium-restricted diet
 - a sudden change in bowel habits that lasts over 14 days
 - kidney disease
 - stomach pain, nausea or vomiting

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or failure to have a bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL contains: magnesium 500 mg
- avoid freezing
- store at room temperature tightly closed

Inactive ingredients

purified water, sodium hypochlorite

Questions or comments? 1-800-540-3765

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007
©TOPCO GPMA1120 QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcarebrand.com
PRODUCT OF USA

*This product is not manufactured or distributed by Bayer Healthcare LLC, distributors of Phillips'® brand Milk of Magnesia.

✓ QUALITY GUARANTEED

996-19112-0 REV GC1220



MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-649
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name			Strength	
WATER (UNII: 059QF0KO0R)				
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-649-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/18/2020	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M007	12/01/2020		

Labeler - TOP CARE (TOPCO ASSOCIATES LLC) (006935977)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(36800-649)

Revised: 11/2023

TOP CARE (TOPCO ASSOCIATES LLC)