

GERI-LANTA MAXIMUM STRENGTH- aluminum hydroxide, magnesium hydroxide, dimethicone suspension
Geri-Care Pharmaceuticals, Corp

gerilanta max

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400mg (equivalent to dried gel, USP)

Magnesium hydroxide 400 mg

Simethicone 40mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if

symptoms last more than 2 weeks

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use

- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not exceed 12 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: **magnesium 165 mg, sodium 5 mg**
- do not freeze
- store at room temperature tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package Label



Drug Facts TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

Active Ingredients (In each 5 mL teaspoonful)	Purposes
Aluminum hydroxide (equivalent to dried gel, USP) 400 mg	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

Uses relieves • heartburn • sour stomach • acid indigestion • the symptoms referred to as gas

Warnings
Ask a doctor before use if you have
• kidney disease • a magnesium-restricted diet
Ask a doctor or pharmacist before use if you are taking a prescription drug. Antacids may interact with certain prescription drugs.
Stop use and ask a doctor if symptoms last for more than 2 weeks
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before use
• adults and children 12 years of age and older: take 2 to 4 teaspoonfuls between meals as needed, at bedtime, or as directed by a doctor
• do not exceed 12 teaspoonfuls in a 24 hour period or use the maximum dosage for more than 2 weeks
• children under 12 years of age: ask a doctor

Other Information
• each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg
• do not freeze • store at room temperature tightly closed

Inactive Ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

*This product is not manufactured or distributed by the owner of the registered trademark MYLANTA®.
Distributed by
Geri-Care Pharmaceuticals
1650 63rd Street
Brooklyn, NY 11204
REV 0519 2-910-06122-1

Barcode: N 3 57896 61912 1

GERI-LANTA MAXIMUM STRENGTH

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-619
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	LEMON (Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-619-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	01/01/2000	

Labeler - Geri-Care Pharmaceuticals, Corp (611196254)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(57896-619) , pack(57896-619)