

**GOOD NEIGHBOR PHARMACY CHILDRENS IBUPROFEN- ibuprofen suspension**  
**Amerisource Bergen**

-----  
**Amerisource Bergen Children's Ibuprofen Oral Suspension Drug Facts**

**Active ingredient (in each 5 mL = 1 teaspoonful)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.

- use only enclosed measuring cup
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart		
Weight (lbs)	Age (yrs)	Dose (teaspoonful or mL)
under 2 years		ask a doctor
24-35 lbs	2-3 years	1 tsp or 5 mL
36-47 lbs	4-5 years	1 1/2 tsp or 7.5 mL
48-59 lbs	6-8 years	2 tsp or 10 mL
60-71 lbs	9-10 years	2 1/2 tsp or 12.5 mL
72-95 lbs	11 years	3 tsp or 15 mL

### Other information

- each teaspoon contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

### Inactive ingredients

anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

### Questions or comments?

1-800-719-9260

### Principal Display Panel

Compare to Children's Motrin® active ingredient

FOR AGES 2 TO 11

children's

Ibuprofen Oral Suspension

100 mg per 5 mL (teaspoonful)


pain reliever/fever reducer (NSAID)

Lasts up to 8 hours

bubble gum flavor

Alcohol Free

4 fl oz (120 mL)



# children's Ibuprofen

Oral Suspension

100 mg per 5 mL  
(teaspoonful)

pain reliever/  
fever reducer (NSAID)

NDC 24385-361-26

Compare to  
Children's Motrin®  
active ingredient\*



FOR AGES 2 TO 11

# children's Ibuprofen

Oral Suspension

100 mg per 5 mL  
(teaspoonful)

pain reliever/  
fever reducer (NSAID)

Lasts up to  
8 hours

bubble gum flavor

Alcohol Free

4 fl oz  
(120 mL)



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

### Drug Facts

#### Active ingredient Purposes

(in each 5 mL = 1 teaspoonful)  
Ibuprofen 100 mg (NSAID)\*.....Pain reliever/  
fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Uses

temporarily:  
relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache  
reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug

takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

takes more or for a longer time than directed

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

#### Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn

### Drug Facts (continued)

- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

#### Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

#### When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed measuring cup
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

### Drug Facts (continued)

#### Dosing Chart

Weight (lbs)	Age (yrs)	Dose (teaspoonful or mL)
under 2 years ask a doctor		
24-35 lbs	2-3 years	1 tsp or 5 mL
36-47 lbs	4-5 years	1 1/2 tsp or 7.5 mL
48-59 lbs	6-8 years	2 tsp or 10 mL
60-71 lbs	9-10 years	2 1/2 tsp or 12.5 mL
72-95 lbs	11 years	3 tsp or 15 mL

#### Other information

- each teaspoon contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

**Inactive ingredients** anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum


#### Questions or comments? 1-800-719-9260

\*\*Good Neighbor Pharmacy® Ibuprofen Oral Suspension 100 mg per 5 mL (teaspoonful) is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

GLUTEN FREE

Distributed By  
AmerisourceBergen  
1300 Morris Drive  
Chesterbrook, PA 19087

Visit us at  
[www.goodneighborpharmacy.com](http://www.goodneighborpharmacy.com)



ABC# 307-715



LOT NO.

EXP.

: 36626 29 C8

## GOOD NEIGHBOR PHARMACY CHILDRENS IBUPROFEN

ibuprofen suspension

### Product Information

Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:24385-361
Route of Administration	ORAL	DEA Schedule	

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (IBUPROFEN)	IBUPROFEN	100 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS CITRIC ACID	
FD&C RED NO. 40	
GLYCERIN	
HIGH FRUCTOSE CORN SYRUP	
HYPROMELLOSES	
POLYSORBATE 80	
WATER	
SODIUM BENZOATE	
SORBITOL	
XANTHAN GUM	

**Product Characteristics**

Color	PINK (light)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24385-361-26	1 in 1 CARTON		
1		120 mL in 1 BOTTLE		
2	NDC:24385-361-34	1 in 1 CARTON		
2		240 mL in 1 BOTTLE		
3	NDC:24385-361-25	1 in 1 CARTON		
3		150 mL in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074937	05/13/2002	

**Labeler** - Amerisource Bergen (007914906)