

**UNISHIELD PAIN ZAPPER- acetaminophen, aspirin, caffeine,  
salicylamide tablet**  
**Unishield**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Unishield Pain Zapper**

**Drug Facts**

**Active ingredients (in each tablet)**

Aspirin (NSAID)\* 162 mg

Caffeine 32.4 mg

Salicylamide (NSAID)\* 152 mg

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

Pain reliever/fever reducer

Adjuvant

Pain reliever/fever reducer

**Uses**

**temporarily relieved minor aches and pains associated with**

■ headaches ■ muscular aches ■ minor arthritis pain ■ back pain ■ colds ■  
toothache ■ menstrual cramps

**Temporarily reduces fever.**

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin and Salicylamide may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin or allergic reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains nonsteroidal anti-inflammatory drugs (NSAIDs), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for pain for more than 10 days unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor
- with any other pain reliever/fever reducer)

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems (such as heartburn, upset stomach or stomach pain)
- you have heart disease, liver cirrhosis, or kidney disease
- you have asthma or high blood pressure
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- anticoagulation
- diabetes

- gout
- arthritis

**Ask a doctor or pharmacist before use if you have**

- liver disease

**When using this product do not exceed recommended dose**

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- symptoms do not improve
- pain or fever gets worse
- ringing in the ears or loss of hearing occurs
- new symptoms occur

**If pregnant or breast-feeding**, ask a health care professional before use. **It is especially important not to use aspirin and salicylamide during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

**Overdose warning:**

Taking more than the recommended dose (overdose) may cause liver damage.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

■ **do not use more than directed**

**Adults and children(12 years and older):** take 2 tablets every 4 hours as needed with a full glass of water. Do not exceed 8 tablets in 24 hours or as directed by a doctor.

**Children under 12 years of age:** Ask a doctor

### ***Other information***

- store at room temperature 59°-86°F (15°-30°C)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

### ***Inactive ingredients***

FD&C yellow #6, magnesium stearate, microcrystalline cellulose, povidone, starch, stearic acid.

**Questions or Comments?** 800-480-5855

# Unishield Pain Zapper Label

Unishield

Pain Zapper

Extra Strength Pain Relief

100 Tablets/ 2 Tablets Per Packet

Acetaminophen 110 mg. Aspirin (NSAID) 162 mg

Caffeine 32.4 mg. Salicylamide (NSAID) 152 mg

Pull To Open

This Package is for Households without Young Children.

- Fast effective relief from tough headaches

- Non-drowsy formula contains caffeine

Fast Acting Advanced Formula

Tamper Evident Unit Dose Packets

The image shows a simulated view of the Unishield Pain Zapper product packaging. The front panel features the Unishield logo, the product name 'Pain Zapper', and 'Extra Strength Pain Relief'. It lists the active ingredients: Acetaminophen 110 mg, Aspirin (NSAID) 162 mg, Caffeine 32.4 mg, and Salicylamide (NSAID) 152 mg. A prominent warning states 'THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.' and includes a 'Pull To Open' instruction. The back panel contains detailed 'Drug Facts', 'Directions', and 'Other Information' sections.

UNISHIELD PAIN ZAPPER			
acetaminophen, aspirin, caffeine, salicylamide tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49314-8113
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	162 mg
SALICYLAMIDE (UNII: EM8BM710ZC) (SALICYLAMIDE - UNII:EM8BM710ZC)		SALICYLAMIDE	152 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	110 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)		CAFFEINE	32.4 mg

## Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
STARCH, CORN (UNII: O8232NY3SJ)	

## Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	FR;2
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49314-8113-3	50 in 1 BOX	03/01/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/01/2021	

**Labeler** - Unishield (790677053)

**Registrant** - Unifirst First Aid Corporation (832947092)