MEDI-PATCH WITH LIDOCAINE RX- capsaicin, lidocaine, menthol, and methyl salicylate patch

Two Hip Consulting, LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

CAPSAICIN 0.035%

LIDOCAINE 0.5%

MENTHOL 5%

METHYL SALICYLATE 20%

PURPOSE

TOPICAL ANALGESIC

TOPICAL ANESTHETIC

USES: FOR THE TEMPORARY RELIEF OF MINOR ACHES AND PAINS OF MUSCLES AND JOINTS ASSOCIATED WITH ARTHRITIS, SIMPLE BACKACHE, STRAINS, SPRAINS, MUSCLE SORENESS AND STIFFNESS.

DIRECTIONS:

USE AS DIRECTED. PRIOR TO USE, CLEAN AND DRY AFFECTED AREA. REMOVE PROTECTIVE FILM AND APPLY DIRECTLY TO AFFECTED AREA. DO NOT USE MORE THAN 3 TIMES A DAY. THOROUGHLY WASH HANDS AFTER APPLICATION. RESEAL POUCH AFTER OPENING.

Inactive Ingredients: ACRLIC ACID, ALUMINUM HYDROXIDE, CARMELLOSE SODIEM, 2-ETHYLHEXYL ACRYLATE, GLYCERIN, ISOPROPYL MYRISTATE, METHYL ACRYLATE, NONOXYNOL-30, POLYACRYLATE, POLYACRYLIC ACID, POLYSORBATE 80, SORBITAN SESQUIOLEATE, STARCH, TALC, TARTARIC ACID, TITANIUM DIOXIDE, WATER.

OTHER INFORMATION: STORE BELOW 90°F/32°C

QUESTIONS? 888 907-2833

IF CONDITION WORSENS, OR IF SYMPTOMS PERSIST FOR MORE THAN 7 DAYS OR CLEAR UP AND OCCUR AGAIN WITHIN A FEW DAYS, DISCONTINUE USE OF THIS PRODUCT AND CONSULT A DOCTOR.

IF PREGNANT OR NURSING ASK A HEALTH PROFESSIONAL BEFORE USE.

KEEP OUT OF REACH OF CHILDREN.

Warnings: For external use only. Use only as directed. Avoid contact with eyes and mucous membranes. Do not use with heating devices or pads. Do not cover or bandage tightly. If swallowed, call poison control. If contact does occur with eyes rinse with cold water and call a doctor.

DO NOT USE:

- ON WOUNDS OR DAMAGED SKIN.
- CHILDREN UNDER 12 YEARS OLD, CONSULT A DOCTOR.
- IF YOU ARE ALLERGIC TO ANY INGREDIENTS IN THIS PRODUCT.

Drug Facts

Active ingredients	Purpose
Capsaicin 0.035%	Topical Analgesic
Lidocaine 0.5%	Topical Anesthetic
Menthol 5%	Topical Analgesic
Methyl Salicylate 20%	Topical Analgesic

Uses: For the temporary relief of minor aches and pains of muscles and joints associated with arthritis, simple backache, strains, sprains, muscle soreness and stiffness.

Warnings:

For external use only.

- Use only as directed •Avoid contact with eyes and mucous membranes •Do not use with heating devices or pads• Do not bandage tightly•If swallowed, call poison control •If contact does occur with eyes, rinse with cold water and call a doctor.
- Do not use: On wounds or damaged skin. Children under 12 years old, consult a doctor. If you are allergic to any ingredients in this product.
- If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.

If pregnant or nursing ask a health professional before use. Keep out of the reach of children.

Directions: Use only as directed. Prior to use, clean & dry affected area. Remove protective film & apply directly to affected area. Do not use more than 3 times a day. Thoroughly wash hands after application. Reseal pouch after opening.

Other information: Store below 90°F/32°C

Inactive ingredients: Acrylic acid, Aluminum
Hydroxide, Carmellose Sodium, 2-Ethylhexyl Acrylate, Glycerin,
Isopropyl Myristate, Methyl Acrylate, Nonoxynol-30,
Polyacrylate, Polyacrylic Acid, Polysorbate 80, Sorbitan
Sesquioleate, Starch, Talc, Tartaric Acid, Titanium Dioxide,
Water(245-114)

Questions? 888 907-2833

Manufactured for: Medi-Sulting.com 547 Cityscape Place San Jose, CA 95136 Made in China www.medi-derm.net

NDC 76074 153 21

MEDI-PATCH WITH LIDOCAINE RX

capsaicin, lidocaine, menthol, and methyl salicylate patch

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76074-153
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety Ingredient Name Basis of Strength METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:0414PZ4LPZ) METHYL SALICYLATE 10 g MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A) MENTHOL 2.5 g

CAPSAICIN (UNII: S07O44R1ZM) (CAPSAICIN - UNII:S07O44R1ZM)	CAPSAICIN	0.0175 g
LIDO CAINE (UNII: 98 PI200987) (LIDO CAINE - UNII:98 PI200987)	LIDOCAINE	0.25 g

Inactive Ingredients		
Ingredient Name	Strength	
ACRYLIC ACID (UNII: J94PBK7X8S)		
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0)		
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)		
ETHYLHEXYL ACETATE (UNII: 2C7K8OA8SB)		
GLYCERIN (UNII: PDC6A3C0OX)		
ISOPROPYL MYRISTATE (UNII: 0 RE8 K4LNJS)		
METHYL ACRYLATE (UNII: WC487PR91H)		
NONOXYNOL-30 (UNII: JJX07DG188)		
SODIUM POLYACRYLATE (2500000 MW) (UNII: 05I15JNI2J)		
POLYACRYLIC ACID (800000 MW) (UNII: D016NSZ87U)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
SORBITAN SESQUIOLEATE (UNII: 0 W8 RRI5W5A)		
STARCH, CORN (UNII: O8232NY3SJ)		
TALC (UNII: 7SEV7J4R1U)		
TARTARIC ACID (UNII: W48881119H)		
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)		
WATER (UNII: 059QF0KO0R)		

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76074-153-41	1 in 1 POUCH		
1	NDC:76074-153-21	1 in 1 PATCH		
2	NDC:76074-153-51	5 in 1 BOX		
2	NDC:76074-153-21	1 in 1 PATCH		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/18/2012	

Labeler - Two Hip Consulting, LLC (965352896)

Registrant - Two Hip Consulting, LLC (965352896)

Revised: 9/2013 Two Hip Consulting, LLC