

COUGH AND SORE THROAT DAYTIME- acetaminophen, dextromethorphan hbr liquid **Kinray LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cough and Sore Throat Drug Facts

Active ingredients (in each 15 mL tablespoon)

Acetaminophen 500 mg

Dextromethorphan HBr 15 mg

Purpose

Pain reliever / Fever reducer

Cough suppressant

Uses

- sore throat
- headache
- minor aches and pains
- cough due to a cold
- temporarily relieves fever

Warnings

Alcohol Warning: if you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take Acetaminophen or other pain relievers/ fever reducers.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount of this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep
- if you are allergic to acetaminophen or any of the ingredients in this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

overdose warning: taking more than recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Take only as recommended - see overdose warning
- use dose cup or Tabela spoon (TBSP) every 6 hours as needed
- do not exceed 4 doses in 24-hour period

| | |
|--|---|
| Adults and children 12 years of age and over | <ul style="list-style-type: none"> • take 2 tablespoons (tbsp) 30 ml in dose cup provided every 6 hours as needed |
| Children under 12 years | <ul style="list-style-type: none"> • do not use this product in children under 12 years of age: this will provide more than the recommended dose (overdose) and may cause liver damage |

Other information

- Store at room temperature 15°- 30°C (59° - 86°F)
- Protect from freezing
- each tablespoon contains: sodium 21 mg

TAMPER EVIDENT: DO NOT USE IF PRINTED INNER CAP SEAL IS BROKEN OR

MISSING

Inactive ingredients

carboxy methyl cellulose sodium, citric acid, FD&C Blue #1, flavor, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium saccharin, sorbitol and sucrose

Questions or comments?

1-845-692-5799

Principal Display Panel



Cough & Sore Throat

Drug Facts

| Active Ingredients (in each 15 mL tablespoon) | Purpose |
|---|-------------------------------|
| Acetaminophen 500 mg..... | Pain reliever / Fever reducer |
| Dextromethorphan HBr, 15 mg..... | Cough Suppressant |

Uses

temporarily relieves: ■ sore throat ■ headache ■ minor aches and pain
■ cough due to a cold ■ temporarily reduces fever

Warnings:

Alcohol Warning: if you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take Acetaminophen or other pain relievers/ fever reducers.

Liver Warning: This product contains Acetaminophen. Severe liver damage may occur if you take:
■ more than 4 doses in 24 hours, which is the maximum daily amount of this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product ■ to make a child sleep ■ if you are allergic to acetaminophen or any of the ingredients in this product.

Ask a doctor before use if you have

■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ a sodium-restricted diet

Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin

When using this product do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when operating machinery

Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

Keep out of reach of children

overdose warning: taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions: Take only as recommended - see overdose warning ■ use dose cup or Tablespoon (TBSP) every 6 hours as needed ■ do not exceed 4 doses in 24-hours period

| | |
|--|---|
| Adults and children 12 years of age and over | take 2 tablespoons (tbsp) 30 ml in dose cup provided every 6 hours apart |
| Children under 12 years | do not use this product in children under 12 years of age: this will provide more than the recommended dose (overdose) and may cause liver damage |

Other information ■ Store at room temperature 15° - 30°C (59° - 86°F) ■ Protect from freezing ■ each tablespoons (15 mL) contains: sodium 21 mg

TAMPER EVIDENT: DO NOT USE IF PRINTED INNER CAP SEAL IS BROKEN OR MISSING.

Inactive Ingredients: carboxy methyl cellulose sodium, citric acid, FD&C Blue #1, flavor, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium saccharin, sorbitol and sucrose

Questions or comments? 1-845-692-5799

†This product is not manufactured or distributed by the owner of the registered trademark Tylenol®. All Preferred Plus Pharmacy Products are satisfaction guaranteed or your money back from retailer where purchased.

Manufactured By:

RIJ PHARMACEUTICAL CORP.

40 Commercial Avenue

Middletown, NY 10941

Distributed By: **Kinray, LLC.**

Whitestone, NY 11357

©2014 Kinray LLC. All Rights Reserved,
PREFERRED PLUS, PREFERRED PLUS++++
and the PREFERRED PLUS logo are trademarks
and/or registered trademarks of Kinray LLC.
All other marks are property of their
respective owners.

Rev. 7/13



Cough & Sore Throat

COUGH AND SORE THROAT DAYTIME

acetaminophen, dextromethorphan hbr liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:61715-076 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|--------------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 500 mg in 15 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 15 mg in 15 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311) | |
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| MENTHOL (UNII: L7T10EIP3A) | |
| POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCROSE (UNII: C151H8M554) | |

Product Characteristics

| | | | |
|-----------------|--------------|---------------------|--|
| Color | BLUE (clear) | Score | |
| Shape | | Size | |
| Flavor | MENTHOL | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:61715-076-08 | 236 mL in 1 BOTTLE | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| | | | |

Labeler - Kinray LLC (012574513)

Revised: 5/2014

Kinray LLC