UP AND UP NAPROXEN SODIUM- naproxen sodium capsule, liquid filled Target Corporation

Target Corporation Naproxen Sodium Capsules, 220 mg (NSAID) Drug Facts

Active ingredient (in each capsule)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

- or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:	 take 1 capsule every 8 to 12 hours while symptoms last for the first dose you may take 2 capsules within the first hour do not exceed 2 capsules in any 8- to 12-hour period do not exceed 3 capsules in a 24-hour period
children under 12 years:	• ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)
- do not use if printed foil under cap is broken or missing
- read all directions and warnings before use. Keep carton.

Inactive ingredients

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions?

Call 1-800-910-6874

Package/Label Principal Display Panel

naproxen sodium capsules, 220 mg (NSAID)

pain reliever/fever reducer


Compare to the active ingredient of Aleve® Liquid Gels

strength to last 12 hours

ACTUAL SIZE

LIQUID GELS** {Replace "#" with the number of liquidgels in the product}

(**LIQUID-FILLED CAPSULES)



UP AND UP NAPROXEN SODIUM

naproxen sodium capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:11673-742
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
NAPRO XEN SO DIUM (NAPRO XEN)	NAPROXEN SODIUM	220 mg	

Inactive Ingredients			
Ingredient Name	Strength		
FD&C BLUE NO. 1			
GELATIN			
GLYCERIN			
LACTIC ACID			
MANNITOL			
POLYETHYLENE GLYCOLS			
POVIDONES			
PROPYLENE GLYCOL			
WATER			
SORBITAN			
SORBITOL			

Product Characteristics			
Color	BLUE (Clear)	Score	no score
Shape	CAPSULE	Size	20 mm
Flavor		Imprint Code	742
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:11673-742-27	1 in 1 CARTON			
1		80 in 1 BOTTLE			
2	NDC:11673-742-06	1 in 1 CARTON			
2		160 in 1 BOTTLE			
3	NDC:11673-742-58	1 in 1 CARTON			
3		40 in 1 BOTTLE			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021920	01/05/2011	

Labeler - Target Corporation (006961700)

Revised: 2/2013 Target Corporation