IBUPROFEN- ibuprofen tablet, film coated WALGREEN CO.

Walgreens 44-392

Active ingredient (in each tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - menstrual cramps
 - the common cold
 - backache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen maay cause a severe allergic reaction, especially in people allergy to aspirin. Sympoms may include:

- facial swelling
- rash
- hives
- asthma (wheezing)
- shock
- skin reddening
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- vou have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- any new symptoms appear
- redness or swelling is present in the painful area
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- used by expiration date on package

Inactive ingredients

carnauba wax, cellulose, corn starch, FD&C yellow #6 aluminum lake, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or Comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Well at

Walgreens

WALGREENS PHARMACIST RECOMMENDED

NDC 0363-0392-17

Ibuprofen 200

Ibuprofen Tablets USP 200 mg / Pain Reliever / Fever Reducer (NSAID)

300

TABLETS

Compare to Motrin® IB Tablets active ingredient III


TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

50844 REV0411A39217


Walgreens Pharmacist Survey Study, Noevember 2010.

IIIThis product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Motrin® IB Tablets.

FRONT



BACK



Walgreens 44-392

IBUPROFEN

Inactive Ingredients

Product Information Product Type HUMAN OTC DRUG LABEL Item Code (Source) NDC:0363-0392 Route of Administration ORAL DEA Schedule

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IBUPRO FEN (IBUPRO FEN)	IBUPROFEN	200 mg	

Ingredient Name	Strength
CARNAUBA WAX	
FD&C YELLOW NO. 6	
HYPROMELLOSES	
LACTOSE	
MAGNESIUM STEARATE	
POLYDEXTROSE	
POLYETHYLENE GLYCOLS	
SODIUM STARCH GLYCOLATE TYPE A POTATO	
STEARIC ACID	
TITANIUM DIO XIDE	

Product Characteristics			
Color	ORANGE	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	44;392
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0392-15	1 in 1 CARTON		
1		50 in 1 BOTTLE		
2	NDC:0363-0392-08	1 in 1 CARTON		
2		24 in 1 BOTTLE		
3	NDC:0363-0392-12	1 in 1 CARTON		
3		100 in 1 BOTTLE		
4	NDC:0363-0392-14	500 in 1 BOTTLE		
5	NDC:0363-0392-16	1000 in 1 BOTTLE		
6	NDC:0363-0392-17	300 in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/30/2002	

Labeler - WALGREEN CO. (008965063)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		038154464	PACK(0363-0392)	

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		832867894	MANUFACTURE(0363-0392)	

Revised: 10/2013 WALGREEN CO.