

# **CLEAR LAX- polyethylene glycol 3350 powder, for solution** **H E B**

## **HEB Clearlax® Drug Facts**

### **Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 g (cap filled to line)

### **Purpose**

Osmotic Laxative

### **Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### **Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

### **Do not use**

if you have kidney disease, except under the advice and supervision of a doctor

### **Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug

### **When using this product**

you may have loose, watery, more frequent stools

### **Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

### **If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - use once a day
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - do not combine with starch-based thickeners used for difficulty swallowing
  - ensure that the powder is fully dissolved before drinking
  - do not drink if there are any clumps
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients**

none

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Compare to MiraLAX® active ingredient

H-E-B®

Clearlax®

Polyethylene Glycol 3350

Powder for Solution

Osmotic Laxative

- Relieves Occasional Constipation/Irregularity


• Softens Stool

Unflavored Powder

Grit Free

NET WT. 17.9 OZ (1 LB 1.9 OZ) 510g

30 ONCE-DAILY DOSES



## CLEAR LAX

polyethylene glycol 3350 powder, for solution

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-306
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

### Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-306-03	510 g in 1 BOTTLE; Type 0: Not a Combination Product	10/08/2009	
2	NDC:37808-306-01	119 g in 1 BOTTLE; Type 0: Not a Combination Product	10/08/2009	
3	NDC:37808-306-02	238 g in 1 BOTTLE; Type 0: Not a Combination Product	10/08/2009	
4	NDC:37808-306-18	153 g in 1 BOTTLE; Type 0: Not a Combination Product	08/16/2013	01/29/2020
5	NDC:37808-306-17	289 g in 1 BOTTLE; Type 0: Not a Combination Product	08/16/2013	
6	NDC:37808-306-19	595 g in 1 BOTTLE; Type 0: Not a Combination Product	07/31/2013	05/20/2021
7	NDC:37808-306-04	850 g in 1 BOTTLE; Type 0: Not a Combination Product	09/04/2014	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	10/08/2009	

**Labeler** - H E B (007924756)

Revised: 6/2023

H E B