HI VETIC- tolnaftate solution Sato Pharmaceutical Co., Ltd.

Hi Vetic Soln

Purpose Antifungal

Uses

■ for effective treatment of most athlete's foot, jock itch, and ringworm ■prevents the recurrence of most athlete's foot with daily use

Do not use

■on children under 2 years of age unless directed by a doctor

Warnings For external use only Extremely flammable. Keep away from fire or flame

When using this product

■avoid contact with the eyes

Keep our of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Stop use and ask a doctor if

- ■irritation occurs
- there is no improvement within 4 weeks for athlete's foot and ringworm, and within 2 weeks for jock itch

Directions

- ■for the tratment of athlete's foot, jock itch, and ringworm
- ■cleanse the affected area with soap and water and dry throughly
- ■apply a thin layer of the product over affected area twice daily (morning and night), or as directed by a doctor
- ■to prevent athlete's foot
- ■cleanse the feet with soap and water and dry throughly
- ■apply a thin layer of the product to the feet once or twice daily (morning and/or night)
- ■supervise children in the use of this product
- ■for athlete's foot
- ■pay special attention to spaces between the toes
- ■wear well-fitting, ventilated shoes
- change shoes and socks at least once daily
- ■for athlete's foot and ringworm, use daily for 4 weeks.

- ■for jock itch, use daily for 2 weeks.
- ■if condition persists longer, consult a doctor
- ■this product is not effective on the scalp or nails

Other information

- keep tightly closed
- ■cap bottle tightly and store at room temperature away from heat

Inactive ingredients acetone, citric acid, diethyl sebacate, fragrance, isopropyl alcohol, lauromacrogol 400, propyl gallate, water

Active ingredient

Tolnaftate 1%

hiveticsolncart.jpg

ANTIFUNGAL REMEDY (TOLNAFTATE)

hi VETIC° SOLUTION

★For your safety use only if carton is sealed.

Drug Facts

Active ingredient

Purpose

Uses

■ for effective treatment of most athlete's foot, jock itch, and ringworm ■ prevents the recurrence of most athlete's foot with daily use

Warnings
For external use only
Extremely flammable. Keep away from fire or flame

Do not use
■ on children under 2 years of age unless directed by

When using this product avoid contact with the eyes

Stop use and ask a doctor if

■ irritation occurs
■ there is no improvement within 4 weeks for athlete's foot and ringworm, and within 2 weeks for jock itch

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Directions

 To the treatment of athlete's foot, jock itch, and ringworm cleanse the affected area with scap and water and dry throughly
 apply a thin layer of the product over affected area twice daily (morning and night), or as directed by a doctor
 to prevent athlete's foot
 cleanse the feet with scap and water and dry throughly
 apply a thin layer of the product to the feet once or twice daily (morning and/or night)
 supervise children in the use of this product
 for athlete's foot
 pay special attention to spaces between the toes
 wear well-fitting, ventilated shoes
 change shoes and socks at least once daily
 for athlete's foot and ringworm, use daily for 4 weeks.
 If condition persists ingrer, consuit a doctor
 this product is not effective on the scalp or nails

 Other Information

Other information

keep tightly closed
 cap bottle tightly and store at room temperature away from heat

Inactive ingredients acetone, citric acid, diethyl sebacate, fragrance, isopropyl alcohol, lauromacrogol 400, propyl gallate, water

Manufactured by

SATO PHARMACEUTICAL CO., LTD.

1-5-27 MOTOAKASAKA MINATO-KU TOKYO, JAPAN

医薬品表示(続き)

- さい。 ■少なくとも1日1回は 靴とソックスをはき替え てください。 ■水虫、ぜにたむしの治療 にはかれる。
- には4週間毎日続けてご使用ください。 ■いんきんたむしの治療には2週間毎日続けてご使用ください。 ■症状が長期間持続する場合は医師には1月ください。 ■症状が長期間持続する場合は医師にとれば、 本剤は関す、爪には効果はありません。

その他の情報

■ふたをしっかり閉め、室温 に保存し、熱を避けてください。

その他の成分

アセトン、クエン酸、ジエチルセ バケート、香料、イソプロパノー ル、ラウロマクロゴール400、 没食子酸プロビル、精製水

Reports of serious side effects associated with use of this product can be sent to: SATO PHARMACEUTICAL_INC 20501 Earl Street, Suite 3, Tomance, CA 90503

LOT 整装配品

US-R

sato

NDC 49873-050-02

ANTIFUNGAL TOLNAFTATE) SOLUTION



水虫治療薬 (トルナフテート)



0.67 FL OZ (20 mL)

★安全のために、外箱ののり付けしたふたが開け られていた場合には、使 用しないでください。

医薬品表示

有効成分・分量 配合目的 トルナフテート1% … 抗真菌剤

効能·効果

■水虫、いんきんたむし、ぜ にたむしの治療。 ■日々 の使用による水虫の予防。

使用上の注意

外用にのみご使用ください。 可燃性。火気に近づけない でください。

使用しないでください。 ■医師の指示がない限り2 才以下の小児には使用しないでください。

使用の際、ご注意ください。 ■目に入れないようにして ください。

使用を中止し、医師にご相 談ください。

■炎症が生じた場合 ■水虫、ぜにたむしが4週間 以内、いんきんたむしが2週間 以内に改善されない場合

全ての医薬品は小児の手の 至くの医染血は小児保管している。 最かないところで使うして、 はいださまった場合には、 でしまった場合には、 に専門家又はポイズご相 にすったとなった。 トロールでください。

用法·用量

- 用ください。

HI VETIC

tolnaftate solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:49873-050

Route of Administration

TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TOLNAFTATE (UNII: 06KB629TKV) (TOLNAFTATE - UNII:06KB629TKV)	TOLNAFTATE	1 g in 100 mL

Inactive Ingredients		
Ingredient Name	Strength	
ACETONE (UNII: 1364PS73AF)		
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
DIETHYL SEBACATE (UNII: I41B9FJK6V)		
ISOPROPYL ALCOHOL (UNII: ND2M416302)		
POLIDOCANOL (UNII: 0AWH8BFG9A)		
PROPYL GALLATE (UNII: 8D4SNN7V92)		
WATER (UNII: 059QF0KO0R)		

Packaging				
t Item Code Package Description		Marketing Start Date	Marketing End Date	
NDC:49873-050- 02	1 in 1 CARTON	08/14/1990		
1	20 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M005	08/14/1990	

Labeler - Sato Pharmaceutical Co., Ltd. (690575642)

Establishment			
Name	Address	ID/FEI	Business Operations
Sato Pharmaceutical Co., Ltd.		715699133	manufacture(49873-050) , label(49873-050) , pack(49873-050)

Revised: 12/2023 Sato Pharmaceutical Co., Ltd.