ALL DAY RELIEF- naproxen sodium tablet, film coated Kroger Company

Kroger Co. All Day Relief Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	 take 1 caplet every 8 to 12 hours while symptoms last for the first dose you may take 2 caplets within the first hour do not exceed 2 caplets in any 8- to 12-hour period do not exceed 3 caplets in a 24-hour period
Children under 12 years	• ask a doctor

Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-800-632-6900

Principal Display Panel

COMPARE TO active ingredient of ALEVE® CAPLETS

See side panel

SEE NEW WARNINGS INFORMATION

Strength to Last 12 Hours

All Day Relief

Naproxen Sodium Tablets, USP 220 mg

Pain Reliever - Fever Reducer (NSAID)

Actual Size

**Capsule-Shaped Tablets





Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use, it is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Kep out of read of children. In case of Keep out of read of children. In case of Children of the control children.

while symptoms last
I not the first lobes you may take
2 caplets within the first hour
4 on ot exceed 2 caplets in any
8- to 12-hour period
6 on ot exceed 3 caplets in a
24-hour period do not take more than directed
 the smallest effective dose should be used
 drink a full glass of water with each dose ■ take 1 caplet every 8 to 12 hour


ask a doctor

Other information

each caplet contains: sodium 21 mg

store at 20-25°C (68-77°F). Avoid high humidity
and excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide



DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purposes Naproxen sodium 220 mg Pain reliever/

(naproxen 200 mg) (NSAID)*.....fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 muscular aches
 - backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
 shock

skin reddening ■ rash
■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product


- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear







ALL DAY RELIEF

naproxen sodium tablet, film coated

Product Information

Product TypeHUMAN OTC DRUG LABELItem Code (Source)NDC:30142-368Route of AdministrationORALDEA Schedule

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength
NAPRO XEN SO DIUM (NAPRO XEN) NAPRO XEN SO DIUM 220 mg

Inactive Ingredients			
Ingredient Name	Strength		
HYPROMELLOSES			
MAGNESIUM STEARATE			
CELLULO SE, MICRO CRYSTALLINE			
POLYETHYLENE GLYCOLS			
POVIDONES			
TALC			
TITANIUM DIO XIDE			

Product Characteristics			
Color	BLUE (Light Blue)	Score	no score
Shape	CAPSULE (oval)	Size	12mm
Flavor		Imprint Code	L368
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1 N	NDC:30142-368-71	1 in 1 CARTON		
1		50 in 1 BOTTLE		

2 NDC:30142-368-62	1 in 1 CARTON	
2	24 in 1 BOTTLE	
3 NDC:30142-368-78	1 in 1 CARTON	
3	100 in 1 BOTTLE	
4 NDC:30142-368-90	500 in 1 BOTTLE	
5 NDC:30142-368-76	1 in 1 CARTON	
5	120 in 1 BOTTLE	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA074661	01/16/1997		

Labeler - Kroger Company (006999528)

Revised: 12/2012 Kroger Company