GREEN GUARD IBUPRO RELIEF- ibuprofen tablet, film coated Unifirst First Aid Corporation

Green Guard – UNIFIRST FIRST AID PRODUCTS

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves minor aches and pains associated with

- headache
- toothache
- backache
- menstrual cramps
- common cold
- muscular aches
- minor arthritis pain

Temporarily reduces fever.

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- skin reddening
- asthma (wheezing)
- facial swelling
- rash
- shock
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

• take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children: (12 years and older)

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:

Do not give to children under 12 years of age.

Other information

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

carnauba wax*, cellulose*, colloidal silicon dioxide*, corn starch*, croscarmellose sodium*, hypromellose*, iron oxide red*, lactose*, magnesium stearate, microcrystalline cellulose, polydextrose*, polyethylene glycol, polyvinyl alcohol*, povidone*, pregelatinized starch*, silica*, sodium lauryl sulfate*, sodium starch glycolate*, stearic acid*, talc*, titanium dioxide, triacetate*, triacetin*

*may contain

Questions or comments? 1-800-869-6970

100R Green Guard Ibuprofen Label

Ibupro relief

Compare active ingredient to Advil®

Green Guard®

Pull Out

Ibuprofen 200 mg/Ibuprofeno 200 mg

Pain Reliever/Fever Reducer (NSAID)

Alivia el Dolor/Reduce la Fiebre (AINE)

Advil® is a Registered Trademark of Pfizer Consumer Healthcare

125 packets of 2 Tablets

250 Tablets

Order# 2251

Distributed by Green Guard®

St. Louis, MO 63045

250 Tablets 125 Packets of 2 Tablets

GHAND

CHEEN

Alivia el Dolor/Reduce la Fiebre (AINE) Pain Reliever/Fever Reducer (NSAID)

> Ibuprofeno 200mg /gm002 n5forqudi

ojdngi

Información del medicamento (continuación)

No se administre

- otro producto que alivia el dolor/reduce la fiebre
- inmediatamente antes o después de una cirugia del corazón

Consulte con un médico antes de consumirlo, si

- la advertencia acerca del sangrado estomacal debe aplicarse tiene antecedentes de problemas estomacales tales como
- acidez estomacal

 tiene la presión sanguinea alta, enfermedad del corazón,
- cirrosis del hígado o enfermedad de los riñones está tomando un diurético
- presenta problemas o graves efectos secundarios al tomar medicamentos para aliviar el dolor o para reducir la fiebre padece de asma

Consulte con su médico o farmaceuta antes de administrarlo, si usted está

- tomando aspirina a causa de un ataque al corazón o un ataque cerebral, ya que el ibuprofeno pudiera disminuir el
- beneficio de la aspirina

 tomando cualquier otra medicina
- se encuentra bajo cuidado médico debido a un estado de

Al consumir este medicamento

- pudiera aumentar el riesgo de sufrir un ataque al corazón o cerebro-vascular si consume más o por más tiempo de lo que esté indicado
- ingiera alimentos o leche si se presenta malestar estomacal

Deje de tomar este producto y consulte con el médico si se presentan cualquiera de los siguientes síntomas de sangrado estomacal:

- tiene dolor estomacal que no mejora
 el dolor empeora o perdura por más de 10 días
- la fiebre empeora o perdura por más de 3 días aparece en la parte adolorida enrojecimiento o hinchazón
- aparece cualquier sintoma nuevo

Si está embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto. Es sobre do importante que no se consuma ibuprofeno durante los

Información del medicamento (continuación)

timos 3 meses del embarazo, a menos que el médico así se lo haya especificado, ya que esto pudiera causar problemas a la criatura que está por nacer o traer complicaciones durante el parto.

Mantenga este y todos los medicamentos fuera del alcance de los niños. En caso de ocurrir una sobredosis obtenga nmediata atención médica o llame a un centro de control de envenenamiento (Poison Control Center).

Indicaciones

- no se debe exceder la dosis de consumo recomendada
- debe consumirse la menor dosis efectiva
 no debe consumirse por más de 10 días, a menos que el médico así lo señale (ver las Advertencias)

Adultos y niños: (de 12 años y mayores)

Tome 1 tableta cada 4 a 6 horas mientras persisten los síntomas. Si el dolor o la fiebre no responden con 1 tableta, puede consumir 2 tabletas. No consuma más de 6 tabletas en 24 horas, a menos que el médico lo señale.

12 años:

No se administre a niños menores de 12 años de edad.

Otra información

- lea toda la información del producto antes de consi
- almacenamiento: Consérvese a temperatura de 68-77°F (20-25°C)
- evite el calor que exceda de 104°F (superior a 40°C)
- paquetes con sellado evidente
- no use ningún paquete abierto o roto

Ingredientes inactivos cera de camauba*, celulosa do coloidal de silicio*, almidón de maíz*, croscaramelosa sódica*, hipromelosa*, óxido de hierro rojo*, lactosa*, estearato de magnesio, celulosa microcristalina, polidextrosa*, polietine[glicol, alcohol polivinilico", povidona", almidón pregelatinizado", silice" laurilsulfato sódico", almidón glicolato sódico", ácido estéarico" talco*, dióxido de titanio, triacetato*, triacetina*

¿Tiene alguna pregunta o comentario? Llam al 1-800-869-6970

CONSERVE LA CAJA PARA OBTENER LA INFORMACIÓN COMPLETA DEL PRODUCTO



Ibuprofen 200mg/Ibuprofeno 200mg


Pain Reliever/Fever Reducer (NSAID) Alivia el Dolor/Reduce la Fiebre (AINE)

Advil® is a Registered Trademark of Pfizer Consumer Healthcare

125 Packets of 2 Tablets 250 Tablets

Order #2251

Distributed by Green Guard®, St Louis, MO 63045



RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION



de que ocurra esto aumentan si

■ tiene la edad de 60 o más años

ibuprofeno, naproxeno, u otros)

■ si ha tenido úlceras estomacales o problemas de sangrado

si se consume más o por más tiempo de lo indicado

torna medicamentos para el adelgazamiento de la sangre (anticoagulantes) o esteroides
 toma otras medicinas, con o sin receta médica, que contenga un AINE (aspirina,

toma 3 o más bebidas alcohólicas diariamente mientras consume este producto.

GREEN GUARD IBUPRO RELIEF

ibuprofen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:47682-029
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Ibuprofen (Ibuprofen)	Ibupro fen	200 mg	

Inactive Ingredients			
Ingredient Name	Strength		
CELLULO SE, MICRO CRYSTALLINE			
CROSCARMELLOSE SODIUM			
MAGNESIUM STEARATE			
POLYETHYLENE GLYCOLS			

POLYVINYL ALCOHOL	
SILICON DIO XIDE	
STARCH, CORN	
TALC	
TITANIUM DIO XIDE	

Product Characteristics				
Color	white (white)	Score	no score	
Shape	ROUND (ROUND)	Size	10 mm	
Flavor		Imprint Code	115	
Contains				

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-029-48	125 in 1 BOX, UNIT-DOSE		
1	NDC:47682-029-99	2 in 1 PACKET		
2	NDC:47682-029-33	50 in 1 BOX, UNIT-DOSE		
2	NDC:47682-029-99	2 in 1 PACKET		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	06/25/2012	

GREEN GUARD IBUPRO RELIEF

ibuprofen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:47682-014
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name Basis of Strength Strength			
Ibuprofen (Ibuprofen)	Ibupro fen	200 mg	

Inactive Ingredients		
Ingredient Name	Strength	
CELLULO SE, MICRO CRYSTALLINE		
HYPROMELLOSES		
LACTOSE		
MAGNESIUM STEARATE		
POLYDEXTROSE		

POLYETHYLENE GLYCOLS	
POVIDONE	
SILICON DIO XIDE	
SODIUM STARCH GLYCOLATE TYPE A POTATO	
STARCH, CORN	
STEARIC ACID	
TITANIUM DIO XIDE	
TRIACETIN	

Product Characteristics			
Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	10 mm
Flavor		Imprint Code	44;352
Contains			

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:47682-014-48	125 in 1 BOX, UNIT-DOSE				
1	NDC:47682-014-99	2 in 1 PACKET				
2	NDC:47682-014-33	50 in 1 BOX, UNIT-DOSE				
2	NDC:47682-014-99	2 in 1 PACKET				

Marketing Information						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
ANDA	ANDA075139	12/30/2008				

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 9/2012 Unifirst First Aid Corporation