

**EQUATE NAPROXEN SODIUM- naproxen sodium tablet, film coated**  
**Wal-Mart Stores Inc**

-----  
**Wal-Mart Naproxen Sodium Tablets, 200 mg Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - it feels like the pill is stuck in your throat
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"><li>• take 1 caplet every 8 to 12 hours while symptoms last</li><li>• for the first dose you may take 2 caplets within the first hour</li><li>• do not exceed 2 caplets in any 8- to 12-hour period</li><li>• do not exceed 3 caplets in a 24-hour period</li></ul>
Children under 12 years	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

### Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

### Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

### Questions or comments?

1-888-287-1915

### Principal Display Panel

EASY OPEN CAP

Compare to Aleve® Caplets Active Ingredient

Naproxen Sodium Tablets, 220 mg

Pain Reliever/Fever Reducer (NSAID)

Strength to Last 12 Hours

PACKAGE NOT CHILD RESISTANT

220 mg

225 CAPLETS†

†Capsule Shaped Tablets

Actual Size

**Drug Facts (continued)**  
**Inactive Ingredients:** FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, titanium dioxide  
**Questions or comments?**  
1-888-281-1915  
Gluten-Free

Satisfaction guaranteed - For quantities or comments please call 1-888-281-1915

**Contains Wheat:** Some capsules contain wheat gluten. The product is not manufactured or distributed by Bayer HealthCare LLC, Ambler, PA 19004-0001. H457

EASY OPEN CAP

NDC 49035-140-83



Compare to Aleve® Capsule Active Ingredients\*\*

# Naproxen Sodium

Tablets, 220 mg  
Pain Reliever/Fever Reducer (NSAID)  
Strength to Last 12 Hours



PACKAGE NOT CHILD RESISTANT

220 mg 225 CAPSULES  
\*Capsule Shaped Tablet

Actual Size

DO NOT USE IF PRINTED FOL UNLESS CAP IS BROKEN OR MISSING

**Drug Facts**  
**Active ingredient (in each caplet):** Naproxen sodium 220 mg (naproxen 200 mg (NSAID))...over reducer  
\*Compare to Aleve® capsules and ibuprofen 200 mg

**Uses**  
Temporarily relieves minor aches and pains due to:  
■ minor pain of arthritis ■ muscular aches  
■ headache ■ menstrual cramps ■ headache  
■ backache ■ the common cold  
■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Naproxen sodium may cause a severe allergic reaction in some people. Stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs, stop use and seek medical help if an allergic reaction occurs.

**Directions**  
Take 1 or 2 caplets 2 to 4 times a day with food or milk. If stomach upset occurs, take with food or milk. If stomach upset occurs, take with food or milk. If stomach upset occurs, take with food or milk. If stomach upset occurs, take with food or milk.

**Other information**  
Keep this and all other medicines away from children. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.

**Ask a doctor or pharmacist before use if you are:**  
■ taking any other medicine for any serious condition  
■ taking any other medicine for any serious condition  
■ taking any other medicine for any serious condition

**When using this product:**  
■ Take with food or milk if stomach upset occurs.  
■ The risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

**Stop use and ask a doctor if:**  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint  
■ vomit blood  
■ have bloody or black stools  
■ have



FEEL BACK HERE ▶

14083 2E F2

### **Drug Facts** (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If **pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

(1-800-222-1222)

- Directions**
- do not take more than directed
  - the smallest effective dose should be used
  - drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years

- ask a doctor

### **Other information**

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).



# EQUATE NAPROXEN SODIUM

naproxen sodium tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG LABEL	<b>Item Code (Source)</b>	NDC:49035-140
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
NAPROXEN SODIUM (NAPROXEN)	NAPROXEN SODIUM	200 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
HYPROMELLOSES	
MAGNESIUM STEARATE	
CELLULOSE, MICROCRYSTALLINE	
POLYETHYLENE GLYCOLS	
POVIDONES	
TALC	
TITANIUM DIOXIDE	

## Product Characteristics

<b>Color</b>	BLUE (light blue)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	L368
<b>Contains</b>			

## Packaging

#	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:49035-140-83	225 in 1 BOTTLE		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA074661	08/09/2013	

**Labeler** - Wal-Mart Stores Inc (051957769)