


## SUCRALFATE- sucralfate oral suspension AvKARE

### Sucralfate Oral Suspension

#### DESCRIPTION

Sucralfate Oral Suspension contains sucralfate, USP and sucralfate is an  $\alpha$ -D-glucopyranoside,  $\beta$ -D-fructofuranosyl-, octakis-(hydrogen sulfate), aluminum complex.



Sucralfate Oral Suspension for oral administration contains 1 g of sucralfate, USP per 10 mL.

Sucralfate Oral Suspension also contains: cherry flavor, colloidal silicon dioxide, FD&C Red #40, glycerin, methylcellulose, methylparaben, microcrystalline cellulose, purified water, simethicone emulsion, and sorbitol solution. Therapeutic category: antiulcer.

#### CLINICAL PHARMACOLOGY

Sucralfate is only minimally absorbed from the gastrointestinal tract. The small amounts of the sulfated disaccharide that are absorbed are excreted primarily in the urine.

Although the mechanism of sucralfate's ability to accelerate healing of duodenal ulcers remains to be fully defined, it is known that it exerts its effect through a local, rather than systemic, action. The following observations also appear pertinent:

1. Studies in human subjects and with animal models of ulcer disease have shown that sucralfate forms an ulcer-adherent complex with proteinaceous exudate at the ulcer site.
2. *In vitro*, a sucralfate-albumin film provides a barrier to diffusion of hydrogen ions.
3. In human subjects, sucralfate given in doses recommended for ulcer therapy inhibits pepsin activity in gastric juice by 32%.

*In vitro*, sucralfate adsorbs bile salts.

These observations suggest that sucralfate's antiulcer activity is the result of formation

of an ulcer-adherent complex that covers the ulcer site and protects it against further attack by acid, pepsin, and bile salts. There are approximately 14 to 16 mEq of acid-neutralizing capacity per 1 g dose of sucralfate.

## CLINICAL TRIALS

In a multicenter, double-blind, placebo-controlled study of Sucralfate Oral Suspension, a dosage regimen of 1 g (10 mL) four times daily was demonstrated to be superior to placebo in ulcer healing.

Results from Clinical Trials Healing Rates for Acute Duodenal Ulcer				
Treatment	n	Week 2 Healing Rates	Week 4 Healing Rates	Week 8 Healing Rates
Sucralfate Oral Suspension	145	23 (16%) *	66 (46%) †	95 (66%) ‡
Placebo	147	10 (7%)	39 (27%)	58 (39%)

\* $P=0.016$  † $P=0.001$  ‡ $P=0.0001$

Equivalence of Sucralfate Oral Suspension to sucralfate tablets has not been demonstrated.

## INDICATIONS AND USAGE

Sucralfate Oral Suspension is indicated in the short-term (up to 8 weeks) treatment of active duodenal ulcer.

## CONTRAINDICATIONS

Sucralfate Oral Suspension is contraindicated for patients with known hypersensitivity reactions to the active substance or to any of the excipients.

## WARNINGS

Fatal complications, including pulmonary and cerebral emboli have occurred with inappropriate intravenous administration of Sucralfate Oral Suspension. Administer Sucralfate Oral Suspension only by the oral route. Do not administer intravenously.

## PRECAUTIONS

The physician should read the "**PRECAUTIONS**" section when considering the use of sucralfate oral suspension in pregnant or pediatric patients, or patients of childbearing potential.

Duodenal ulcer is a chronic, recurrent disease. While short-term treatment with sucralfate can result in complete healing of the ulcer, a successful course of treatment with sucralfate should not be expected to alter the post healing frequency or severity of duodenal ulceration.

Episodes of hyperglycemia have been reported in diabetic patients. Close monitoring of glycemia in diabetic patients treated with Sucralfate Oral Suspension is recommended. Adjustment of the anti-diabetic treatment dose during the use of Sucralfate Oral Suspension might be necessary.

### **Special Populations: Chronic Renal Failure and Dialysis Patients**

When sucralfate is administered orally, small amounts of aluminum are absorbed from the gastrointestinal tract. Concomitant use of sucralfate with other products that contain aluminum, such as aluminum-containing antacids, may increase the total body burden of aluminum. Patients with normal renal function receiving the recommended doses of sucralfate and aluminum-containing products adequately excrete aluminum in the urine. Patients with chronic renal failure or those receiving dialysis have impaired excretion of absorbed aluminum. In addition, aluminum does not cross dialysis membranes because it is bound to albumin and transferrin plasma proteins. Aluminum accumulation and toxicity (aluminum osteodystrophy, osteomalacia, encephalopathy) have been described in patients with renal impairment. Sucralfate should be used with caution in patients with chronic renal failure.

### **Drug Interactions**

Some studies have shown that simultaneous sucralfate administration in healthy volunteers reduced the extent of absorption (bioavailability) of single doses of the following: cimetidine, digoxin, fluoroquinolone antibiotics, ketoconazole, l-thyroxine, phenytoin, quinidine, ranitidine, tetracycline, and theophylline. Subtherapeutic prothrombin times with concomitant warfarin and sucralfate therapy have been reported in spontaneous and published case reports. However, two clinical studies have demonstrated no change in either serum warfarin concentration or prothrombin time with the addition of sucralfate to chronic warfarin therapy.

The mechanism of these interactions appears to be nonsystemic in nature, presumably resulting from sucralfate binding to the concomitant agent in the gastrointestinal tract. In all cases studied to date (cimetidine, ciprofloxacin, digoxin, norfloxacin, ofloxacin, and ranitidine), dosing the concomitant medication 2 hours before sucralfate eliminated the interaction. Due to Sucralfate Oral Suspension's potential to alter the absorption of some drugs, Sucralfate Oral Suspension should be administered separately from other drugs when alterations in bioavailability are felt to be critical. In these cases, patients should be monitored appropriately.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Chronic oral toxicity studies of 24 months' duration were conducted in mice and rats at doses up to 1 g/kg (12 times the human dose).

There was no evidence of drug-related tumorigenicity. A reproduction study in rats at doses up to 38 times the human dose did not reveal any indication of fertility impairment. Mutagenicity studies were not conducted.

### **Pregnancy**

#### **Teratogenic effects. Pregnancy Category B.**

Teratogenicity studies have been performed in mice, rats, and rabbits at doses up to 50 times the human dose and have revealed no evidence of harm to the fetus due to

sucralfate. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when sucralfate is administered to a nursing woman.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

### **Geriatric Use**

Clinical studies of Sucralfate Oral Suspension did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (see **DOSAGE AND ADMINISTRATION**).

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function (see **PRECAUTIONS Special Populations: Chronic Renal Failure and Dialysis Patients**). Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **ADVERSE REACTIONS**

Adverse reactions to sucralfate tablets in clinical trials were minor and only rarely led to discontinuation of the drug. In studies involving over 2700 patients treated with sucralfate, adverse effects were reported in 129 (4.7%).

Constipation was the most frequent complaint (2%). Other adverse effects reported in less than 0.5% of the patients are listed below by body system:

**Gastrointestinal:** diarrhea, dry mouth, flatulence, gastric discomfort, indigestion, nausea, vomiting

**Dermatological:** pruritus, rash

**Nervous System:** dizziness, insomnia, sleepiness, vertigo

**Other:** back pain, headache

**Post-marketing** cases of hypersensitivity have been reported with the use of Sucralfate Oral Suspension, including anaphylactic reactions, dyspnea, lip swelling, edema of the mouth, pharyngeal edema, pruritus, rash, swelling of the face and urticaria.

Cases of bronchospasm, laryngeal edema and respiratory tract edema have been reported with an unknown oral formulation of sucralfate.

Cases of hyperglycemia have been reported with sucralfate.

Bezoars have been reported in patients treated with sucralfate. The majority of patients had underlying medical conditions that may predispose to bezoar formation (such as delayed gastric emptying) or were receiving concomitant enteral tube feedings.

## **OVERDOSAGE**

Due to limited experience in humans with overdosage of sucralfate, no specific treatment recommendations can be given. Acute oral studies in animals, however, using doses up to 12 g/kg body weight, could not find a lethal dose. Sucralfate is only minimally absorbed from the gastrointestinal tract. Risks associated with acute overdosage should, therefore, be minimal. In rare reports describing sucralfate overdose, most patients remained asymptomatic. Those few reports where adverse events were described included symptoms of dyspepsia, abdominal pain, nausea, and vomiting.

## **DOSAGE AND ADMINISTRATION**

**Active Duodenal Ulcer.** The recommended adult oral dosage for duodenal ulcer is 1 g (10 mL) four times per day. Sucralfate Oral Suspension should be administered on an empty stomach.

Antacids may be prescribed as needed for relief of pain but should not be taken within one-half hour before or after Sucralfate Oral Suspension.

While healing with sucralfate may occur during the first week or two, treatment should be continued for 4 to 8 weeks unless healing has been demonstrated by x-ray or endoscopic examination.

**Elderly.** In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy (see **PRECAUTIONS Geriatric Use**).

Call your doctor for medical advice about side effects. You may report side effects to AvKARE at 1-855-361-3993 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## **HOW SUPPLIED**

Sucralfate Oral Suspension **1 g/10 mL** is supplied as a pink colored, cherry flavored suspension available in bottles of 420 mL (NDC 42291-781-42).

## **SHAKE WELL BEFORE USING. AVOID FREEZING.**

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Rx Only

Manufactured for:

AvKARE

Pulaski, TN 38478

Manufactured by:  
Amneal Pharmaceuticals LLC  
Branchburg, NJ 08876  
Mfg. Rev. 08-2020-01 AV 01/21

## PRINCIPAL DISPLAY PANEL

NDC 42291-781-42

**Sucralfate Oral Suspension**  
1 g/10 mL

Each 10 mL (2 teaspoons)  
Sucralfate Oral Suspension contains  
1 g sucralfate, USP.

**Rx only**    **One 420 mL Bottle**

Manufactured for: AVKARE, Pulaski, TN 38478  
Manufactured by: Amneal Pharmaceuticals LLC  
Branchburg, NJ 08876  
Mfg. Rev. 12-2019-01 AV 01/21

NDC 42291-781-42

**Sucralfate Oral Suspension**

**1 g/10 mL**

**SHAKE WELL BEFORE USING**  
**AVOID FREEZING**

**FOR ORAL ADMINISTRATION ONLY**

Each 10 mL (2 teaspoons) Sucralfate Oral Suspension contains 1 g sucralfate, USP.

**Usual Dosage:** Recommended adult dosage is 1 g (2 teaspoons) four times a day on an empty stomach. See package insert for full prescribing information.

**WARNING:** Keep out of reach of children.

**Pharmacist:** Dispense as is.

**Storage:** Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

**Rx only**                      **One 420 mL Bottle**

**AVKARE.**

1" x 2"  
Non-Varnish area for  
Lot No. and Exp. Date

3 N  
42291  
78142  
9

## SUCRALFATE

sucralfate oral suspension

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42291-781
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SUCRALFATE</b> (UNII: XX73205DH5) (SUCRALFATE - UNII:XX73205DH5)	SUCRALFATE	1 g in 10 mL

Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			20 mg in 10 mL	
METHYLCELLULOSE (4000 MPA.S) (UNII: MRJ667KA5E)			25 mg in 10 mL	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)			100 mg in 10 mL	
DIMETHICONE (UNII: 92RU3N3Y1O)			88 mg in 10 mL	
FD&C RED NO. 40 (UNII: WZB9127XOA)			0.05 mg in 10 mL	
GLYCERIN (UNII: PDC6A3C0OX)			990 mg in 10 mL	
METHYLPARABEN (UNII: A2I8C7HI9T)			20 mg in 10 mL	
SORBITOL (UNII: 506T60A25R)			2000 mg in 10 mL	
WATER (UNII: 059QF0KO0R)			7011.95 mg in 10 mL	
Product Characteristics				
Color		pink	Score	
Shape			Size	
Flavor		CHERRY	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-781-42	420 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/16/2021	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA209356	04/16/2021	

**Labeler -** AvKARE (796560394)