## UP AND UP ACETAMINOPHEN INFANTS- acetaminophen suspension Target Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### **Target Corporation Infants' Acetaminophen Drug Facts**

#### Active ingredient (in each 5 mL)

Acetaminophen 160 mg

#### **Purpose**

Pain reliever/fever reducer

#### Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

#### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if your child has ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if your child has

liver disease

#### Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

#### When using this product

do not exceed recommended dose (see overdose warning)

#### Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

#### Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

#### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

| Dosing Chart |               |              |  |
|--------------|---------------|--------------|--|
| Weight (lb)  | Age (yr)      | Dose (mL)*   |  |
| under 24     | under 2 years | ask a doctor |  |
| 24-35        | 2-3 years     | 5 mL         |  |

<sup>\*</sup> or as directed by a doctor

**Attention:** use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

#### Other information

- store at 20-25°C (68-77°F)
- · do not use if printed neckband is broken or missing

#### **Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

#### Questions or comments?

1-888-547-7400

#### Package/Label Principal Display Panel

see new warnings

Compare to active ingredient in Infants' Tylenol® Oral Suspension

value size

for ages 2 to 3 years

dye-free

infants'

acetaminophen 160 mg per 5 mL

fever reducer/pain reliever

oral suspension

no high fructose corn syrup

non-staining

ibuprofen free

alcohol free

aspirin free

Use only with enclosed syringe

See side panel for more information

SYRINGE INCLUDED

GRAPE FLAVOR

**AGES 2 TO 3 YEARS** 

4 FL OZ (118 mL)



#### UP AND UP ACETAMINOPHEN INFANTS

acetaminophen suspension

# Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:11673-133 Route of Administration ORAL

| Active Ingredient/Active Moiety   |                   |                |  |
|---|-------------------|----------------|--|
| Ingredient Name   | Basis of Strength | Strength       |  |
| ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D) | ACETAMINOPHEN     | 160 mg in 5 mL |  |

| Inactive Ingredients                      |          |
|---|----------|
| Ingredient Name                           | Strength |
| ANHYDRO US CITRIC ACID (UNII: XF417D3PSL) |          |

| CALCIUM SULFATE (UNII: WAT0 DDB 505)               |  |
|--|--|
| CARRAGEENAN (UNII: 5C69 YCD2YJ)                    |  |
| GLYCERIN (UNII: PDC6A3C0OX)                        |  |
| CELLULO SE, MICRO CRYSTALLINE (UNII: OP1R32D61U)   |  |
| CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679 OBS 311) |  |
| PROPYLENE GLYCOL (UNII: 6 DC9 Q16 7 V3)            |  |
| PROPYLPARABEN (UNII: Z8IX2SC1OH)                   |  |
| WATER (UNII: 059QF0KO0R)                           |  |
| SODIUM BENZOATE (UNII: OJ245FE5EU)                 |  |
| SORBITOL (UNII: 506T60A25R)                        |  |
| SUCRALOSE (UNII: 96K6UQ3ZD4)                       |  |
| SODIUM PHO SPHATE, TRIBASIC (UNII: A752Q30A6X)     |  |
|  |  |

| Product Characteristics |       |              |  |
|-------------------------|-------|--------------|--|
| Color                   | WHITE | Score        |  |
| Shape                   |       | Size         |  |
| Flavor                  | GRAPE | Imprint Code |  |
| Contains                |       |              |  |

| P | Packaging        |   |                             |                           |  |
|---|------------------|---|-----------------------------|---------------------------|--|
| # | Item Code        | Package Description                                   | <b>Marketing Start Date</b> | <b>Marketing End Date</b> |  |
| 1 | NDC:11673-133-16 | 1 in 1 CARTON   | 04/22/2015                  |                           |  |
| 1 |                  | 59 mL in 1 BOTTLE; Type 0: Not a Combination Product  |                             |                           |  |
| 2 | NDC:11673-133-26 | 1 in 1 CARTON   | 02/15/2018                  |                           |  |
| 2 |                  | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product |                             |                           |  |

| Marketing Information   |  |                      |                    |
|-------------------------|--|----------------------|--------------------|
| Marketing Category      | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC monograph not final | part343                                  | 04/22/2015           |                    |
|                         |  |                      |                    |

### Labeler - Target Corporation (006961700)

Revised: 5/2018 Target Corporation