IBUPROFEN- ibuprofen tablet, coated Shopko Stores Operating Co., LLC

Shopko Ibuprofen Tablets 200 mg Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen,

- or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older	 take 1 tablet every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 tablet, 2 tablets may be used do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	• ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- see end panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, iron oxides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Motrin® IB active ingredient

Ibuprofen

Tablets 200 mg

Pain Reliever/Fever Reducer (NSAID)

See New Warnings Information

Actual Size

50 Coated Tablets



SHOPKO®

**Compare to Motrin® IB active ingredient

Ibuprofen

Tablets 200 mg

Pain Reliever/Fever Reducer (NSAID)

See New Warnings Information



Actual Size

50 Coated Tablets

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, iron oxides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, stearic acid, talc, titanium dioxide

C Questions or comments? 1-800-719-9260

KEEP CARTON FOR REFERENCE

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Motrin® IB.

Distributed by Shopko Stores Operating Co., LLC, 700 Pilgrim Way Green Bay, WI 54307 ©2013 Shopko Stores Operating Co., LLC www.shopko.com

SHOPKO Satisfaction Guaranteed

By Refund Or Exchange



Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use, it is especially important not to use fulprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because if may cause problems in the unborn child or complications during delivery.

Kepp out of reach of children, in case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

do not take more than directed

the smallest effective dose should be used adults and adults and all as lake it labels are yet of hours children

12 years
and older
and older
and older
unless directed by a doctor

children under 12 years

Other information


The read all warnings and directions before use

The store at 25-5°C (68-77F)

The store at 25-5°C (68-77F)

The store at 25-5°C (68-77F)

The store at 25-5°C (104-7F)



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)

Purposes

lbuprofen 200 mg (NSAID)*.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
 toothache
 backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives facial swelling asthma (wheezing)

■ shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis. or kidney disease
- you have asthma
 you are taking a diuretic

Ask a doctor or pharmacist before use if you are


- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint vomit blood have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear





IBUPROFEN

ibuprofen tablet, coated

Product Information

 Product Type
 HUMAN OTC DRUG LABEL
 Item Code (Source)
 NDC:37012-074

 Route of Administration
 ORAL
 DEA Schedule

Active Ingredient/Active Moiety Ingredient Name Basis of Strength Strength IBUPROFEN (IBUPROFEN) BUPROFEN 200 mg

Inactive Ingredients			
Strength			

Product Characteristics						
Color	ORANGE	Score	no score			
Shape	ROUND	Size	10 mm			
Flavor		Imprint Code	I2			
Contains						

Pa	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:37012-074-71	1 in 1 CARTON					
1		50 in 1 BOTTLE					

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077349	08/19/2013	

Labeler - Shopko Stores Operating Co., LLC (023252638)

Revised: 9/2013 Shopko Stores Operating Co., LLC