

FAMOTIDINE- famotidine tablet, film coated
Valu Merchandisers Company

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Famotidine USP, 10 mg

Famotidine USP, 20 mg

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

Famotidine USP, 10 mg

- with other acid reducers
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Famotidine USP, 20 mg

- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor.
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

Famotidine USP, 10 mg

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Famotidine USP, 20 mg

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

OTHER INFORMATION

- store at 20° to 25°C (68° to 77°F)
- protect from moisture
- read the directions and warnings before use
- keep the carton. It contains important information.
- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.**
- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**

INACTIVE INGREDIENTS

Colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide red (only for Famotidine USP, 10 mg), magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PATIENT INFORMATION

- **1 tablet** relieves heartburn due to acid indigestion
- Famotidine prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn - causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

PRINCIPAL DISPLAY PANEL

COMPARE TO THE ACTIVE INGREDIENT OF ORIGINAL STRENGTH PEPCID®AC[†]

Best Choice® Brand

ORIGINAL STRENGTH

ACID CONTROLLER

FAMOTIDINE TABLETS USP, 10 mg

ACID REDUCER

Just One Tablet Prevents & Relieves Heartburn Due to Acid Indigestion

30 TABLETS

DISTRIBUTED BY: VALU MERCHANDISERS, CO.

5085657/R0411

10 mg, 30's

COMPARE TO THE ACTIVE INGREDIENT OF ORIGINAL STRENGTH PEPCID[®] AC[†]

Best Choice[®] Brand

MAXIMUM STRENGTH

ACID CONTROLLER

FAMOTIDINE TABLETS USP, 20 mg

ACID REDUCER

Just One Tablet Prevents & Relieves Heartburn Due to Acid Indigestion

25 TABLETS

DISTRIBUTED BY: VALU MERCHANDISERS, CO.

5078117/0310



ACID CONTROLLER

FAMOTIDINE TABLETS USP, 20 mg
ACID REDUCER

MAXIMUM
STRENGTH



COMPARE TO THE ACTIVE
INGREDIENT OF
MAXIMUM STRENGTH PEPCID® AC[†]

ACID CONTROLLER

FAMOTIDINE TABLETS USP, 20 mg
ACID REDUCER

MAXIMUM
STRENGTH

Just One Tablet Prevents & Relieves
Heartburn Due to Acid Indigestion

25 TABLETS



Tips for Managing Heartburn

- Do not lie flat or bend over after eating ■ Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime ■ Raise the head of your bed ■ Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals ■ If overweight, lose weight ■ Quit smoking

Expiration Date:

Batch No.

NON VARNISH

TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOLIO UNDER CAP IS OPEN OR TORN.
See end panel for batch number and expiration date.

BEST CHOICE



0 170038 161832 4



100% GUARANTEED

0310

PROUDLY DISTRIBUTED BY:
VALLI MERCHANTSERS, CO.
KANSAS CITY, MO 64111
www.avvbrands.com

■ 1 tablet relieves heartburn due to acid indigestion
■ Famotidine prevents heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Questions? call 1-800-406-7984

Inactive ingredients colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, pregelatinized starch, talc, titanium dioxide

Other information
■ read the directions and warnings before use
■ store at 20° to 25° C (68° to 77° F)
■ keep the carton. It contains important information.
■ protect from moisture

Drug Facts (continued)

Directions
■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn.
■ do not use more than 2 tablets in 24 hours
■ children under 12 years: ask a doctor


Warnings
■ Do not use ■ with other acid reducers ■ if you have kidney disease, except under the advice and supervision of a doctor ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating, or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness
Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days
Keep out of reach of children.
If pregnant or breast-feeding, ask a health professional before use.
In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Uses
■ relieves heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages
■ prevents heartburn associated with acid indigestion brought on by eating or drinking

Active ingredient (in each tablet)
Famotidine USP, 20 mg
Acid reducer

Purpose
Acid reducer



5078117



5078117

20 mg, 25's

FAMOTIDINE

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-035
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	035
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-035-31	30 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	06/25/2010	

FAMOTIDINE

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-036
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
COLLOIDAL SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	036
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-036-25	25 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090283	07/23/2010	

Labeler - Valu Merchandisers Company (868703513)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(63941-035, 63941-036)

Revised: 2/2013

Valu Merchandisers Company