

**MULTI SYMPTOM DAYTIME- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled  
Greenbrier International, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Multi-Symptom Daytime Liquid Capsules 10 softgels**

***Drug Facts***

**Active ingredients**

***Active ingredients (in each softgel)***

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purposes**

***Purposes***

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks daily while using this product

### **Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, lasts for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, see a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product, do not use more than directed.**

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed - see Overdose Warning
- do not exceed 4 doses per 24 hours

|                                     |                                     |
|-------------------------------------|-------------------------------------|
| adults & children 12 years and over | 2 Softgels with water every 4 hours |
| children 4 to under 12 years        | ask a doctor                        |
| children under 4 years              | do not use                          |

### Other information

- store at room temperature

### Inactive ingredients

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, methyl paraben, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

### Questions or comments?

1-888-952-0050

### Distributed by:

**Greenbrier International, Inc.**

**Chesapeake, VA 23320**

### PRINCIPAL DISPLAY PANEL - 10 Softgel Carton

ASSURED

COMPARE TO ACTIVE INGREDIENTS IN VICKS<sup>®</sup> DAYQUIL<sup>®</sup>

**Multi-Symptom**

**DayTime**

**Liquid Capsules**

- **Acetaminophen- Pain Reliever/Fever Reducer**
- **Dextromethorphan HBr- Cough Suppressant**
- **Phenylephrine HCl -Nasal Decongestant**

Fever, Headache, Sore Throat, Coughing, Minor Aches and Pains, & Nasal congestion

**10 Softgels**

COMPARE TO ACTIVE INGREDIENT IN VICKS DAYQUIL<sup>®</sup>

**THIS PRODUCT IS PACKAGED IN A CHILD RESISTANT AND TAMPER EVIDENT**

**PACKAGE. USE ONLY IF BLISTERS ARE INTACT.**

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

\*This product is not manufactured or distributed by Procter & Gamble distributor of Vicks® DayQuil® Cold & Flu LiquiCaps®

MULTI-SYMPTOM

ASSURED<sup>3</sup>

# DayTime Liquid Capsules

ASSURED<sup>3</sup>

COMPARE TO ACTIVE INGREDIENTS IN  
VICKS<sup>®</sup> DAYQUIL<sup>®</sup>\*

MULTI-SYMPTOM

# DayTime Liquid Capsules

- Acetaminophen - Pain Reliever / Fever Reducer
- Dextromethorphan HBr - Cough Suppressant
- Phenylephrine HCl - Nasal Decongestant

**Fever, Headache, Sore Throat,  
Coughing, Minor Aches and Pains,  
& Nasal Congestion**



Actual Size

10 softgels

Lot:

Exp:



Distributed by:  
Greenbrier International, Inc.  
Chesapeake, VA 23320

06110DTRFC

LB949  
R0811

THIS PRODUCT IS PACKAGED IN A CHILD RESISTANT AND TAMPER EVIDENT PACKAGE.  
USE ONLY IF BLISTERS ARE INTACT.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

**Drug Facts** (continued)

**Active ingredients (in each softgel) Purposes**  
 Acetaminophen 325 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr 10 mg.....Cough suppressant  
 Phenylephrine HCl 5 mg.....Nasal decongestant

**Uses**  
 temporarily relieves common cold/flu symptoms:  
 irritation, sore throat, headache, minor aches and pains, fever

**Warnings**  
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product.  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks daily while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash.  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use** ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Directions** ■ Take only as directed - see Overdose warning adults & children, even if you do not notice any signs or symptoms. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults & children.

**Other information** ■ store at room temperature

**Inactive ingredients** FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, methyl paraben, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan, titanium dioxide

**Questions or comments?** 1-888-952-0650  
 distributor of Vicks DayQuil Cold & Flu LiquidCaps<sup>®</sup>  
 This product is not manufactured or distributed by Procter & Gamble

**Drug Facts**

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.  
 When using this product, do not use more than directed.  
 Stop use and ask a doctor if  
 ■ you get nervous, dizzy or sleepless  
 ■ pain, nasal congestion or cough gets worse or lasts more than 7 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present ■ new symptoms occur  
 ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.  
 ■ if pregnant or breast-feeding, ask a health professional before use.  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks daily while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash.  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied with or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use** ■ with any other products containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema

Failure to follow these warnings could result in serious consequences.

# MULTI SYMPTOM DAYTIME

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

## Product Information

|                         |                |                    |                |
|-------------------------|----------------|--------------------|----------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:33992-6110 |
| Route of Administration | ORAL           |                    |                |

## Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength             | Strength |
|---|-------------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)                    | ACETAMINOPHEN                 | 325 mg   |
| Dextromethorphan Hydrobromide (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS) | Dextromethorphan Hydrobromide | 10 mg    |
| Phenylephrine Hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)      | Phenylephrine Hydrochloride   | 5 mg     |

## Inactive Ingredients

| Ingredient Name                         | Strength |
|---|----------|
| FD&C Red No. 40 (UNII: WZB9127XOA)      |          |
| FD&C Yellow No. 6 (UNII: H77VEI93A8)    |          |
| gelatin (UNII: 2G86QN327L)              |          |
| glycerin (UNII: PDC6A3C0OX)             |          |
| METHYLPARABEN (UNII: A2I8C7HI9T)        |          |
| polyethylene glycols (UNII: 3WJQ0SDW1A) |          |
| POVIDONES (UNII: FZ989GH94E)            |          |
| propylene glycol (UNII: 6DC9Q167V3)     |          |
| water (UNII: 059QF0K00R)                |          |
| sorbitol (UNII: 506T60A25R)             |          |
| titanium dioxide (UNII: 15FIX9V2JP)     |          |

## Product Characteristics

|          |        |              |          |
|----------|--------|--------------|----------|
| Color    | ORANGE | Score        | no score |
| Shape    | OVAL   | Size         | 21mm     |
| Flavor   |        | Imprint Code | 512      |
| Contains |        |              |          |

## Packaging

| # | Item Code        | Package Description                                     | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:33992-6110-1 | 1 in 1 CARTON   | 01/18/2018           |                    |
| 1 |                  | 10 in 1 BLISTER PACK; Type 0: Not a Combination Product |                      |                    |

## Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341                                  | 01/18/2018           |                    |

**Labeler** - Greenbrier International, Inc. (610322518)

Revised: 12/2017

Greenbrier International, Inc.