

**MILK OF MAGNESIA- magnesium hydroxide liquid**  
**Rij Pharmaceutical Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active Ingredient (in each 5 mL) One teaspoonful)**

Magnesium Hydroxide 400 mg

**Purpose**

Saline laxative/Antacid

**Use**

**Laxative**

- relieves occasional constipation (irregularity).
- generally produces bowel movement in ½ to 6 hours.

**Antacid**

- relieves acid indigestion, sour stomach, heartburn.

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- stomach pain, nausea or vomiting
- notice a sudden change in bowel habits that persists over a period of 2 weeks
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Antacid may interact with certain prescription drugs.

**When using this product**

- it may have a laxative effect
- do not exceed the maximum recommended daily dosage in a 24-hour period

**Stop use and ask a doctor if**

**Laxative**

- you have rectal bleeding or failure to have a bowel movement after use. These may indicate a serious condition.
- you need to use a laxative for more than one week

## Antacid

- symptoms last for more than 2 weeks.

### If pregnant or breast-feeding,

ask a health professional before use.

### Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

## Directions

- shake well before use

	Laxative use:	Antacid use
	Do not use dosage cup for ages under 12 years. Follow each dose with a full glass (8 oz) of liquid.	Do not use dosage cup. Follow each dose with a little water.
adults and children 12 years and older	2-4 TBSP (30-60 mL) once a day	1-3 tsp (5-15 mL) up to 4 times a day
6 to 11 years	1-2 TBSP (15-30 mL) once a day	ask a doctor
2 to 5 years	1-3 tsp (5-15 mL) once a day	ask a doctor
under 2 years	ask a doctor	ask a doctor

## Other information

- TAMPER EVIDENT: Do not use if breakaway band on cap is broken or missing.
- Store at room temperature 15° - 30°C (59° - 86°F)
- keep tightly closed
- keep from freezing
- each teaspoon (5 mL) contains: magnesium 166 mg
- each teaspoon (5 mL) contains: sodium 1 mg

## Inactive ingredients

citric acid, flavor, purified water, sodium hypochlorite.

## Principal Display Panel



NDC# 53807-401-16

Compare to the Active Ingredient in Phillip's Milk of Magnesia®†

# MILK OF Magnesia

ANTACID • LAXATIVE

For the Relief of:

- Constipation
- Acid Indigestion
- Sour Stomach
- Heartburn

MINT



16 FL. OZ (473 ml)

### Drug Facts

**Active ingredient (in each 5 ml)(One teaspoonful)** Purpose  
Magnesium Hydroxide 400 mg .....Saline laxative/Antacid

**Uses** **Laxative** ■ relieves occasional constipation (irregularity)  
■ generally produces bowel movement in 1/2 to 6 hours  
**Antacid** ■ relieves acid indigestion, sour stomach, heartburn

### Warnings

**Ask a doctor before use if you have** ■ kidney disease ■ stomach pain, nausea or vomiting ■ notice a sudden change in bowel habits that persists over a period of 2 weeks ■ a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Antacid may interact with certain prescription drugs.

**When using this product** ■ it may have a laxative effect. ■ do not exceed the maximum recommended daily dosage in a 24-hour period

### Stop use and ask a doctor if

**Laxative** ■ you have rectal bleeding or failure to have a bowel movement after use. These may indicate a serious condition  
■ you need to use a laxative for more than one week

**Antacid** ■ symptoms last for more than 2 weeks

**If pregnant or breast-feeding**, ask health professional before use  
**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center immediately.

### Directions

 ■ shake well before use

	Laxative use:	Antacid use:
	Do not use dosage cup for ages under 12 years. Follow each dose with a full glass (8 oz) of liquid.	Do not use dosage cup. Follow each dose with a little water.
adults and children 12 years and older	2-4 TBSP (30-60 mL) once a day	1-3 tsp (5-15 mL) up to 4 times a day
6 to 11 years	1-2 TBSP (15-30 mL) once a day	ask a doctor
2 to 5 years	1-3 tsp (5-15 mL) once a day	ask a doctor
under 2 years	ask a doctor	ask a doctor

**Other information** ■ TAMPER EVIDENT: Do not use if breakaway band on cap is broken or missing. ■ Store at room temperature 15°-30°C (59°-86°F) ■ keep tightly closed ■ keep from freezing ■ each teaspoon (5 mL) contains: magnesium 166 mg ■ each teaspoon (5 mL) contains: sodium 1 mg

**Inactive ingredient:** citric acid, flavor, purified water, sodium hypochlorite.

† This product is not manufactured or distributed by Bayer Corporation, owner of the registered trademark Phillip's® Milk of Magnesia. All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from retailer where purchased. L REV. 03/12

Manufactured by:  
RJ PHARMACEUTICAL CORP.  
40 Commercial Avenue  
Middletown, NY 10941  
Distributed by:  
Kinray, Inc.,  
Whitestone, NY 11357



## MILK OF MAGNESIA

magnesium hydroxide liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53807-401
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	<b>MAGNESIUM HYDROXIDE</b>	400 mg in 5 mL
---	----------------------------	-------------------

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM HYPOCHLORITE</b> (UNII: DY38VHM5OD)	

<b>Product Characteristics</b>			
<b>Color</b>	WHITE (opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	PEPPERMINT	<b>Imprint Code</b>	
<b>Contains</b>			

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53807-401-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	
2	NDC:53807-401-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/1999	

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	03/16/1999	

**Labeler** - Rij Pharmaceutical Corporation (144679156)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
Rij Pharmaceutical Corporation		144679156	manufacture(53807-401)