

ARTHRITIS PAIN RELIEF- acetaminophen tablet, film coated, extended release

Meijer Distribution Inc

Meijer Distribution, Inc. Arthritis Pain Relief Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults	<ul style="list-style-type: none">• take 2 caplets every 8 hours with water• swallow whole; do not crush, chew, split or dissolve• do not take more than 6 caplets in 24 hours• do not use for more than 10 days unless directed by a doctor
under 18 years of age	<ul style="list-style-type: none">• ask a doctor

Other information

- store at 20-25°C (68-77°F).
- meets the requirements of USP Dissolution Test 2

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Principal Display Panel

meijer®

Compare to Tylenol® 8HR Arthritis Pain active ingredient

PACKAGE NOT CHILD-RESISTANT

EASY OPEN CAP

8 HOUR arthritis pain relief

Acetaminophen

Extended-Release Tablets, 650 mg

Pain Reliever | Fever Reducer

For the Temporary Relief of Minor Arthritis Pain

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

150 Caplets* 650 mg each

*Capsule-Shaped Tablets

meijer

NDC 41250-966-47

Compare to Tylenol®
8HR Arthritis Pain
active ingredient**

PACKAGE NOT
CHILD-RESISTANT

EASY OPEN CAP

8 HOUR

arthritis
pain relief

Acetaminophen
Extended-Release Tablets, 650 mg
Pain Reliever | Fever Reducer

For the Temporary Relief of
Minor Arthritis Pain

150 Caplets*

650 mg each

*Capsule-Shaped Tablets

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN

Drug Facts

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Active ingredient (in each caplet)

Purpose

Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - premenstrual and menstrual cramps
 - the common cold
 - headache
 - toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

under 18 years
of age

- ask a doctor

Other information

- store at 20-25°C (68-77°F)
- meets the requirements of USP *Dissolution Test 2*

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, prege latinized starch, stearic acid, titanium dioxide

Questions or comments? 1-800-719-9260

Contains No Aspirin
U.S. Patent 7,897,172

**This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Tylenol®.

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com

PID 430128



9647 6E F 8

ARTHRITIS PAIN RELIEF

acetaminophen tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-966
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L544
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-966-47	150 in 1 BOTTLE; Type 0: Not a Combination Product	09/05/2007	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075077	09/05/2007	

Labeler - Meijer Distribution Inc (006959555)

