

# **DROXIDOPA- droxidopa capsule**

## **Bionpharma Inc.**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use DROXIDOPA CAPSULES safely and effectively. See full prescribing information for DROXIDOPA CAPSULES.**

**DROXIDOPA capsules, for oral use**

**Initial U.S. Approval: 2014**

#### **WARNING: SUPINE HYPERTENSION**

*See full prescribing information for complete boxed warning.*

**Monitor supine blood pressure prior to and during treatment and more frequently when increasing doses. Elevating the head of the bed lessens the risk of supine hypertension, and blood pressure should be measured in this position. If supine hypertension cannot be managed by elevation of the head of the bed, reduce or discontinue Droxidopa capsule [see Warnings and Precautions (5.1)].**

#### **INDICATIONS AND USAGE**

Droxidopa capsule is indicated for the treatment of orthostatic dizziness, lightheadedness, or the “feeling that you are about to black out” in adult patients with symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (Parkinson's disease [PD], multiple system atrophy, and pure autonomic failure), dopamine beta-hydroxylase deficiency, and non-diabetic autonomic neuropathy. Effectiveness beyond 2 weeks of treatment has not been established. The continued effectiveness of Droxidopa capsule should be assessed periodically ( 1).

#### **DOSAGE AND ADMINISTRATION**

- Starting dose is 100 mg three times during the day ( 2.1)
- Titrate by 100 mg three times daily, up to a maximum dose of 600 mg three times daily ( 2.1)
- Take consistently with or without food ( 2.1)
- To reduce the potential for supine hypertension, elevate the head of the bed and give the last dose at least 3 hours prior to bedtime ( 2.1)
- Take Droxidopa capsule whole ( 2.1)

#### **DOSAGE FORMS AND STRENGTHS**

100 mg, 200 mg, and 300 mg capsules ( 3)

#### **CONTRAINDICATIONS**

History of hypersensitivity to the drug or its ingredients ( 4)

#### **WARNINGS AND PRECAUTIONS**

- Droxidopa capsule may cause supine hypertension and may increase cardiovascular risk if supine hypertension is not well-managed ( 5.1)
- Hyperpyrexia and confusion ( 5.2)
- May exacerbate symptoms in patients with existing ischemic heart disease, arrhythmias, and congestive heart failure ( 5.3)
- Allergic reactions ( 5.4)

#### **ADVERSE REACTIONS**

- The most common adverse reactions (>5% and ≥3% compared to placebo) are headache, dizziness, nausea, and hypertension (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Bionpharma Inc. at 1-888-235-2466 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

#### **DRUG INTERACTIONS**

Use of DOPA decarboxylase inhibitors may require dose adjustments for Droxidopa capsule ( 7.2)

----- **USE IN SPECIFIC POPULATIONS** -----

- Lactation: Breastfeeding not recommended ( 8.2)
- Patients with Renal Impairment: Dosing recommendations cannot be provided for patients with GFR less than 30 mL/min ( 8.6)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 10/2019**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**WARNING: SUPINE HYPERTENSION**

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

2.1 Dosing Information

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Supine Hypertension

5.2 Hyperpyrexia and Confusion

5.3 Ischemic Heart Disease, Arrhythmias, and Congestive Heart Failure

5.4 Allergic Reactions

**6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Post marketing Experience

**7 DRUG INTERACTIONS**

7.1 Drugs that Increase Blood Pressure

7.2 Parkinson's Medications

7.3 Non-selective MAO inhibitors

**8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

**10 OVERDOSAGE**

10.1 Symptoms

10.2 Treatment

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

**14 CLINICAL STUDIES**

14.1 Studies in Neurogenic Orthostatic Hypotension

## 16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

16.2 Storage and Handling

## 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### **WARNING: SUPINE HYPERTENSION**

#### **WARNING: SUPINE HYPERTENSION**

**Monitor supine blood pressure prior to and during treatment and more frequently when increasing doses. Elevating the head of the bed lessens the risk of supine hypertension, and blood pressure should be measured in this position. If supine hypertension cannot be managed by elevation of the head of the bed, reduce or discontinue Droxidopa capsule [see *Warnings and Precautions (5.1)*].**

## 1 INDICATIONS AND USAGE

Droxidopa capsule is indicated for the treatment of orthostatic dizziness, lightheadedness, or the “feeling that you are about to black out” in adult patients with symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (Parkinson's disease [PD], multiple system atrophy, and pure autonomic failure), dopamine beta-hydroxylase deficiency, and non-diabetic autonomic neuropathy. Effectiveness beyond 2 weeks of treatment has not been established. The continued effectiveness of Droxidopa capsule should be assessed periodically.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Dosing Information

The recommended starting dose of Droxidopa capsule is 100 mg, taken orally three times daily: upon arising in the morning, at midday, and in the late afternoon at least 3 hours prior to bedtime (to reduce the potential for supine hypertension during sleep). Administer Droxidopa capsule consistently, either with food or without food. Take Droxidopa capsule whole. Titrate to symptomatic response, in increments of 100 mg three times daily every 24 to 48 hours up to a maximum dose of 600 mg three times daily (i.e., a maximum total daily dose of 1,800 mg).

Monitor supine blood pressure prior to initiating Droxidopa and after increasing the dose.

Patients who miss a dose of Droxidopa should take their next scheduled dose.

### **3 DOSAGE FORMS AND STRENGTHS**

Droxidopa capsules are available in 100 mg, 200 mg, and 300 mg strengths as specified below.

- 100 mg: Hard gelatin, size '4' capsules with opaque white body and opaque light blue cap, imprinted with "733" in black ink on cap and "100 mg" in black ink on body, containing off white to light brown powder.
- 200 mg: Hard gelatin, size '2' capsule, with an opaque white body and opaque light yellow cap, imprinted with "734" in black ink on cap and "200 mg" in black ink on body, containing off white to light brown powder.
- 300 mg: Hard gelatin, size '1' capsule, with an opaque white body and opaque light green cap, imprinted with "735" in black ink on cap and "300 mg" in black ink on body, containing off white to light brown powder.

### **4 CONTRAINDICATIONS**

Droxidopa capsule is contraindicated in patients who have a history of hypersensitivity to the drug or its ingredients [see *Warnings and Precautions (5.4)*].

### **5 WARNINGS AND PRECAUTIONS**

#### **5.1 Supine Hypertension**

Droxidopa capsule therapy may cause or exacerbate supine hypertension in patients with nOH. Patients should be advised to elevate the head of the bed when resting or sleeping. Monitor blood pressure, both in the supine position and in the recommended head-elevated sleeping position. Reduce or discontinue Droxidopa capsule if supine hypertension persists. If supine hypertension is not well-managed, Droxidopa capsule may increase the risk of cardiovascular events, particularly stroke.

#### **5.2 Hyperpyrexia and Confusion**

Postmarketing cases of a symptom complex resembling neuroleptic malignant syndrome (NMS) have been reported with Droxidopa capsule use during post marketing surveillance. Observe patients carefully when the dosage of Droxidopa capsule is changed or when concomitant levodopa is reduced abruptly or discontinued, especially if the patient is receiving neuroleptics.

NMS is an uncommon but life-threatening syndrome characterized by fever or hyperthermia, muscle rigidity, involuntary movements, altered consciousness, and mental status changes. The early diagnosis of this condition is important for the appropriate management of these patients.

#### **5.3 Ischemic Heart Disease, Arrhythmias, and Congestive Heart Failure**

Droxidopa capsule may exacerbate existing ischemic heart disease, arrhythmias, and congestive heart failure. Careful consideration should be given to this potential risk prior to initiating therapy in patients with these conditions.

#### **5.4 Allergic Reactions**

Hypersensitivity reactions including anaphylaxis, angioedema, bronchospasm, urticaria and rash have been reported in post marketing experience. Some of these reactions resulted in emergency treatment. If a hypersensitivity reaction occurs, discontinue the drug and initiate appropriate therapy.

This product contains FD&C Yellow No. 5 (tartrazine) which may also cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity [see *Contraindications (4)*].

## 6 ADVERSE REACTIONS

The following adverse reactions with Droxidopa capsule are included in more detail in the Warnings and Precautions section of the label:

- Supine Hypertension [see *Warnings and Precautions (5.1)*]
- Hyperpyrexia and Confusion [see *Warnings and Precautions (5.2)*]
- May exacerbate existing ischemic heart disease, arrhythmias, and congestive heart failure [see *Warnings and Precautions (5.3)*]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The safety evaluation of Droxidopa capsule is based on two placebo-controlled studies 1 to 2 weeks in duration (Studies 301 and 302), one 8-week placebo-controlled study (Study 306), and two long-term, open-label extension studies (Studies 303 and 304). In the placebo-controlled studies, a total of 485 patients with Parkinson's disease, multiple system atrophy, pure autonomic failure, dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy were randomized and treated, 245 with Droxidopa capsule and 240 with placebo [see *Clinical Studies (14)*].

#### Placebo-Controlled Experience

The most commonly observed adverse reactions (those occurring at an incidence of greater than 5% in the Droxidopa capsule group and with at least a 3% greater incidence in the Droxidopa capsule group than in the placebo group) in Droxidopa capsule-treated patients during the three placebo-controlled trials were headache, dizziness, nausea, and hypertension. The most common adverse reactions leading to discontinuation from Droxidopa capsule were hypertension or increased blood pressure and nausea.

**Table 1 Most Common Adverse Reactions Occurring More Frequently in the Droxidopa capsule Group**

	Study 301 and Study 302 (1 to 2 Weeks Randomized Treatment)		Study 306 (8 to 10 Weeks Randomized Treatment)	
	Placebo	Droxidopa	Placebo	Droxidopa

	(N=132) n (%)	capsule (N=131) n (%)	(N=108) n (%)	capsule (N=114) n (%)
Headache	4 (3.0)	8 (6.1)	8 (7.4)	15 (13.2)
Dizziness	2 (1.5)	5 (3.8)	5 (4.6)	11 (9.6)
Nausea	2 (1.5)	2 (1.5)	5 (4.6)	10 (8.8)
Hypertension	0	2 (1.5)	1 (0.9)	8 (7.0)

Note: n=number of patients. Adverse reactions that were reported in greater than 5% of patients in the Droxidopa capsule group and with at least a 3% greater incidence in the Droxidopa capsule group than in the placebo group were from Study 306.

#### Long-Term, Open-Label Trials with Droxidopa capsule

In the long-term, open-label extension studies, a total of 422 patients, mean age 65 years, were treated with Droxidopa capsule for a mean total exposure of approximately one year. The commonly reported adverse events were falls (24%), urinary tract infections (15%), headache (13%), syncope (13%), and dizziness (10%).

## 6.2 Post marketing Experience

The following adverse reactions have been identified during post-approval use of Droxidopa capsule. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Cardiac Disorders:* Chest pain

*Eye Disorders:* Blurred vision

*Gastrointestinal Disorders:* Pancreatitis, abdominal pain, vomiting, diarrhea

*General Disorders and Administration Site Conditions:* Fatigue

*Nervous System Disorders:* Cerebrovascular accident

*Psychiatric Disorders:* Psychosis, hallucination, delirium, agitation, memory disorder

## 7 DRUG INTERACTIONS

### 7.1 Drugs that Increase Blood Pressure

Administering Droxidopa capsule in combination with other agents that increase blood pressure (e.g., norepinephrine, ephedrine, midodrine, and triptans) would be expected to increase the risk for supine hypertension.

### 7.2 Parkinson's Medications

Dopa-decarboxylase inhibitors may require dose adjustments for Droxidopa capsule.

### 7.3 Non-selective MAO inhibitors

The concomitant use of selective MAO-B inhibitors, such as rasagiline or selegiline, was permitted in the Droxidopa capsule clinical trials. However, based on mechanism of action, the use of non-selective MAO inhibitors and linezolid should be avoided as there is a potential for increased blood pressure when taken with Droxidopa capsule.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no available data on use of Droxidopa capsule in pregnant women and risk of major birth defects or miscarriage. Droxidopa capsule did not produce significant reproductive toxicity in pregnant female rats or rabbits or in their fetuses. However, when pregnant female rats were dosed during days 7-17 of gestation (the period of fetal organogenesis) with doses of Droxidopa capsule corresponding to 0.3, 1 and 3 times the maximum recommended daily dose of 1,800 mg in a 60 kg patient, based on body surface area, and when their male and female offspring (who were exposed only during fetal life) were subsequently bred, the female offspring exhibited a dose-dependent reduction in the number of live fetuses across all three doses and an increased number of embryonic/fetal deaths at the two higher doses (see Data).

The estimated background risk of major birth defects and miscarriage in the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

#### Data

##### *Animal Data*

During a multigenerational reproductive toxicity study in rats, pregnant females were dosed during days 7-17 of gestation (the period of fetal organogenesis) with doses of Droxidopa capsule corresponding to 0.3, 1 and 3 times the maximum recommended daily dose of 1,800 mg in a 60 kg patient. Reduced weight gain, renal lesions, and a small number of deaths were observed in females treated with the two higher doses. When their male and female offspring (who were exposed to Droxidopa capsule only during fetal life) were subsequently bred, the female offspring exhibited a dose-dependent reduction in the number of live fetuses across all three doses and an increased number of embryonic/fetal deaths at the two higher doses.

### **8.2 Lactation**

#### Risk Summary

There is no information regarding the presence of Droxidopa or its active metabolite(s) in human milk, the effects of Droxidopa capsule on the breastfed child, nor the effects of Droxidopa capsule on milk production/excretion. Droxidopa is present in rat milk with peak concentrations seen 4 hours after oral drug administration and drug excretion into milk still occurring 48 hours after administration (see Data). However, due to species-specific differences in lactation physiology, animal lactation data typically do not reliably predict levels in humans. Because of the potential for serious adverse reactions, including reduced weight gain in breastfed infants, advise a woman not to breastfeed

during treatment with Droxidopa capsule.

## Data

### *Animal Data*

In rats, oral administration of droxidopa resulted in excretion into breast milk with peak concentrations seen 4 hours after administration, and excretion still occurring 48 hours after administration. When the drug was administered to nursing dams during the period of lactation at a dose corresponding to 3 times the maximum recommended daily dose of 1,800 mg in a 60 kg patient when based on body surface area, reduced weight gain and reduced survival were observed in the offspring. Despite the observed decreased weight gain, physical development was normal (with respect to timing and organ morphology).

### **8.4 Pediatric Use**

The safety and effectiveness of Droxidopa capsule in pediatric patients have not been established.

### **8.5 Geriatric Use**

A total of 197 patients with symptomatic nOH aged 75 years or above were included in the Droxidopa capsule clinical program. No overall differences in safety or effectiveness were observed between these patients and younger patients, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

### **8.6 Renal Impairment**

Droxidopa capsule and its metabolites are primarily cleared renally. Patients with mild or moderate renal impairment (GFR greater than 30 mL/min) were included in clinical trials and did not have a higher frequency of adverse reactions. Clinical experience with Droxidopa capsule in patients with severe renal function impairment (GFR less than 30 mL/min) is limited.

## **10 OVERDOSAGE**

### **10.1 Symptoms**


There have been cases of overdose reported during postmarketing surveillance. A patient ingested 7,700 mg of droxidopa and experienced a hypertensive crisis that resolved promptly with treatment. Another patient treated with a total daily dose of 2,700 mg of Droxidopa capsule experienced hypertension and an intracranial hemorrhage.

### **10.2 Treatment**

There is no known antidote for Droxidopa capsules overdose. In case of an overdose that may result in an excessively high blood pressure, discontinue Droxidopa capsule and treat with appropriate symptomatic and supportive therapy. Counsel patients to remain in a standing or seated position until their blood pressure drops below an acceptable limit.

## 11 DESCRIPTION

Droxidopa capsules contain droxidopa, which is a synthetic amino acid precursor of norepinephrine, for oral administration. Chemically, droxidopa is (-)-threo-3-(3,4-Dihydroxyphenyl)-L-serine. It has the following structural formula:



Droxidopa is an odorless, tasteless, white to off-white crystals or crystalline powder. It is slightly soluble in water, and practically insoluble in methanol, glacial acetic acid, ethanol, acetone, ether, and chloroform. It is soluble in dilute hydrochloric acid. It has a molecular weight of 213.19 and a molecular formula of  $C_9H_{11}NO_5$ .

Droxidopa capsules also contain the following inactive ingredients: mannitol, corn starch, and magnesium stearate. The capsule shell is printed with black ink. The black inks contain shellac, dehydrated alcohol, isopropyl alcohol, butyl alcohol, propylene glycol, strong ammonia solution, black iron oxide, potassium hydroxide and purified water. The capsule shell contains the following inactive ingredients: 100 mg – gelatin, titanium dioxide, FD&C Blue No. 2, black iron oxide, red iron oxide, water and SLS; 200 mg – gelatin, titanium dioxide, FD&C Blue No. 2, black iron oxide, iron oxide yellow, water and SLS; 300 mg – gelatin, titanium dioxide, FD&C Blue No. 1, FD&C Yellow No.,5 (tartrazine), FD&C Red No. 40, water and SLS. Droxidopa capsules differ in size and color by strength [see *Dosage Forms and Strengths (3)*].

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The exact mechanism of action of Droxidopa capsule in the treatment of neurogenic orthostatic hypotension is unknown. Droxidopa is a synthetic amino acid analog that is directly metabolized to norepinephrine by dopa-decarboxylase, which is extensively distributed throughout the body. Droxidopa is believed to exert its pharmacological effects through norepinephrine and not through the parent molecule or other metabolites. Norepinephrine increases blood pressure by inducing peripheral arterial and venous vasoconstriction. Droxidopa in humans induces small and transient rises in plasma norepinephrine.

### 12.2 Pharmacodynamics

Peak droxidopa plasma concentrations are associated with increases in systolic and diastolic blood pressures. Droxidopa has no clinically significant effect on standing or supine heart rates in patients with autonomic failure.

#### Cardiac Electrophysiology

No prolongation of the QTc interval was observed with Droxidopa capsule at single oral

doses up to 2,000 mg, as shown in a dedicated thorough QT study.

## **12.3 Pharmacokinetics**

### Absorption

Peak plasma concentrations ( $C_{max}$ ) of droxidopa were reached by 1 to 4 hours post-dose (mean of approximately 2 hours) in healthy volunteers. High-fat meals have a moderate impact on droxidopa exposure with  $C_{max}$  and area under the plasma concentration-time curve (AUC) decreasing by 35% and 20%, respectively. The  $C_{max}$  was delayed by approximately 2 hours with a high-fat meal.

### Distribution

Pre-clinical studies suggest that droxidopa can cross the blood-brain barrier. Droxidopa exhibits plasma protein binding of 75% at 100 ng/mL and 26% at 10,000 ng/mL. The estimated apparent volume of distribution of droxidopa is about 200 L in humans.

### Elimination

The total clearance of droxidopa after oral administration (CL/F) was approximately 400 mL/hr following administration of a single 300 mg dose.

### Metabolism

The metabolism of droxidopa is mediated by catecholamine pathway and not through the cytochrome P450 system. Droxidopa is initially converted to methoxylated dihydroxyphenylserine (3-OM-DOPS), a major metabolite, by catechol-O-methyltransferase (COMT), to norepinephrine by DOPA decarboxylase (DDC), or to protocatechualdehyde by DOPS aldolase. After oral dosing in humans, plasma norepinephrine levels peak within 3 to 4 hours but are generally very low (less than 1 ng/mL) and variable with no consistent relationship with dose. The contribution of the metabolites of droxidopa other than norepinephrine to its pharmacological effects is not well understood.

### Excretion

The mean elimination half-life of droxidopa is approximately 2.5 hours in humans. The major route of elimination of droxidopa and its metabolites is via the kidneys in both animals and in humans. Studies in animals with radiolabeled drug showed that ~75% of the administered radioactivity was excreted in urine within 24 hours of oral dosing.

### Specific Populations

There are no clinically relevant effects of age, body mass index, or sex on the pharmacokinetics of droxidopa. A population pharmacokinetic analysis suggests that hepatic function, assessed by aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase, and total bilirubin, did not influence the exposure to droxidopa. The controlled clinical trials included patients with mild to moderate renal impairment. No dose adjustments are required in patients with mild to moderate renal impairment.

### Drug Interaction Studies

No dedicated drug-drug interaction studies were performed for droxidopa. Patients in the Phase 3 trials with Droxidopa capsule received concomitant levodopa/carbidopa, dopamine agonists, MAO-B inhibitors, COMT inhibitors and other medications used to

treat Parkinson's disease. Carbidopa, a peripheral dopa-decarboxylase inhibitor, could prevent the conversion of Droxidopa to norepinephrine outside of the central nervous system (CNS). Patients taking Droxidopa capsule with L-DOPA/dopa-decarboxylase inhibitor combination drugs had decreased clearance of Droxidopa, an increase in overall exposure (AUC) to droxidopa of approximately 100%, and an increase in overall exposure to 3-OM-DOPS of approximately 50%. However, in clinical trials, it was found that the decreased clearance was not associated with a significant need for a different treatment dose or increases in associated adverse events. Dopamine agonists, amantadine derivatives, and MAO-B inhibitors do not appear to affect Droxidopa clearance, and no dose adjustments are required.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies have been conducted at dosages up to 1,000 mg/kg/day in mice and up to 100 mg/kg/day in rats with no indication of carcinogenic effects. Based on dose per unit body surface area, these two doses correspond to approximately 3 and 0.5 times, respectively, the maximum recommended total daily dose of 1,800 mg in a 60 kg patient. Droxidopa was clastogenic in Chinese hamster ovary cells (chromosome aberration assay), but was not mutagenic in bacteria (Ames assay), and was not clastogenic in a mouse micronucleus assay.

Studies in rats show that droxidopa has no effect on fertility.

### **13.2 Animal Toxicology and/or Pharmacology**

In long-term chronic toxicity studies, rats and mice treated for 52 and 80 weeks, respectively, at doses up to 300 mg/kg/day in rats and 1,000 mg/kg/day in mice had increased incidences of renal and cardiac lesions (rats and mice) and deaths (rats only). The doses at which these effects were not seen represented 0.2 and 0.3 times, in rats and mice, respectively, the maximum recommended total daily dose of 1,800 mg in a 60 kg patient, when based on body surface area.

No signs of toxicity were observed in monkeys or dogs given droxidopa for 13 weeks at doses 32 times (3,000 mg/kg/day) and 37 times (2,000 mg/kg/day), respectively, the maximum human dose.

## **14 CLINICAL STUDIES**

### **14.1 Studies in Neurogenic Orthostatic Hypotension**

Clinical studies (described below) examined the efficacy of Droxidopa capsule in the short-term (1 to 2 weeks) and over longer-term periods (8 weeks; 3 months). Studies 301 and 306B showed a treatment effect of Droxidopa capsule at Week 1, but none of the studies demonstrated continued efficacy beyond 2 weeks of treatment.

Study 306B was a multi-center, double-blind, randomized, placebo-controlled, parallel-group study in patients with symptomatic nOH and Parkinson's disease. Patients entering the study were required to have a decrease of at least 20 mm Hg or 10 mm Hg, respectively, in systolic or diastolic blood pressure, within 3 minutes after standing,

as well as symptoms associated with neurogenic orthostatic hypotension. The study had an initial dose titration period that lasted up to 2 weeks in which patients received placebo or 100 to 600 mg of Droxidopa capsule three times daily, followed by an 8-week treatment period.


Efficacy was measured using the OHSA Item #1 score (“dizziness, lightheadedness, feeling faint, and feeling like you might black out”) at Week 1, in patients who had completed titration and 1 week of maintenance therapy.

A total of 171 patients were enrolled, and 147 patients were included in the efficacy analysis. The mean age was 72 years, and patients were mostly Caucasian. During the study, 94% of placebo-treated patients and 88% on Droxidopa capsule were taking dopa-decarboxylase inhibitors; 17% of placebo-treated patients and 26% on Droxidopa capsule were taking fludrocortisone. There were more premature discontinuations in the Droxidopa capsule group (28%) than in the placebo group (20%).

In both groups, the mean baseline dizziness score was 5.1 on an 11-point scale. At Week 1, patients showed a statistically significant mean 0.9 unit decrease in dizziness with Droxidopa capsule versus placebo ( $P=0.028$ ), but the effect did not persist beyond Week 1. The data at all time points are shown in Figure 1.

Patients receiving Droxidopa capsule also had a greater increase, compared to placebo, in the Week 1 lowest standing systolic blood pressure within 3 minutes after standing (5.6 mm Hg;  $P=0.032$ ).


**Figure 1. Mean Change in OHSA Item 1 Score by Week in Study 306B**



Note: The graph is based on observed data only. The error bars are the 95% confidence interval of the mean change from baseline in OHSA Item 1 scores.

**Figure 2. Distribution of Patients by Change in OHSA Item 1, Baseline to Week 1, in Study 306B**

Figure 2 shows the distribution of changes from Baseline to Week 1 in the OHSA Item #1 score. Overall, the figure shows that patients treated with Droxidopa capsule improved more than those treated with placebo.



Study 301 was a multicenter, multinational, double-blind, randomized, placebo-controlled, parallel-group study in patients with symptomatic neurogenic orthostatic hypotension. The study included an initial open-label dose titration period, a 7-day washout period, and a randomized double-blind 7-day treatment period. To be eligible for enrollment, patients were required to have a decrease in systolic or diastolic blood pressure of at least 20 or 10 mm Hg, respectively, within 3 minutes after standing. The study was enriched, such that only patients who had been identified as “responders” during the titration period were randomized to Droxidopa capsule or placebo. To be considered a responder, a patient had to demonstrate improvement on the OHSA Item #1 score by at least 1 point, as well as an increase in systolic blood pressure of at least 10 mm Hg post-standing, during the open-label dose titration period. Patients who dropped out during the titration period because of side effects or other reasons were also not included in the double-blind portion of the study.

Patients had a primary diagnosis of Parkinson’s disease (n=60), pure autonomic failure (n=36), or multiple system atrophy (n=26). The mean age was 60 years, and most were Caucasian. 45% of patients were taking dopadecarboxylase inhibitors, and 29% were taking fludrocortisone.

Efficacy was measured using the Orthostatic Hypotension Questionnaire (OHQ), a patient-reported outcome that measures symptoms of nOH and their impact on the patient’s ability to perform daily activities that require standing and walking. The OHQ includes OHSA Item #1 as one of several components. A statistically significant treatment effect was not demonstrated on OHQ (treatment effect of 0.4 unit, P=0.19).

The mean baseline dizziness score on OHSA Item #1 (“dizziness, lightheadedness, feeling faint, and feeling like you might black out”) was 5.2 units on an 11-point scale. At Week 1 of treatment, patients showed a mean 0.7 unit decrease in dizziness with Droxidopa capsule versus placebo (P=0.06).

Study 302 (n=101) was a placebo-controlled, 2-week randomized withdrawal study of Droxidopa capsule in patients with symptomatic nOH. Study 303 (n=75) was an extension of Studies 301 and 302, where patients received their titrated dose of Droxidopa capsule for 3 months and then entered a 2-week randomized withdrawal phase. Neither study showed a statistically significant difference between treatment arms on its primary endpoint. Considering these data, the effectiveness of Droxidopa

capsule beyond 2 weeks is uncertain, and patients should be evaluated periodically to determine whether Droxidopa capsule is continuing to provide a benefit.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **16.1 How Supplied**

Droxidopa capsules are supplied in the following dosage strengths:

100 mg: Hard gelatin, size '4' capsules with opaque white body and opaque light blue cap, imprinted with "733" in black ink on cap and "100 mg" in black ink on body, containing off white to light brown powder.

200 mg: Hard gelatin, size 2 capsule, with an opaque white body and opaque light yellow cap, imprinted with "734" in black ink on cap and "200 mg" in black ink on body, containing off white to light brown powder.

300 mg: Hard gelatin, size 1 capsule, with an opaque white body and opaque light green cap, imprinted with "735" in black ink on cap and "300 mg" in black ink on body, containing off white to light brown powder

100 mg 90-count bottle (NDC code# 69452-256-19)

200 mg 90-count bottle (NDC code# 69452-257-19)

300 mg 90-count bottle (NDC code# 69452-258-19)

Dispense in a tight, Light-resistant container. Do not use if the printed seal under the cap is broken or missing.

### **16.2 Storage and Handling**

Droxidopa capsules should be stored at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see *USP Controlled Room Temperature*].

## **17 PATIENT COUNSELING INFORMATION**

### *Elevations in Blood Pressure*

Counsel patients that Droxidopa capsule causes elevations in blood pressure and increases the risk of supine hypertension, which could lead to strokes, heart attacks, and death. Instruct patients to rest and sleep in an upper-body elevated position and monitor blood pressure. Instruct patients how to manage observed blood pressure elevations. To reduce the risk of supine hypertension, in addition to raising the upper body, the late afternoon dose of Droxidopa capsule should be taken at least three hours before bedtime [see *Warnings and Precautions (5.1)*].

### *Concomitant Treatments*

Counsel patients about the concomitant use of drugs to treat other conditions that may have an additive effect with Droxidopa capsule [see *Drug Interactions (7)*].

### *Allergic Reactions*

Counsel patients to discontinue Droxidopa capsule and seek immediate medical attention

if any signs or symptoms of a hypersensitivity reaction such as anaphylaxis, angioedema, bronchospasm, urticaria or rash occur [see *Warnings and Precautions* (5.4)].

### Lactation

Advise women not to breastfeed during treatment with Droxidopa capsule [see *Use in Specific Populations* (8.2)].

### Food

Patients should take Droxidopa capsule the same way each time, either with food or without food [see *Dosage and Administration* (2.1)].

### Missed Dose

If a dose is missed, patients should take the next dose at the regularly scheduled time and should not double the dose.

Distributed by:

Bionpharma Inc.

600 Alexander Road,

Princeton, NJ 08540.

Revised: 10/2019

## Container Label for 100 mg, 200 mg and 300 mg Capsule

<p>NDC 69452-256-19</p> <h1>Droxidopa Capsules</h1> <p><b>100 mg</b></p> <p><b>Rx Only</b></p> <hr/> <p>90 Capsules <b>BIONPHARMA</b></p>	<p>Each capsule contains 100 mg of droxidopa.</p> <p><b>Usual Dosage:</b> See Prescribing information.</p> <p>Store at 20°C to 25°C (68°F to 77°F); Excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].</p> <p>Dispense in a tight, light-resistant container. Do not use if the printed seal under the cap is broken or missing.</p> <p><b>Warning:</b> Keep Out of Reach of Children.</p> <p><b>Manufactured For:</b> Bionpharma Inc. 600 Alexander Road, Princeton, NJ 08540</p> <p>Rev. 12/2018</p>	 <p>N 3 69452 25619 0</p>
---	--	--

NDC 69452-257-19

# Droxidopa Capsules

200 mg

Rx Only

90 Capsules **BIONPHARMA**

Each capsule contains 200 mg of droxidopa.

**Usual Dosage:** See Prescribing information.

Store at 20°C to 25°C (68°F to 77°F);  
Excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Dispense in a tight, light-resistant container.  
Do not use if the printed seal under the cap is broken or missing.

**Warning:** Keep Out of Reach of Children.

**Manufactured For:**

Bionpharma Inc.  
600 Alexander Road,  
Princeton, NJ 08540



Rev. 12/2018  
N 3 69452 25719 7

NDC 69452-258-19

# Droxidopa Capsules

300 mg

Rx Only

90 Capsules **BIONPHARMA**

Each capsule contains 300 mg of droxidopa.

**Usual Dosage:** See Prescribing Information.

Store at 20°C to 25°C (68°F to 77°F);  
Excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

Dispense in a tight, light-resistant container.  
Do not use if the printed seal under the cap is broken or missing.

**Warning:** Contains FD&C Yellow 5

(tartrazine) as color additive.

Keep Out of Reach of Children.

**Manufactured For:**

Bionpharma Inc.  
600 Alexander Road,  
Princeton, NJ 08540



Rev. 04/2019  
N 3 69452 25819 4

## DROXIDOPA

droxidopa capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69452-256
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DROXIDOPA (UNII: J7A92W69L7) (DROXIDOPA - UNII:J7A92W69L7)	DROXIDOPA	100 mg

### Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SHELLAC (UNII: 46N107B710)	

<b>ALCOHOL</b> (UNII: 3K9958V90M)
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)
<b>BUTYL ALCOHOL</b> (UNII: 8PJ61P6TS3)
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>AMMONIA</b> (UNII: 5138Q19F1X)
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)
<b>WATER</b> (UNII: 059QF0K00R)
<b>GELATIN</b> (UNII: 2G86QN327L)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)

### Product Characteristics

<b>Color</b>	white (opaque white body) , blue (opaque light blue cap)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	733;100mg
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69452-256-19	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA213033	10/01/2021	

## DROXIDOPA

droxidopa capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69452-257
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DROXIDOPA</b> (UNII: J7A92W69L7) (DROXIDOPA - UNII:J7A92W69L7)	DROXIDOPA	200 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>BUTYL ALCOHOL</b> (UNII: 8PJ61P6TS3)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>AMMONIA</b> (UNII: 5138Q19F1X)	
<b>FERROSO FERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	

## Product Characteristics

<b>Color</b>	white (opaque white body) , yellow (opaque light yellow cap)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	734;200mg
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69452-257-19	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2021	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA213033	10/01/2021	

## DROXIDOPA

droxidopa capsule

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:69452-258
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DROXIDOPA</b> (UNII: J7A92W69L7) (DROXIDOPA - UNII:J7A92W69L7)	DROXIDOPA	300 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>BUTYL ALCOHOL</b> (UNII: 8PJ61P6TS3)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>AMMONIA</b> (UNII: 5138Q19F1X)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	

**Product Characteristics**

<b>Color</b>	white (opaque white body) , green (opaque light green cap)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	735;300mg
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69452-258-19	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2021	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA213033	10/01/2021	

**Labeler** - Bionpharma Inc. (079637826)

**Registrant** - Bionpharma Inc. (079637826)

**Establishment**

Name	Address	ID/FEI	Business Operations
Appco Pharma LLC (Unit-II)		078510186	manufacture(69452-256, 69452-257, 69452-258)

Revised: 6/2021

Bionpharma Inc.