

**THIAMINE HCL- thiamine hcl injection, solution**  
**HF Acquisition Co LLC, DBA HealthFirst**

-----  
**THIAMINE HCl INJECTION, USP 200 mg/2 mL (100 mg/mL) 2mL VIAL**


**DESCRIPTION**

Thiamine Hydrochloride Injection, USP is a sterile solution of thiamine hydrochloride in Water for Injection for intramuscular (IM) or slow intravenous (IV) administration.

Each mL contains: Thiamine hydrochloride 100 mg; chlorobutanol anhydrous (chloral derivative) 0.5%; monothioglycerol 0.5%; water for injection, q.s. Sodium hydroxide may have been added for pH adjustment (2.5 to 4.5).

Thiamine hydrochloride, or vitamin B 1, occurs as white crystals or crystalline powder that usually has a slight characteristic odor. Freely soluble in water; soluble in glycerin; slightly soluble in alcohol; insoluble in ether and benzene. Thiamine is rapidly destroyed in neutral or alkaline solutions but is stable in the dry state. It is reasonably stable to heat in acid solution.

The chemical name of thiamine hydrochloride is thiazolium,3-[(4-amino-2-methyl-5-pyrimidinyl)methyl]-5-(2-hydroxyethyl)-4-methylchloride, monohydrochloride and it has the following structural formula:



C 12H 17ClN 4OS • HCl M.W. 337.27

## CLINICAL PHARMACOLOGY

The water soluble vitamins are widely distributed in both plants and animals. They are absorbed in man by both diffusion and active transport mechanisms. These vitamins are structurally diverse (derivatives of sugar, pyridine, purines, pyrimidine, organic acid complexes and nucleotide complex) and act as coenzymes, as oxidation-reduction agents, possibly as mitochondrial agents. Metabolism is rapid, and the excess is excreted in the urine.

Thiamine is distributed in all tissues. The highest concentrations occur in liver, brain, kidney and heart. When thiamine intake is greatly in excess of need, tissue stores increase two to three times. If intake is insufficient, tissues become depleted of their vitamin content. Absorption of thiamine following IM administration is rapid and complete.

Thiamine combines with adenosine triphosphate (ATP) to form thiamine pyrophosphate, also known as cocarboxylase, a coenzyme. Its role in carbohydrate metabolism is the

decarboxylation of pyruvic acid in the blood and  $\alpha$ -ketoacids to acetaldehyde and carbon dioxide. Increased levels of pyruvic acid in the blood indicate vitamin B 1 deficiency.

The requirement for thiamine is greater when the carbohydrate content of the diet is raised. Body depletion of vitamin B 1 can occur after approximately three weeks of total absence of thiamine in the diet.

## **INDICATIONS**

Thiamine hydrochloride injection is effective for the treatment of thiamine deficiency or beriberi whether of the dry (major symptoms related to the nervous system) or wet (major symptoms related to the cardiovascular system) variety. Thiamine hydrochloride injection should be used where rapid restoration of thiamine is necessary, as in Wernicke's encephalopathy, infantile beriberi with acute collapse, cardiovascular disease due to thiamine deficiency, or neuritis of pregnancy if vomiting is severe. It is also indicated when giving IV dextrose to individuals with marginal thiamine status to avoid precipitation of heart failure.

Thiamine hydrochloride injection is also indicated in patients with established thiamine deficiency who cannot take thiamine orally due to coexisting severe anorexia, nausea, vomiting, or malabsorption. Thiamine hydrochloride injection is not usually indicated for conditions of decreased oral intake or decreased gastrointestinal absorption, because multiple vitamins should usually be given.

## **CONTRAINDICATIONS**

A history of sensitivity to thiamine or to any of the ingredients in this drug is a contraindication. (See WARNINGS for further information.)

## **WARNINGS**

**WARNING:** This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

Serious hypersensitivity/anaphylactic reactions can occur, especially after repeated administration. Deaths have resulted from IV or IM administration of thiamine (see ADVERSE REACTIONS).

Routine testing for hypersensitivity, in many cases, may not detect hypersensitivity. Nevertheless, a skin test should be performed on patients who are suspected of drug allergies or previous reactions to thiamine, and any positive responders should not receive thiamine by injection.

If hypersensitivity to thiamine is suspected (based on history of drug allergy or

occurrence of adverse reactions after thiamine administration), administer one-hundredth of the dose intradermally and observe for 30 minutes. If no reaction occurs, full dose can be given; the patient should be observed for at least 30 minutes after injection. Be prepared to treat anaphylactic reactions regardless of the precautions taken.

Treatment of anaphylactic reactions includes maintaining a patent airway and the use of epinephrine, oxygen, vasopressors, steroids and antihistamines.

## **PRECAUTIONS**

### General

Simple vitamin B 1 deficiency is rare. Multiple vitamin deficiencies should be suspected in any case of dietary inadequacy.

### Information for Patients

The patient should be advised as to proper dietary habits during treatment so that relapses will be less likely to occur with reduction in dosage or cessation of injection therapy.

### Usage in Pregnancy

Pregnancy Category A—Studies in pregnant women have not shown that thiamine hydrochloride increases the risk of fetal abnormalities if administered during pregnancy. If the drug is used during pregnancy, the possibility of fetal harm appears remote. Because studies cannot rule out the possibility of harm however, thiamine hydrochloride should be used during pregnancy only if clearly needed.

### Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when thiamine hydrochloride is administered to a nursing mother.

## **ADVERSE REACTIONS**

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

An occasional individual may develop a hypersensitivity or life-threatening anaphylactic reaction to thiamine, especially after repeated injections. Collapse and death have been reported. A feeling of warmth, pruritus, urticaria, weakness, sweating, nausea, restlessness, tightness of the throat, angioneurotic edema, cyanosis, pulmonary edema, and hemorrhage into the gastrointestinal tract have also been reported. Some tenderness and induration may follow IM use (see WARNINGS).

## **OVERDOSAGE**

Parenteral doses of 100 to 500 mg singly have been administered without toxic effects. However, dosages exceeding 30 mg three times a day are not utilized effectively.

When the body tissues are saturated with thiamine, it is excreted in the urine as

pyrimidine. As the intake of thiamine is further increased, it appears unchanged in the urine.

## **DOSAGE & ADMINISTRATION**

“Wet” beriberi with myocardial failure must be treated as an emergency cardiac condition, and thiamine must be administered slowly by the IV route in this situation (see WARNINGS).

In the treatment of beriberi, 10 to 20 mg of thiamine hydrochloride are given IM three times daily for as long as two weeks. (See WARNINGS regarding repeated injections of thiamine.) An oral therapeutic multivitamin preparation containing 5 to 10 mg thiamine, administered daily for one month, is recommended to achieve body tissue saturation.

Infantile beriberi that is mild may respond to oral therapy, but if collapse occurs, doses of 25 mg may cautiously be given IV.

Poor dietary habits should be corrected and an abundant and well-balanced dietary intake should be prescribed.

Patients with neuritis of pregnancy in whom vomiting is severe enough to preclude adequate oral therapy should receive 5 to 10 mg of thiamine hydrochloride IM daily.

In the treatment of Wernicke-Korsakoff syndrome, thiamine hydrochloride has been administered IV in an initial dose of 100 mg, followed by IM doses of 50 to 100 mg daily until the patient is consuming a regular, balanced diet. (See WARNINGS regarding repeated injections of thiamine.)

Patients with marginal thiamine status to whom dextrose is being administered should receive 100 mg thiamine hydrochloride in each of the first few liters of IV fluid to avoid precipitating heart failure.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

## **HOW SUPPLIED**

Thiamine Hydrochloride Injection, USP is supplied as follows:

NDC 51662-1523-1, 2 mL in 1 VIAL, MULTI-DOSE, (100 mg/mL)

NDC 51662-1523-2, 2 mL in 1 VIAL, MULTI-DOSE, (100 mg/mL), 1 VIAL PER POUCH

NDC 51662-1523-2, 2 mL in 1 VIAL, MULTI-DOSE, (100 mg/mL), 1 VIAL PER POUCH, 25 POUCHES PER CASE

Store at 20° to 25°C (68° to 77° F) [see USP Controlled Room Temperature].

PROTECT FROM LIGHT.

Use only if solution is clear and seal intact.

## **SPL UNCLASSIFIED**



Lake Zurich, IL 60047

[www.fresenius-kabi.com/us](http://www.fresenius-kabi.com/us)

45819H

Revised: September 2019

**PRINCIPAL DISPLAY PANEL - 51662-1523-1, SERIALIZED VIAL LABELING AND VIAL LABELING**

(01) 00351662152317

(17) 220831

(21) 79744521100

(10) 6023511

See manufacturer's package insert  
Distributed by HF Acquisition Co., LLC  
Mukilteo, WA 98275



RX ONLY



NDC 63323-013-01 1302

**Thiamine HCl**  
Injection, USP

**200 mg per 2 mL**  
(100 mg per mL)

For IM or IV use.  
Protect from light.

2 mL

Multiple Dose Vial

Rx only

NDC 63323-013-02 1302

**THIAMINE HCl**  
**INJECTION, USP**

**200 mg/2 mL**

(100 mg/mL)

For IM or IV Use

Rx only

**2 mL** Multiple Dose Vial

**Sterile**

Each mL contains: Thiamine HCl 100 mg; chlorobutanol anhydrous (chloral derivative) 0.5%; monothioglycerol 0.5%; Water for Injection q.s. Sodium hydroxide may have been added for pH adjustment. Contains no more than 400 mcg/L of aluminum. Usual Dosage: See insert. Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. **PROTECT FROM LIGHT.** Use only if solution is clear and seal intact.



**APP Pharmaceuticals, LLC**  
Schaumburg, IL 60173

42599I



401818H

LOT/EXP



**PRINCIPAL DISPLAY PANEL NDC 51662-1523-2 POUCH LABELING**  
VIAL LABEL



401818H

LOT/EXP



**POUCH LABEL**

THIAMINE HCL INJECTION, USP 200 mg/2 mL  
(100 mg/mL) 2 mL VIAL

NDC: 51662-1523-2  
LOT: 123456  
EXP: 2030-01-01

(01) 00351662152324  
(10) 123456  
(17) 300101  
(21) 118267750034

FOR INTRAMUSCULAR USE OR INTRAVENOUS USE. STERILE. EACH mL  
CONTAINS: THIAMINE HCL 100mg; CHLOROBUTANOL ANHYDROUS  
(CHLORAL DERIVATIVE) 0.5%; MONOTHIOGLYCEROL 0.5%; WATER FOR  
INJECTION, q.s. SODIUM HYDROXIDE MAY HAVE BEEN ADDED FOR pH  
ADJUSTMENT. CONTAINS NO MORE THAN 400 mcg/L OF ALUMINUM. USUAL  
DOSAGE: SEE PACKAGE INSERT. STORE AT 20° TO 25°C (68° TO 77°F)  
[SEE USP CONTROLLED ROOM TEMPERATURE]. PROTECT FROM LIGHT. USE  
ONLY IF SOLUTION IS CLEAR AND SEAL INTACT.

See manufacturer's package insert  
ORIGINAL MFG NDC: 63323-013-02  
RX ONLY  
Manufactured by HF Acquisition Co., LLC  
Mukilteo, WA 98275



## PRINCIPAL DISPLAY PANEL NDC 51662-1523-3 CASE LABELING

### CASE LABELING

THIAMINE HCL INJECTION, USP 200 mg/2 mL  
(100 mg/mL) 2mL VIAL (CASE OF 25)

NDC: 51662-1523-3  
LOT: 123456  
EXP: 2030-01-01

(01) 40351662152339  
(10) 123456  
(17) 300101  
(21) 129066573464

FOR INTRAMUSCULAR USE OR INTRAVENOUS USE. STERILE. EACH mL  
CONTAINS: THIAMINE HCL 100mg; CHLOROBUTANOL ANHYDROUS  
(CHLORAL DERIVATIVE) 0.5%; MONOTHIOGLYCEROL 0.5%; WATER FOR  
INJECTION, q.s. SODIUM HYDROXIDE MAY HAVE BEEN ADDED FOR pH  
ADJUSTMENT. CONTAINS NO MORE THAN 400 mcg/L OF ALUMINUM. USUAL  
DOSAGE: SEE PACKAGE INSERT. STORE AT 20° TO 25°C (68° TO 77°F)  
[SEE USP CONTROLLED ROOM TEMPERATURE]. PROTECT FROM LIGHT. USE  
ONLY IF SOLUTION IS CLEAR AND SEAL INTACT.

See manufacturer's package insert

ORIGINAL MFG NDC: 63323-013-02

RX ONLY

Manufactured by HF Acquisition Co., LLC  
Mukilteo, WA 98275

CASE



SERIALIZED RFID LABELING

(01)40351662152339  
(17)230919  
(21)351662141212  
(10)HF1234TESTING

SEE MANUFACTURE'S INSERT  
DISTRIBUTED BY HF ACQUISITIONS CO., LLC  
MUKILTEO, WA 98275

RX ONLY



## THIAMINE HCL

thiamine hcl injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:51662-1523(NDC:63323- 013)
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>THIAMINE HYDROCHLORIDE</b> (UNII: M572600E5P) (THIAMINE ION - UNII: 4ABT0J945J)	THIAMINE HYDROCHLORIDE	100 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>CHLOROBUTANOL</b> (UNII: HM4YQM8WRC)	
<b>MONOTHIOGLYCEROL</b> (UNII: AAO1P0WSXJ)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51662-1523-1	2 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product	06/14/2021	
2	NDC:51662-1523-3	25 in 1 CASE	11/10/2022	
2	NDC:51662-1523-2	1 in 1 POUCH		
2		2 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA080556	06/14/2021	

**Labeler** - HF Acquisition Co LLC, DBA HealthFirst (045657305)

**Registrant** - HF Acquisition Co LLC, DBA HealthFirst (045657305)

## Establishment

Name	Address	ID/FEI	Business Operations
HF Acquisition Co LLC, DBA HealthFirst		045657305	relabel(51662-1523)

Revised: 1/2024

HF Acquisition Co LLC, DBA HealthFirst