# SUNMARK URINARY PAIN RELIEF MAXIMUM STRENGTH - phenazopyridine hydrochloride tablet

McKesson

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

#### **Drug Facts**

PHENAZOPYRIDINE HYDROCHLORIDE 97.5 MG

#### **PURPOSE**

PHENAZOPYRIDINE HYDROCHLORIDE

**URINARY ANALGESIC** 

#### WARNINGS

DO NOT EXCEED RECOMMENDED DOSAGE

#### **ASK DOCTOR**

ASK DOCTOR BEFORE USE

IF YOU HAVE KIDNEY DISEASE

ALLERGIES TO FOODS, PRESERVATIVES OR DYES

HAD A HYPERSENSITIVE REACTION TO PHENAZOPYRIDINE

#### WHEN USING

WHEN USING THIS PRODUCT

STOMACH UPSET MAY OCCUR, TAKING THIS PRODUCT WITH OR AFTER MEALS MAY REDUCE STOMACH UPSET

YOUR URINE WILL BECOME REDDISH ORANGE IN COLOR.THIS IS NOT HARMFUL, BUT CARE SHOULD BE TAKEN TO

AVOID STAINING CLOTHING OR OTHER ITEMS.

#### **STOP USE**

STOP USE AND ASK A DOCTOR

IF YOUR SYMPTOMS LAST FOR MORE THAN 2 DAYS

YOU SUSPECT YOU ARE HAVING AN ADVERSE REACTION TO THE MEDICATION

#### PREGNANCY OR BREAST FEEDING

ASK A HEALTH PROFESSIONAL BEFORE US

#### KEEP OUT OF REACH OF CHILDREN

IN CASE OF OVERDOSE ,GET MEDICAL HELP OR CONTACT A POISON CONTROL CENTER RIGHT AWAY.

#### **INDICATIONS & USAGE**

Use; fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections.

#### **INACTIVE INGREDIENT**

Lactose, magnesium silicate, magnesium stearate, microcrystalline cellulose,


pharmaceutical glaze, and sodium starch glycolate.

#### **DOSAGE & ADMINISTRATION**

Adults and Children 12 years of age and over; take 2 tablets 3 times daily with a full glass of water, with or after meals as needed.

Children under 12 years of age; consult a doctor.

Do not use for more than 2 days (12 tablets) without consulting a Doctor.



#### SUNMARK URINARY PAIN RELIEF MAXIMUM STRENGTH

phenazopyridine hydrochloride tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-941	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
<b>PHENAZO PYRIDINE HYDRO CHLO RIDE</b> (UNII: 0 EWG668 W17) (PHENAZO PYRIDINE - UNII: K2J09 EMJ52)	PHENAZOPYRIDINE HYDROCHLORIDE	97.5 mg		

Inactive Ingredients			
Ingredient Name	Strength		
LACTOSE (UNII: J2B2A4N98G)			
MAGNESIUM SILICATE (UNII: 9B9691B2N9)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			

Product Characteristics				
Color	red	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	975	
Contains				

l	Packaging				
	# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
	1 NDC:49348-941-01	1 in 1 CARTON	11/18/2010		
	1 NDC:49348-941-02	12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		11/18/2010		

## Labeler - McKesson (116956644)

### **Registrant** - Reese Pharmaceutical Co (004172052)

Establishment			
Name	Address	ID/FEI	Business Operations
Reese Pharmaceutical Co		004172052	relabel(49348-941), repack(49348-941)

Establishment				
Name	Address	ID/FEI	Business Operations	
Pharbest		557054835	manufacture(49348-941)	

Revised: 12/2020 McKesson