

CVS PHARMACY CHILDRENS GLYCERIN SUPPOSITORIES LAXATIVE- glycerin liquid
CVS PHARMACY, INC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredient (in each 2.7 - mL average delivered dose)

Glycerin 2.8 g.....Hyperosmotic Laxative

Use

- or relief of occasional constipation
- this product generally produces bowel movement in 15 minutes to 1 hour

Warnings

For rectal use only

May cause rectal discomfort or a burning sensation

Ask a doctor before using any laxative if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop using this product and consult a doctor if you have

- rectal bleeding
- no bowel movement after use of a laxative

These symptoms may indicate a serious condition.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

Single daily dosage

children 2 to under 6 years	1 suppository or as directed by a doctor
children under 2 years	consult a doctor

Left-side position. Place child on left side with knees bent and arms resting comfortably.

Knee-chest position. Have child kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.

Administering the laxative: CAUTION: REMOVE GREEN PROTECTIVE COVER BEFORE INSERTING. Hold unit upright, grasping bulb of unit with fingers. Grasp green protective cover with other hand, twist one revolution and pull green cover off and discard. With steady pressure, gently

insert tip into rectum with a slight side-to-side movement, with tip pointing toward navel.
Discontinue use if resistance is encountered. Forcing the tip can result in injury. Insertion may be easier if child receiving suppository bears down as if having a bowel movement. This helps relax the muscles around the anus. Squeeze the bulb until nearly all liquid is expelled. While continuing to squeeze the bulb, remove the tip from the rectum and discard unit. It is not necessary to empty unit completely. The unit contains more than the amount of liquid needed for effective use. A small amount of liquid will remain in the unit after squeezing.

Other Information

- Tamper Evident: If the bottle is open do not use

Inactive Ingredients

edetate disodium, purified water

Questions or comments?

Call 1-877-545-3635



CVS PHARMACY CHILDRENS GLYCERIN SUPPOSITORIES LAXATIVE

glycerin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-006	
Route of Administration	RECTAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
Glycerin (UNII: PDC6A3C0OX) (Glycerin - UNII:PDC6A3C0OX)		Glycerin	2.8 g	
Inactive Ingredients				
Ingredient Name			Strength	
EDETATE DISODIUM (UNII: 7FLD91C86K)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-006-06	4 in 1 CARTON		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part334	06/03/2002		

Labeler - CVS PHARMACY, INC. (062312574)

Revised: 2/2012

CVS PHARMACY, INC.