

**SEVERE COLD AND COUGH NIGHTTIME- acetaminophen, diphenhydramine hcl, phenylephrine hcl liquid**  
**Family Dollar (FAMILY WELLNESS)**

-----  
**Drug Facts**

**Active ingredients (in each 30 mL)**

**Acetaminophen 650 mg**

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

**Purposes**

**Pain reliever/fever reducer**

Antihistamine/Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to a cold
  - minor aches and pains
  - headache
  - sore throat
  - runny nose
  - sneezing
  - itchy, watery eyes due to hay fever
  - nasal and sinus congestion
  - itching of the nose or throat
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing diphenhydramine, even one used on skin
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- heart disease
- thyroid disease
- high blood pressure
- diabetes
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking**

- the blood thinning drug warfarin
- sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcohol drinks
- be careful when driving a motor vehicle or operating machinery
- marked drowsiness may occur
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

- **do not take more than directed (see overdose warning)**
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- mL=milliliter
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL every 4 hours
- children under 12 years of age; do not use

## **Other information**

- each 30 mL contains: potassium 10 mg
- each 30 mL contains: sodium 14 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## **Inactive ingredients**

acesulfame potassium, alcohol, citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavors, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

## **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

COMPARE TO THE ACTIVE INGREDIENTS OF THERA® EXPRESSMAX® NIGHTTIME SEVERE COLD AND COUGH\*

Severe Cold & Cough

NIGHTTIME

**Acetaminophen** (Pain Reliever/Fever Reducer)

Diphenhydramine HCl (Cough Suppressant/Antihistamine)

Phenylephrine HCl (Nasal Decongestant)

For Relief of:

- Nasal Congestion
- Headache
- Sore Throat Pain
- Cough
- Body Ache
- Fever
- Runny Nose

Alcohol Content: 10%

MAXIMUM STRENGTH

PER 4 HOUR DOSE

BERRY FLAVOR

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by GSK Consumer Health, distributor Theraflu® ExpressMax® Nighttime Severe Cold & Cough.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC

500 VOLVO PARKWAY, CHESAPEAKE, VA 23320

**Product Label**

COMPARE TO THE ACTIVE INGREDIENTS OF THERAFLU® EXPRESSMAX® NIGHTTIME SEVERE COLD & COUGH\*

**FAMILY wellness™**

NDC 55319-602-83

**Severe Cold & Cough**

**NIGHTTIME**

**Acetaminophen** (Pain Reliever/Fever Reducer)  
**Diphenhydramine HCl** (Cough Suppressant/Antihistamine)  
**Phenylephrine HCl** (Nasal Decongestant)

**For Relief of:**

- Nasal Congestion • Headache
- Sore Throat Pain • Cough
- Body Ache • Fever • Runny Nose

**BERRY FLAVOR**

Alcohol Content: 10%  
 MAXIMUM STRENGTH  
 PER 4 HOUR DOSE

**8.3 FL OZ (245 mL)**

PLD-8399B LB007952

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® ExpressMax® Nighttime Severe Cold & Cough.

**DISTRIBUTED BY: MIDWOOD BRANDS, LLC**  
 500 VOLVO PARKWAY, CHESAPEAKE, VA 23320

**NOT 100% SATISFIED?**

Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

SKU 0913244

PLD-A399B  
 LB007953



**Drug Facts**

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/ Cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Uses**

- temporarily relieves these symptoms due to a cold
  - minor aches and pains
  - headache
  - sore throat
  - runny nose
  - sneezing
  - itchy, watery eyes due to hay fever
  - nasal and sinus congestion
  - itching of the nose or throat
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Drug Facts (continued)**

**Do not use**

- to make a child sleep
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- high blood pressure
- heart disease
- diabetes
- thyroid disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking**

- the blood thinning drug warfarin
- sedatives or tranquilizers

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**When using this product**

- do not exceed recommended dosage
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- marked drowsiness may occur
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses (180 mL) in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
  - 30 mL every 4 hours
- children under 12 years of age: do not use

**Other information**

- each 30 mL contains: potassium 10 mg
- each 30 mL contains: sodium 14 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** acesulfame potassium, alcohol, citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, sodium benzoate, sodium citrate

**Questions or comments?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**FAMILY WELLNESS Severe Cold & Cough Berry Flavor**

**SEVERE COLD AND COUGH NIGHTTIME**

acetaminophen, diphenhydramine hcl, phenylephrine hcl liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55319-602
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MALTITOL</b> (UNII: D65DG142WK)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-602-83	245 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/30/2020	07/31/2025

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2020	07/31/2025

**Labeler** - Family Dollar (FAMILY WELLNESS) (024472631)

